This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

+ *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.

+ *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.

+ *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.

+ *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at `http://books.google.com/`

Educ T 128.73.314

THE NATIONAL SERIES OF STANDARD SCHOOL BOOKS

COMPRISES STANDARD WORKS

In every department of instruction and of every grade. The teacher in want of a book for any particular purpose or class, will always find the best of its kind in our catalogue. No other series even claims to be as complete as this. None is so extensive or so judiciously selected. Among so many volumes a high standard of merit is maintained, as it is our aim never to permit our imprint upon a poor or unworthy book. It is also our plan to make books not for a class or sect, but for the whole country—unobjectionable to parties and creeds, while inculcating the great principles of political freedom and Christianity, upon which all right-minded persons are agreed. Hence, and from their almost universal circulation, the name—"National Series." Among the principal volumes are

Pa … self. *The*
… ry want of
… in smaller
… companion

Da … *rveying,*
… have been
… hed to take

Ba … s to follow.
… forgotten,
… ping about

M … *ysical.*
… ries. From
… he wants.

St … *losophy,*
… l precedent;

Cl … lessons and

W … *French,*
… with signal

Se … *rations,*
… ant editions.

E … ts published

E … ed Authors,
… guage—De-
… on—North-

H … —Berard's
… cal Hist.

P … l Steel Pens
… llen's Map

N … y—Porter's
… 's Botany—

I … y—Hunting-
ton's Fine Arts—Champlin's Political Economy—Mansfield's Government Manual—Alden's Ethics—Brooks' Manual of Devotion—Tracy's School Record, &c.

The Teacher's Library consists of over 30 volumes of strictly professional literature, as Page's Theory and Practice—Holbrook's Normal Methods—Northend's Teacher's Assistant, &c.

A DESCRIPTIVE CATALOGUE of all these and many more may be obtained by enclosing a stamp to the Publishers,

A. S. BARNES & COMPANY,
National Educational Publishers,
… AM STREET, NEW YORK.

3 2044 097 010 888

Harvard College Library

FROM

....

....

....

THE WORMAN SERIES IN MODERN LANGUAGE.

A COMPLETE COURSE IN GERMAN.

By JAMES H. WORMAN, A.M.

EMBRACING

ELEMENTARY GERMAN GRAMMAR,
COMPLETE GERMAN GRAMMAR,
GERMAN READER,
GERMAN COPY-BOOKS, GERMAN ECHO.

IN PREPARATION,

HISTORY OF GERMAN LITERATURE,
GERMAN AND ENGLISH LEXICON.

I. THE GERMAN GRAMMARS of Worman are widely preferred on account of their clear, explicit method (on the conversation plan), introducing a system of analogy and comparison with the learners' own language and others commonly studied.

The arts of speaking, of understanding the spoken language, and of correct pronunciation, are treated with great success.

The new classifications of nouns and of irregular verbs are of great value to the pupil. The use of heavy type to indicate etymological changes, is new. The Vocabulary is *synonymical*—also a new feature.

II. WORMAN'S GERMAN READER contains progressive selections from a wide range of the very best German authors, including three complete plays, which are usually purchased in separate form for advanced students who have completed the ordinary Reader.

It has Biographies of eminent authors, Notes after the text, References to all German Grammars in common use, and an adequate Vocabulary; also, Exercises for translation into the German.

III. WORMAN'S GERMAN ECHO (*Deutsches Echo*) is entirely a new thing in this country. It presents familiar colloquial exercises without translation, and will teach fluent conversation in a few months of diligent study.

No other method will ever make the student *at home* in a foreign language. By this he thinks in, as well as speaks it. For the time being he is *a German* through and through. The laborious process of translating his thoughts no longer impedes free unembarrassed utterance.

WORMAN'S COMPLETE FRENCH COURSE

IS INAUGURATED BY

L'ÉCHO DE PARIS,

Or, "French Echo;" on a plan identical with the German Echo described above. This will be followed in due course by the other volumes of

THE FRENCH SERIES,

VIZ.:

A COMPLETE GRAMMAR, | *A FRENCH READER,*
AN ELEMENTARY GRAMMAR, | *A FRENCH LEXICON,*
A HISTORY OF FRENCH LITERATURE.

WORMAN'S WORKS

are adopted as fast as published by many of the best institutions of the country. In completeness, adaptation, and homogeneity for consistent courses of instruction, they are simply

UNRIVALED.

NEW

ELEMENTARY ALGEBRA:

EMBRACING

THE FIRST PRINCIPLES OF THE SCIENCE.

BY

CHARLES DAVIES, LL.D.,
PROFESSOR OF HIGHER MATHEMATICS, COLUMBIA COLLEGE

A. S. BARNES & COMPANY,
NEW YORK AND CHICAGO.

1873.

Educ T 128.73.314

DAVIES' MATHEMATICS.

THE WEST POINT COURSE,

And Only Thorough and Complete Mathematical Series.

IN THREE PARTS.

I. COMMON SCHOOL COURSE.

Davies' Primary Arithmetic.—The fundamental principles displayed in Object Lessons.

Davies' Intellectual Arithmetic.—Referring all operations to the unit 1 as the only tangible basis for logical development.

Davies' Elements of Written Arithmetic.—A practical introduction to the whole subject. Theory subordinated to Practice.

Davies' Practical Arithmetic.*—The most successful combination of Theory and Practice, clear, exact, brief, and comprehensive.

II. ACADEMIC COURSE.

Davies' University Arithmetic.*—Treating the subject exhaustively as *a science*, in a logical series of connected propositions.

Davies' Elementary Algebra.*—A connecting link, conducting the pupil easily from arithmetical processes to abstract analysis.

Davies' University Algebra.*—For institutions desiring a more complete but not the fullest course in pure Algebra.

Davies' Practical Mathematics.—The science practically applied to the useful arts, as Drawing, Architecture, Surveying, Mechanics, etc.

Davies' Elementary Geometry.—The important principles in simple form, but with all the exactness of vigorous reasoning.

Davies' Elements of Surveying.—Re-written in 1870. The simplest and most practical presentation for youths of 12 to 16.

III. COLLEGIATE COURSE.

Davies' Bourdon's Algebra.*—Embracing Sturm's Theorem, and a most exhaustive and scholarly course.

Davies' University Algebra.*—A shorter course than Bourdon, for Institutions have less time to give the subject.

Davies' Legendre's Geometry.—Acknowledged *the only* satisfactory treatise of its grade. 300,000 copies have been sold.

Davies' Analytical Geometry and Calculus.—The shorter treatises, combined in one volume, are more available for American courses of study.

Davies' Analytical Geometry. } The original compendiums, for those de-
Davies' Diff. & Int. Calculus. } siring to give full time to each branch.

Davies' Descriptive Geometry.—With application to Spherical Trigonometry, Spherical Projections, and Warped Surfaces.

Davies' Shades, Shadows, and Perspective.—A succinct exposition of the mathematical principles involved.

Davies' Science of Mathematics.—For teachers, embracing

I. GRAMMAR OF ARITHMETIC,
II. OUTLINES OF MATHEMATICS,
III. LOGIC AND UTILITY OF MATHEMATICS,
IV. MATHEMATICAL DICTIONARY.

* Keys may be obtained from the Publishers by Teachers only.

Entered, according to Act of Congress, in the year 1859, by
CHARLES DAVIES,
In the Clerk's Office of the District Court of the United States for the Southern District of New York.

N. E. A.

HARVARD COLLEGE LIBRARY
GIFT OF THE
AMERICAN ANTIQUARIAN SOCIETY
Sep. 8, 1938

PREFACE.

ALGEBRA naturally follows Arithmetic in a course of scientific studies. The language of figures, and the elementary combinations of numbers, are acquired at an early age. When the pupil passes to a new system, conducted by letters and signs, the change seems abrupt; and he often experiences much difficulty before perceiving that Algebra is but Arithmetic written in a different language.

It is the design of this work to supply a connecting link between Arithmetic and Algebra; to indicate the unity of the methods, and to conduct the pupil from the arithmetical processes to the more abstract methods of analysis, by easy and simple gradations. The work is also introductory to the University Algebra, and to the Algebra of M. Bourdon, which is justly considered, both in this country and in Europe, as the best text-book on the subject, which has yet appeared.

In the Introduction, or Mental Exercises, the language of figures and letters are both employed. Each Lesson is so arranged as to introduce a single principle, not known before, and the whole is so combined as to prepare the pupil, by a thorough system of mental training, for those processes of reasoning which are peculiar to the algebraic analysis.

It is about twenty years since the first publication of the ELEMENTARY ALGEBRA. Within that time, great changes have taken place in the schools of the country. The systems of mathematical instruction have been improved, new methods have been developed, and these require corresponding modifications in the text-books. Those modifications have now been made, and this work will be permanent in its present form.

Many changes have been made in the present edition, at the suggestion of teachers who have used the work, and favored me with their opinions, both of its defects and merits. I take this opportunity of thanking them for the valuable aid they have rendered me. The criticisms of those engaged in the daily business of teaching are invaluable to an author; and I shall feel myself under special obligation to all who will be at the trouble to communicate to me, at any time, such changes, either in methods or language, as their experience may point out. It is only through the cordial co-operation of teachers and authors—by joint labors and mutual efforts—that the text-books of the country can be brought to any reasonable degree of perfection.

A Key to this volume has been prepared for the use of *Teachers only.*

CONTENTS.

CHAPTER I.

DEFINITIONS AND EXPLANATORY SIGNS.

PAGES

Algebra—Definitions—Explanation of the Signs 33–41
Examples in writing Algebraic expressions 41
Interpretation of Algebraic language 42

CHAPTER II.

FUNDAMENTAL OPERATIONS.

Addition—Rule—Examples 43–50
Subtraction—Rule—Examples—Remarks 50–56
Multiplication—Monomials—Polynomials 56–63
Division—Monomials.. 63–68
Signification of the symbol a^0 68–70
Division of Polynomials—Examples 71–75

CHAPTER III.

USEFUL FORMULAS. FACTORING, ETC.

Formulas (**1**), (**2**), (**3**), (**4**), (**5**), and (**6**) 76–79
Factoring 79–81
Greatest Common Divisor................................. 81–84
Least Common Multiple 84–87

CHAPTER IV

FRACTIONS.

Transformation of Fractions 89
To Reduce an Entire Quantity to a Fractional Form 90

PAGES.
To Reduce a Fraction to its Lowest Terms 90
To Reduce a Fraction to a Mixed Quantity.................... 92
To Reduce a Mixed Quantity to a Fraction 93
To Reduce Fractions to a Common Denominator 94
Addition of Fractions 96-98
Subtraction of Fractions 98-99
Multiplication of Fractions 99-102
Division of Fractions....................................... 102-105

CHAPTER V.

EQUATIONS OF THE FIRST DEGREE.

Definition of an Equation—Different Kinds 105-106
Transformation of Equations—First and Second 106-110
Solution of Equations—Rule 110-114
Problems involving Equations of the First Degree 115-130
Equations involving Two Unknown Quantities 130-131
Elimination—By Addition—By Subtraction—By Comparison.. 131-143
Problems involving Two Unknown Quantities 143-148
Equations involving Three or more Unknown Quantities...... 148-159

CHAPTER VI.

FORMATION OF POWERS.

Definition of Powers... 160-161
Powers of Monomials 161-163
Powers of Fractions 163-165
Powers of Binomials 165-167
Of the Terms—Exponents—Coefficients 167-170
Binomial Formula—Examples 170-172

CHAPTER VII.

SQUARE ROOT. RADICALS OF THE SECOND DEGREE.

Definition—Perfect Squares—Rule—Examples............... 173-179
Square Root of Fractions.................................... 179-181
Square Root of Monomials.................................... 181-183
Imperfect Squares, or Radicals 183-187
Addition of Radicals... 187-189
Subtraction of Radicals 189-190

PAGES.
Multiplication of Radicals 190–191
Division of Radicals 191–192
Square Root of Polynomials 193–197

CHAPTER VIII.

EQUATIONS OF THE SECOND DEGREE.

Equations of the Second Degree—Definition—Form 198–200
Incomplete Equations 200–209
Complete Equations—Rule 209–211
Four Forms 211–227
Four Properties 227–229
Formation of Equations of the Second Degree 229–231
Numerical Values of the Roots 231–236
Problems 236–240
Equations involving more than One Unknown Quantity 241–250
Problems 250–254

CHAPTER IX.

ARITHMETICAL AND GEOMETRICAL PROPORTION.

Ways in which Two Quantities may be Compared 255
Arithmetical Proportion and Progression 256–257
Last Term 257–260
Sum of the Extremes—Sum of Series 260–262
The Five Numbers—To find any number of Means 262–265
Geometrical Proportion 267
Various Kinds of Proportion 268–278
Geometrical Progression 278–280
Last Term—Sum of Series 280–285
Progression having an Infinite Number of Terms 285–288
The Five Numbers—To find One Mean 288–289

CHAPTER X.

LOGARITHMS.

Theory of Logarithms 290–295

SUGGESTIONS TO TEACHERS.

1. THE Introduction is designed as a mental exercise. If thoroughly taught, it will train and prepare the mind of the pupil for those higher processes of reasoning, which it is the peculiar province of the algebraic analysis to develop.

2. The statement of each question should be made, and every step in the solution gone through with, without the aid of a slate or black-board; though perhaps, in the beginning, some aid may be necessary to those unaccustomed to such exercises.

3. Great care must be taken to have every principle on which the statement depends, carefully analyzed; and equal care is necessary to have every step in the solution distinctly explained.

4. The reasoning process is the logical connection of distinct apprehensions, and the deduction of the consequences which follow from such a connection. Hence, the basis of all reasoning must lie in distinct elementary ideas.

5. Therefore, to teach one thing at a time—to teach that thing well—to explain its connections with other things, and the consequences which follow from such connections, would seem to embrace the whole art of instruction.

ELEMENTARY ALGEBRA.

INTRODUCTION.

MENTAL EXERCISES.

LESSON I.

1. John and Charles have the same number of apples; both together have twelve: how many has each?

Analysis.—Let x denote the number which John has; then, since they have an equal number, x will also denote the number which Charles has, and twice x, or $2x$, will denote the number which both have, which is 12. If twice x is equal to 12, x will be equal to 12 divided by 2, which is 6; therefore, each has 6 apples.

WRITTEN.

Let x denote the number of apples which John has; then,

$$x + x = 2x = 12; \text{ hence, } x = \frac{12}{2} = 6.$$

Note.—When x is written with the sign $+$ before it, it is read *plus* x: and the line above, is read, x *plus* x *equals* 12.

NOTE.—When x is written by itself, it is read one x, and is the same as, $1x$;

x or $1x$,	means once	x,	or one x,
$2x$,	" twice	x,	or two x,
$3x$,	" three times	x,	or three x,
$4x$,	" four times	x,	or four x,
&c.,	&c.,		&c.

2. What is $x + x$ equal to?
3. What is $x + 2x$ equal to?
4. What is $x + 2x + x$ equal to?
5. What is $x + 5x + x$ equal to?
6. What is $x + 2x + 3x$ equal to?

7. James and John together have twenty-four peaches, and one has as many as the other: how many has each?

ANALYSIS.—Let x denote the number which James has; then, since they have an equal number, x will also denote the number which John has, and twice x will denote the number which both have, which is 24. If twice x is equal to 24, x will be equal to 24 divided by 2, which is 12; therefore, each has 12 peaches.

WRITTEN.

Let x denote the number of peaches which James has; then,

$$x + x = 2x = 24; \text{ hence, } x = \frac{24}{2} = 12.$$

VERIFICATION.

A Verification is the operation of proving that the number found will satisfy the conditions of the question. Thus,

$$\overset{\text{James' apples.}}{12} + \overset{\text{John's apples.}}{12} = 24.$$

NOTE.—Let the following questions be *analyzed*, *written*, and *verified*, in *exactly* the *same manner* as the above.

8. William and John together have 36 pears, and one has as many as the other: how many has each?

9. What number added to itself will make 20?

10. James and John are of the same age, and the sum of their ages is 32: what is the age of each?

11. Lucy and Ann are twins, and the sum of their ages is 16: what is the age of each?

12. What number is that which added to itself will make 30?

13. What number is that which added to itself will make 50?

14. Each of two boys received an equal sum of money at Christmas, and together they received 60 cents: how much had each?

15. What number added to itself will make 100?

16. John has as many pears as William; together they have 72: how many has each?

17. What number added to itself will give a sum equal to 46?

18. Lucy and Ann have each a rose bush with the same number of buds on each; the buds on both number 46: how many on each?

LESSON II.

1. John and Charles together have 12 apples, and Charles has twice as many as John: how many has each?

ANALYSIS.—Let x denote the number of apples which John has; then, since Charles has twice as many, $2x$ will denote his share, and $x + 2x$, or $3x$, will denote the number which they both have, which is 12. If $3x$ is equal to 12, x will be equal to 12 divided by 3, which is 4; therefore, John has 4 apples, and Charles, having twice as many, has 8.

WRITTEN.

Let x denote the number of apples John has, then,

$2x$ will denote the number of apples Charles has; and

$x + 2x = 3x = 12$, the number both have; then,

$x = \frac{12}{3} = 4$, the number John has; and,

$2x = 2 \times 4 = 8$, the number Charles has.

VERIFICATION.

$4 + 8 = 12$, the number both have.

2. William and John together have 48 quills, and William has twice as many as John: how many has each?

3. What number is that which added to twice itself, will give a number equal to 60?

4. Charles' marbles added to John's make 3 times as many as Charles has; together they have 51: how many has each?

ANALYSIS.—Since Charles' marbles added to John's make three times as many as Charles has, Charles must have one third, and John two thirds of the whole.

Let x denote the number which Charles has; then $2x$ will denote the number which John has, and $x + 2x$, or $3x$, will denote what they both have, which is 51. Then, if $3x$ is equal to 51, x will be equal to 51 divided by 3, which is 17. Therefore, Charles has 17 marbles, and John, having twice as many, has 34.

WRITTEN.

Let x denote the number of Charles' marbles; then,

$2x$ will denote the number of John's marbles; and

$3x = 51$, the number of both; then,

$x = \frac{51}{3} = 17$, Charles' marbles; and

$17 \times 2 = 34$, John's marbles.

5. What number added to twice itself will make 75?

6. What number added to twice itself will make 57?

7. What number added to twice itself will make 39?

8. What number added to twice itself will give 90?

9. John walks a certain distance on Tuesday, twice as far on Wednesday, and in the two days he walks 27 miles: how far did he walk each day?

10. Jane's bush has twice as many roses as Nancy's: and on both bushes there are 36: how many on each?

11. Samuel and James bought a ball for 48 cents; Samuel paid twice as much as James: what did each pay?

12. Divide 48 into two such parts that one shall be double the other.

13. Divide 66 into two such parts that one shall be double the other.

14. The sum of three equal numbers is 12: what are the numbers?

ANALYSIS.—Let x denote one of the numbers; then, since the numbers are equal, x will also denote each of the others, and x plus x plus x, or $3x$ will denote their sum, which is 12. Then, if $3x$ is equal to 12, x will be equal to 12 divided by 3, which is 4: therefore, the numbers are 4, 4, and 4.

WRITTEN.

Let x denote one of the equal numbers; then,

$$x + x + x = 3x = 12; \text{ and}$$

$$x = \frac{12}{3} = 4.$$

VERIFICATION.

$$4 + 4 + 4 = 12.$$

15. The sum of three equal numbers is 24: what are the numbers?

16. The sum of three equal numbers is 36: what are the numbers?

17. The sum of three equal numbers is 54: what are the numbers?

LESSON III.

1. What number is that which added to three times itself will make 48?

ANALYSIS.—Let x denote the number; then, $3x$ will denote three times the number, and x plus $3x$, or $4x$, will denote the sum, which is 48. If $4x$ is equal to 48, x will be equal to 48 divided by 4, which is 12; therefore, 12 is the required number.

WRITTEN.

Let x denote the number; then,

$3x =$ three times the number; and

$x + 3x = 4x = 48$, the sum: then,

$x = \frac{48}{4} = 12$, the required number.

VERIFICATION.

$$12 + 3 \times 12 = 12 + 36 = 48.$$

NOTE.—All *similar* questions are solved by the same form of analysis.

2. What number added to 4 times itself will give 40?
3. What number added to 5 times itself will give 42?
4. What number added to 6 times itself will give 63?
5. What number added to 7 times itself will give 88?
6. What number added to 8 times itself will give 81?
7. What number added to 9 times itself will give 100?
8. James and John together have 24 quills, and John has three times as many as James: how many has each?
9. William and Charles have 64 marbles, and Charles has 7 times as many as William: how many has each?

10. James and John travel 96 miles, and James travels 11 times as far as John: how far does each travel?

11. The sum of the ages of a father and son is 84 years; and the father is 3 times as old as the son: what is the age of each?

12. There are two numbers of which the greater is 7 times the less, and their sum is 72: what are the numbers?

13. The sum of four equal numbers is 64: what are the numbers?

14. The sum of six equal numbers is 54: what are the numbers?

15. James has 24 marbles; he loses a certain number, and then gives away 7 times as many as he loses which takes all he has: how many did he give away? Verify.

16. William has 36 cents, and divides them between his two brothers, James and Charles, giving one, eight times as many as the other: how many does he give to each?

17. What is the sum of x and $3x$? Of x and $7x$? Of x and $5x$? Of x and $12x$?

LESSON IV.

1. If 1 apple costs 1 cent, what will a number of apples denoted by x cost?

Analysis.—Since one apple costs 1 cent, and since x denotes *any number* of apples, the cost of x apples will be as many cents as there are apples: that is, x cents.

2. If 1 apple costs 2 cents, what will x apples cost?

Analysis.—Since one apple costs 2 cents, and since x denotes the number of apples, the cost will be twice as many cents as there are apples: that is $2x$ cents.

3. If 1 apple costs 3 cents, what will x apples cost?

4. If 1 lemon costs 4 cents, what will x lemons cost?

5. If 1 orange costs 6 cents, what will x oranges cost?

6. Charles bought a certain number of lemons at 2 cents apiece, and as many oranges at 3 cents apiece, and paid in all 20 cents: how many did he buy of each?

ANALYSIS.—Let x denote the number of lemons; then, since he bought as many oranges as lemons, it will also denote the number of oranges. Since the lemons were 2 cents apiece, $2x$ will denote the cost of the lemons; and since the oranges were 3 cents apiece, $3x$ will denote the cost of the oranges; and $2x + 3x$, or $5x$, will denote the cost of both, which is 20 cents. Now, since $5x$ cents are equal to 20 cents, x will be equal to 20 cents divided by 5 cents, which is 4: hence, he bought 4 of each.

WRITTEN.

Let x denote the number of lemons, or oranges; then,

$2x =$ the cost of the lemons; and

$3x =$ the cost of the oranges; hence,

$2x + 3x = 5x = 20$ cents $=$ the cost of lemons and oranges; hence,

$$x = \frac{20 \text{ cents}}{5 \text{ cents}} = 4, \text{ the number of each.}$$

VERIFICATION.

4 lemons at 2 cents each, give, $4 \times 2 = 8$ cents.
4 oranges at 3 cents each, " $4 \times 3 = 12$ cents.
Hence, they both cost, 8 cents $+12$ cents $= 20$ cents.

7. A farmer bought a certain number of sheep at 4 dollars apiece, and an equal number of lambs at 1 dollar apiece, and the whole cost 60 dollars: how many did he buy of each?

8. Charles bought a certain number of apples at 1 cent apiece, and an equal number of oranges at 4 cents apiece, and paid 50 cents in all: how many did he buy of each?

9. James bought an equal number of apples, pears, and lemons; he paid 1 cent apiece for the apples, 2 cents apiece for the pears, and 3 cents apiece for the lemons, and paid 72 cents in all: how many did he buy of each? Verify.

10. A farmer bought an equal number of sheep, hogs, and calves, for which he paid 108 dollars; he paid 3 dollars apiece for the sheep, 5 dollars apiece for the hogs, and 4 dollars apiece for the calves: how many did he buy of each?

11. A farmer sold an equal number of ducks, geese, and turkeys, for which he received 90 shillings. The ducks brought him 3 shillings apiece, the geese 5, and the turkeys 7: how many did he sell of each sort?

12. A tailor bought, for one hundred dollars, two pieces of cloth, each of which contained an equal number of yards. For one piece he paid 3 dollars a yard, and for the other 2 dollars a yard: how many yards in each piece?

13. The sum of three numbers is 28; the second is twice the first, and the third twice the second: what are the numbers? Verify.

14. The sum of three numbers is 64; the second is 3 times the first, and the third 4 times the second: what are the numbers?

LESSON V.

1. If 1 yard of cloth costs x dollars, what will 2 yards cost?

ANALYSIS.—Two yards of cloth will cost twice as much as one yard. Therefore, if 1 yard of cloth costs x dollars, 2 yards will cost twice x dollars, or $2x$ dollars.

2. If 1 yard of cloth costs x dollars, what will 3 yards cost? Why?

3. If 1 orange costs x cents, what will 7 oranges cost? Why? 8 oranges?

4. Charles bought 3 lemons and 4 oranges, for which he paid 22 cents. He paid twice as much for an orange as for a lemon: what was the price of each?

ANALYSIS.—Let x denote the price of a lemon; then, $2x$ will denote the price of an orange; $3x$ will denote the cost of 3 lemons, and $8x$ the cost of 4 oranges; hence, $3x$ plus $8x$, or $11x$, will denote the cost of the lemons and oranges, which is 22 cents. If $11x$ is equal to 22 cents, x is equal to 22 cents divided by 11, which is 2 cents: therefore, the price of 1 lemon is 2 cents, and that of 1 orange 4 cents.

WRITTEN.

Let x denote the price of 1 lemon; then,

$2x =$ " 1 orange; and,

$3x + 8x = 11x = 22$ cts., the cost of lemons and oranges;

hence, $x = \frac{22 \text{ cts.}}{11} = 2$ cts., the price of 1 lemon;

and, $2 \times 2 = 4$ cts., the price of 1 orange.

VERIFICATION.

$3 \times 2 =$ 6 cents, cost of lemons,
$4 \times 4 =$ 16 cents, cost of oranges.
22 cents, total cost.

5. James bought 8 apples and 3 oranges, for which he paid 20 cents. He paid as much for 1 orange as for 4 apples: what did he pay for one of each?

6. A farmer bought 3 calves and 7 pigs, for which he paid 19 dollars. He paid four times as much for a calf as for a pig: what was the price of each?

7. James bought an apple, a peach, and a pear, for which he paid 6 cents. He paid twice as much for the peach as for

the apple, and three times as much for the pear as for the apple: what was the price of each?

8. William bought an apple, a lemon, and an orange, for which he paid 24 cents. He paid twice as much for the lemon as for the apple, and 3 times as much for the orange as for the apple: what was the price of each?

9. A farmer sold 4 calves and 5 cows, for which he received 120 dollars. He received as much for 1 cow as for 4 calves: what was the price of each?

10. Lucy bought 3 pears and 5 oranges, for which she paid 26 cents, giving twice as much for each orange as for each pear: what was the price of each?

11. Ann bought 2 skeins of silk, 3 pieces of tape, and a penknife, for which she paid 80 cents. She paid the same for the silk as for the tape, and as much for the penknife as for both: what was the cost of each?

12. James, John, and Charles are to divide 56 cents among them, so that John shall have twice as many as James, and Charles twice as many as John: what is the share of each?

13. Put 54 apples into three baskets, so that the second shall contain twice as many as the first, and the third as many as the first and second: how many will there be in each.

14. Divide 60 into four such parts that the second shall be double the first, the third double the second, and the fourth double the third: what are the numbers?

LESSON VI.

1. If $2x + x$ is equal to $3x$, what is $3x - x$ equal to? Written, $3x - x = 2x$.

2. What is $4x - x$ equal to? Written,

$$4x - x = 3x.$$

3. What is $8x$ minus $6x$ equal to? Written,

$$8x - 6x = 2x.$$

4. What is $12x - 9x$ equal to? *Ans.* $3x$.

5. What is $15x - 7x$ equal to?

6. What is $17x - 13x$ equal to? *Ans.* $4x$.

7. Two men, who are 30 miles apart, travel towards each other; one at the rate of 2 miles an hour, and the other at the rate of 3 miles an hour: how long before they will meet?

ANALYSIS:—Let x denote the number of hours. Then, since the time, multiplied by the rate, will give the distance, $2x$ will denote the distance traveled by the first, and $3x$ the distance traveled by the second. But the sum of the distances is 30 miles; hence,

$$2x + 3x = 5x = 30 \text{ miles};$$

and if $5x$ is equal to 30, x is equal to 30 divided by 5, which is 6: hence, they will meet in 6 hours.

WRITTEN.

Let x denote the time in hours; then,

$2x =$ the distance traveled by the 1st; and
$3x =$ " " 2d.

By the conditions,

$$2x + 3x = 5x = 30 \text{ miles, the distance apart};$$

hence, $$x = \frac{30}{5} = 6 \text{ hours}.$$

VERIFICATION.

$$\begin{aligned} 2 \times 6 &= 12 \text{ miles, distance traveled by the first.} \\ 3 \times 6 &= \underline{18 \text{ miles}}\text{, distance traveled by the second.} \\ & 30 \text{ miles, whole distance.} \end{aligned}$$

8. Two persons are 10 miles apart, and are traveling in the same direction; the first at the rate of 3 miles an hour, and the second at the rate of 5 miles: how long, before the second will overtake the first?

ANALYSIS.—Let x denote the time, in hours. Then, $3x$ will denote the distance traveled by the first in x hours; and $5x$ the distance traveled by the second. But when the second overtakes the first, he will have traveled 10 miles more than the first: hence,

$$5x - 3x = 2x = 10;$$

if $2x$ is equal to 10, x is equal to 5: hence, the second will overtake the first in 5 hours.

WRITTEN.

Let x denote the time, in hours: then,

$3x =$ the distance traveled by the 1st;

and, $5x =$ " " 2d;

and, $5x - 3x = 2x = 10$ hours;

or, $x = \frac{10}{2} = 5$ hours.

VERIFICATION.

$3 \times 5 = 15$ miles, distance traveled by 1st.

$5 \times 5 = 25$ miles, " " 2d.

$25 - 15 = 10$ miles, distance apart.

9. A cistern, holding 100 hogsheads, is filled by two pipes; one discharges 8 hogsheads a minute, and the other 12: in what time will they fill the cistern?

10. A cistern, holding 120 hogsheads, is filled by 3 pipes; the first discharges 4 hogsheads in a minute, the second 7, and the third 1: in what time will they fill the cistern?

11. A cistern which holds 90 hogsheads, is filled by a pipe which discharges 10 hogsheads a minute; but there is a waste pipe which loses 4 hogsheads a minute: how long will it take to fill the cistern?

12. Two pieces of cloth contain each an equal number of yards; the first cost 3 dollars a yard, and the second 5, and both pieces cost 96 dollars: how many yards in each?

13. Two pieces of cloth contain each an equal number of yards; the first cost 7 dollars a yard, and the second 5; the first

cost 60 dollars more than the second: how many yards in each piece?

14. John bought an equal number of oranges and lemons the oranges cost him 5 cents apiece, and the lemons 3; and he paid 56 cents for the whole: how many did he buy of each kind?

15. Charles bought an equal number of oranges and lemons; the oranges cost him 5 cents apiece, and the lemons 3; he paid 14 cents more for the oranges than for the lemons: how many did he buy of each?

16. Two men work the same number of days, the one receives 1 dollar a day, and the other two: at the end of the time they receive 54 dollars: how long did they work?

LESSON VII.

1. John and Charles together have 25 cents, and Charles has 5 more than John: how many has each?

ANALYSIS.—Let x denote the number which John has; then, $x + 5$ will denote the number which Charles has, and $x + x + 5$, or $2x + 5$, will be equal to 25, the number they both have. Since $2x + 5$ equals 25, $2x$ will be equal to 25 minus 5, or 20, and x will be equal to 20 divided by 2, or 10: therefore, John has 10 cents, and Charles 15.

WRITTEN.

Let x denote the number of John's cents; then,

$x + 5 =$ " Charles' cents; and,

$x + x + 5 = 25$, the number they both have; or,

$2x + 5 = 25$; and,

$2x = 25 - 5 = 20$; hence,

$x = \frac{20}{2} = 10$, John's number; and,

$10 + 5 = 15$, Charles' number.

VERIFICATION.

John's. Charles'.
10 + 15 = 25, the sum.

Charles'. John's.
15 − 10 = 5, the difference.

2. James and John have 30 marbles, and John has 4 more than James: how many has each?

3. William bought 60 oranges and lemons; there were 20 more lemons than oranges: how many were there of each sort?

4. A farmer has 20 more cows than calves; in all he has 36: how many of each sort?

5. Lucy has 28 pieces of money in her purse, composed of cents and dimes; the cents exceed the dimes in number by 16: how many are there of each sort?

6. What number added to itself, and to 9, will make 29?

7. What number added to twice itself, and to 4, will make 25?

8. What number added to three times itself, and to 12, will make 60?

9. John has five times as many marbles as Charles, and what they both have, added to 14, makes 44: how many has each?

10. There are three numbers, of which the second is twice the first, and the third twice the second, and when 9 is added to the sum, the result is 30: what are the numbers?

11. Divide 17 into two such parts that the second shall be two more than double the first: what are the parts?

12. Divide 40 into three such parts that the second shall be twice the first, and the third exceed six times the first by 4: what are the parts?

13. Charles has twice as many cents as James, and John

has twice as many as Charles; if 7 be added to what they all have, the sum will be 28: how many has each?

14. Divide 15 into three such parts that the second shall be 3 times the first, the third twice the second, and 5 over: what are the numbers?

15. An orchard contains three kinds of trees, apples, pears, and cherries; there are 4 times as many pears as apples, twice as many cherries as pears, and if 14 be added, the number will be 40; how many are there of each?

LESSON VIII.

1. John after giving away 5 marbles, had 12 left: how many had he at first?

ANALYSIS.—Let x denote the number; then, x minus 5 will denote what he had left, which was equal to 12. Since x diminished by 5 is equal to 12, x will be equal to 12, increased by 5; that is, to 17: therefore, he had 17 marbles.

WRITTEN.

Let x denote the number he had at first; then,

$x - 5 = 12$, what he had left; and

$x = 12 + 5 = 17$, what he first had.

VERIFICATION.

$17 - 5 = 12$, what were left.

2. Charles lost 6 marbles and has 9 left: how many had he at first?

3. William gave 15 cents to John, and had 9 left: how many had he at first?

4. Ann plucked 8 buds from her rose bush, and there were 19 left: how many were there at first?

5. William took 27 cents from his purse, and there were 13 left: how many were there at first?

6. The sum of two numbers is 14, and their difference is 2: what are the numbers?

ANALYSIS.—The difference of two numbers, added to the less, will give the greater. Let x denote the *less* number; then, $x + 2$, will denote the greater, and $x + x + 2$, will denote their sum, which is 14. Then, $2x + 2$ equals 14; and $2x$ equals 14 minus 2, or 12: hence, x equals 12 divided by 2, or 6: hence, the numbers are 6 and 8.

VERIFICATION.

$$6 + 8 = 14, \text{ their sum; and}$$
$$8 - 6 = 2, \text{ their difference.}$$

7. The sum of two numbers is 18, and their difference 6: what are the numbers?

8. James and John have 26 marbles, and James has 4 more than John: how many has each?

9. Jane and Lucy have 16 books, and Lucy has 8 more than Jane: how many has each?

10. William bought an equal number of oranges and lemons; Charles took 5 lemons, after which William had but 25 of both sorts: how many did he buy of each?

11. Mary has an equal number of roses on each of two bushes; if she takes 4 from one bush, there will remain 24 on both: how many on each at first?

12. The sum of two numbers is 20, and their difference is 6: what are the numbers?

ANALYSIS.—If x denotes the *greater* number, $x - 6$ will denote the less, and $x + x - 6$ will be equal to 20; hence, $2x$ equals $20 + 6$, or 26, and x equals 26 divided by 2, equals 13; hence the numbers are 13 and 7.

WRITTEN.

Let x denote the greater; then,

$x - 6 =$ the less; and

$x + x - 6 = 20$, their sum; hence,

$$2x = 20 + 6 = 26; \text{ or,}$$

$$x = \frac{26}{2} = 13; \text{ and } 13 - 6 = 7.$$

VERIFICATION.

$$13 + 7 = 20; \text{ and, } 13 - 7 = 6.$$

13. The sum of the ages of a father and son is 60 years, and their difference is just half that number: what are their ages?

14. The sum of two numbers is 23, and the larger lacks 1 of being 7 times the smaller: what are the numbers?

15. The sum of two numbers is 50; the larger is equal to 10 times the less, minus 5: what are the numbers?

16. John has a certain number of oranges, and Charles has four times as many, less seven; together they have 53: how many has each?

17. An orchard contains a certain number of apple trees, and three times as many cherry trees, less 6; the whole number is 30: how many of each sort?

LESSON IX.

1. If x denotes any number, and 1 be added to it, what will denote the sum? *Ans.* $x + 1$.

2. If 2 be added to x, what will denote the sum? If 3 be added, what? If 4 be added? &c.

If to John's marbles, one marble be added, twice his number will be equal to 10: how many had he?

ANALYSIS.—Let x denote the number; then, $x + 1$ will denote the number after 1 is added, and twice this number, or $2x + 2$, will be equal to 10. If $2x + 2$ is equal to 10, $2x$ will be equal to 10 minus 2, or 8; or x will be equal to 4.

WRITTEN.

Let x denote the number of John's marbles; then,

$x + 1 =$ the number, after 1 is added; and

$2(x + 1) = 2x + 2 = 10$; hence,

$$2x = 10 - 2; \text{ or } x = \frac{8}{2} = 4.$$

VERIFICATION.

$$2(4 + 1) = 2 \times 5 = 10.$$

4. Write $x + 2$ multiplied by 3. *Ans.* $3(x + 2)$.
What is the product? *Ans.* $3x+6$.
5. Write $x + 4$ multiplied by 5. *Ans.* $5(x + 4)$.
What is the product? *Ans.* $5x + 20$.
6. Write $x + 3$ multiplied by 4. *Ans.* $4(x + 3)$.
What is the product? *Ans.* $4x + 12$.

7. Lucy has a certain number of books; her father gives her two more, when twice her number is equal to 14: how many has she?

8. Jane has a certain number of roses in blossom; two more bloom, and then 3 times the number is equal to 15: how many were in blossom at first?

9. Jane has a certain number of handkerchiefs, and buys 4 more, when 5 times her number is equal to 45: how many had she at first?

10. John has 1 apple more than Charles, and 3 times John's, added to what Charles has, make 15: how many has each?

ANALYSIS.—Let x denote Charles' apples; then $x + 1$ will denote John's; and $x + 1$ multiplied by 3, added to x, or $3x + 3 + x$, will be equal to 15, what they both had; hence, $4x + 3$ equals 15; and $4x$ equals 15 minus 3, or 12; and $x = 4$. Write, and verify.

11. James has two marbles more than William, and twice his marbles plus twice William's are equal to 16: how many has each?

12. Divide 20 into two such parts that one part shall exceed the other by 4.

13. A fruit-basket contains apples, pears, and peaches; there are 2 more pears than apples, and twice as many peaches as pears; there are 22 in all: how many of each sort?

14. What is the sum of $x + 3x + 2(x + 1)$?

15. What is the sum of $2(x + 1) + 1(x + 1) + x$?

16. What is the sum of $x + 5(x + 8)$?

17. The sum of two numbers is 11, and the second is equal to twice the first plus 2: what are the numbers?

18. John bought 3 apples, 3 lemons, and 3 oranges, for which he paid 27 cents; he paid 1 cent more for a lemon than for an apple, and 1 cent more for an orange than for a lemon: what did he pay for each?

19. Lucy, Mary, and Ann, have 15 cents; Mary has 1 more than Lucy, and Ann twice as many as Mary?

LESSON X.

1. If x denote any number, and 1 be subtracted from it, what will denote the difference? *Ans.* $x - 1$.

If 2 be subtracted, what will denote the difference? If 3 be subtracted? 4? &c.

2. John has a certain number of marbles; if 1 be taken away, twice the remainder will be equal to 12: how many has he?

ANALYSIS.—Let x denote the number; then, $x - 1$ will denote the number after 1 is taken away; and twice this number, or $2(x - 1) = 2x - 2$, will be equal to 12. If $2x$

diminished by 2 is equal to 12, $2x$ is equal to 12 plus 2, or 14; hence, x equals 14 divided by 2, or 7.

WRITTEN.

Let x denote the number; then,

$$x - 1 = \text{the number which remained, and}$$
$$2(x - 1) = 2x - 2 = 12;\ \text{hence,}$$
$$2x = 12 + 2,\ \text{or}\ 14;\ \text{and}\ x = \frac{14}{2} = 7.$$

VERIFICATION.

$$2(7 - 1) = 14 - 2 = 12;\ \text{also,}\ 2(7 - 1) = 2 \times 6 = 12.$$

3. Write 3 times $x - 1$. *Ans.* $3(x - 1)$.
What is the product equal to? *Ans.* $3x - 3$.

4. Write 4 times $x - 2$. *Ans.* $4(x - 2)$.
What is the product equal to? *Ans.* $4x - 8$.

5. Write 5 times $x - 5$. *Ans.* $5(x - 5)$.
What is the product equal to? *Ans.* $5x - 25$.

6. If x denotes a certain number, will $x - 1$ denote a greater or less number? how much less?

7. If $x - 1$ is equal to 4, what will x be equal to?
Ans. $4 + 1$, or 5.

8. If $x - 2$ is equal to 6, what is x equal to?

9. James and John together have 20 oranges; John has 6 less than James: how many has each?

10. A grocer sold 12 pounds of tea and coffee; if the tea be diminished by 3 pounds, and the remainder multiplied by 2, the product is the number of pounds of coffee: how many pounds of each?

11. Ann has a certain number of oranges; Jane has 1 less and twice her number added to Ann's make 13: how many has each?

ANALYSIS.—Let x denote the number of oranges which Ann has; then, $x - 1$ will denote the number Jane has, and $x + 2x - 2$, or $3x - 2$, will denote the number both have, which is 13. If $3x - 2$ equals 13, $3x$ will be equal to $13 + 2$, or 15; and if $3x$ is equal to 15, x will be equal to 15 divided by 3, which is 5: hence, Ann has 5 oranges and Jane 4.

WRITTEN.

Let x denote the number Ann has; then,

$$x - 1 = \text{the number Jane has; and}$$

$$2(x - 1) = 2x - 2 = \text{twice what Jane has; also,}$$

$$x + 2x - 2 = 3x - 2 = 13\text{; hence,}$$

$$3x = 13 + 2 = 15\text{; or } x = \frac{15}{3} = 5.$$

VERIFICATION.

$$5 - 4 = 1\text{; and } 2 \times 4 + 5 = 13.$$

12. Charles and John have 20 cents, and John has 6 less than Charles: how many has each?

13. James has twice as many oranges as lemons in his basket, and if 5 be taken from the whole number, 19 will remain: how many had he of each?

14. A basket contains apples, peaches, and pears; 29 in all. If 1 be taken from the number of apples, the remainder will denote the number of peaches, and twice that remainder will denote the number of pears: how many are there of each sort?

15. If $2x - 5$ equals 15, what is the value of x?

16. If $4x - 5$ is equal to 11, what is the value of x?

17 If $5x - 12$ is equal to 18, what is the value of x?

18. The sum of two numbers is 32, and the greater exceeds the less by 8: what are the numbers?

19. The sum of 2 numbers is 9; if the greater number be diminished by 5, and the remainder multiplied by 3, the product will be the less number: what are the numbers?

20. There are three numbers such that 1 taken from the

first will give the second; the second multiplied by 3 will give the third; and their sum is equal to 26: what are the numbers?

21. John and Charles together have just 31 oranges; if 1 be taken from John's, and the remainder be multiplied by 5, the product will be equal to Charles' number: how many has each?

22. A basket is filled with apples, lemons, and oranges, in all 26; the number of lemons exceed the apples by 2, and the number of oranges is double that of the lemons: how many are there of each?

LESSON XI.

1. John has a certain number of apples, the half of which is equal to 10: how many has he?

Analysis.—Let x denote the number of apples; then, x divided by 2 is equal to 10; if one half of x is equal to 10, twice one-half of x, or x, is equal to twice 10, which is 20; hence, x is equal to 20.

Note.—A similar analysis is applicable to any one of the fractional units. Let each question be solved according to the analysis.

2. John has a certain number of oranges, and one-third of his number is 15: how many has he?

3. If one-fifth of a number is 6, what is the number?

4. If one-twelfth of a number is 9, what is the number?

5. What number added to one-half of itself will give a sum equal to 12?

Analysis.—Denote the number by x; then, x plus one half of x equals 12. But x plus one-half of x equals three halves of x: hence, three halves of x equal 12. If three halves of x equal 12, one-half of x equals one-third of 12, or 4. If one-half of x equals 4, x equals twice 4, or 8, hence, x equals 8.

WRITTEN.

Let x denote the number; then,

$$x + \frac{1}{2}x = \frac{3}{2}x = 12; \text{ then,}$$

$$\frac{1}{2}x = 4, \text{ or } x = 8.$$

VERIFICATION.

$$8 + \frac{8}{2} = 8 + 4 = 12.$$

6. What number added to one-third of itself will give a sum equal to 12?

7. What number added to one-fourth of itself will give a sum equal to 20?

8. What number added to a fifth of itself will make 24?

9. What number diminished by one-half of itself will leave 4? Why?

10. What number diminished by one-third of itself will leave 6?

11. James gave one-seventh of his marbles to William, and then has 24 left: how many had he at first?

12. What number added to two-thirds of itself will give a sum equal to 20?

13. What number diminished by three-fourths of itself will leave 9?

14. What number added to five-sevenths of itself will make 24?

15. What number diminished by seven-eighths of itself will leave 4?

16. What number added to eight-ninths of itself will make 34?

ELEMENTARY ALGEBRA.

CHAPTER I.

DEFINITIONS AND EXPLANATORY SIGNS.

1. QUANTITY is anything which can be increased, diminished, and measured; as number, distance, weight, time, &c.

To measure a thing, is to find how many times it contains some other thing of the same kind, taken as a standard. The assumed standard is called the *unit of measure.*

2. MATHEMATICS is the science which treats of the measurement, properties, and relations of quantities.

In pure mathematics, there are but eight kinds of quantity, and consequently but eight kinds of UNITS, viz.: Units of *Number;* Units of *Currency;* Units of *Length;* Units of *Surface;* Units of *Volume;* Units of *Weight;* Units of *Time;* and Units of *Angular Measure.*

3. ALGEBRA is a branch of Mathematics in which the quantities considered are represented by letters, and the operations to be performed are indicated by signs.

---

1. What is quantity? What is the operation of measuring a thing? What is the assumed standard called?

2. What is Mathematics? How many kinds of quantity are there in the pure mathematics? Name the units of those quantities.

3. What is Algebra?

4. The quantities employed in Algebra are of two kinds, *Known* and *Unknown:*

Known Quantities are those whose values are given; and

Unknown Quantities are those whose values are required.

Known Quantities are generally represented by the leading letters of the alphabet, as, a, b, c, &c.

Unknown Quantities are generally represented by the final letters of the alphabet; as, x, y, z, &c.

When an unknown quantity becomes known, it is often denoted by the same letter with one or more accents; as, x', x'', x'''. These symbols are read: x *prime;* x *second;* x *third, &c.*

5. The SIGN OF ADDITION, +, is called *plus.* When placed between two quantities, it indicates that the second is to be added to the first. Thus, $a + b$, is read, a plus b, and indicates that b is to be added to a. If no sign is written, the sign + is understood.

The sign +, is sometimes called the *positive* sign, and the quantities before which it is written are called *positive quantities,* or *additive quantities.*

6. The SIGN OF SUBTRACTION, —, is called *minus.* When placed between two quantities, it indicates that the second is to be subtracted from the first. Thus, the expression,

---

4. How many kinds of quantities are employed in Algebra? How are they distinguished? What are known quantities? What are unknown quantities? By what are the known quantities represented? By what are the unknown quantities represented? When an unknown quantity becomes known, how is it often denoted?

5. What is the sign of addition called? When placed between two quantities, what does it indicate?

6. What is the sign of subtraction called? When placed between two quantities, what does it indicate?

$c - d$, read *c* minus *d*; indicates that *d* is to be subtracted from *c*. If *a* stands for 6, and *d* for 4, then $a - d$ is equal to $6 - 4$, which is equal to 2.

The sign $-$, is sometimes called the *negative* sign, and the quantities before which it is written are called *negative quantities*, or *subtractive quantities*.

7. The SIGN OF MULTIPLICATION, ×, is read, *multiplied by*, or *into*. When placed between two quantities, it indicates that the first is to be multiplied by the second. Thus, $a \times b$ indicates that *a* is to be multiplied by *b*. If *a* stands for 7, and *b* for 5, then, $a \times b$ is equal to 7×5, which is equal to 35.

The multiplication of quantities is also indicated by simply writing the letters, one after the other; and sometimes, by placing a point between them; thus,

$a \times b$ signifies the same thing as ab, or as $a.b$.
$a \times b \times c$ signifies the same thing as abc, or as $a.b.c$.

8. A FACTOR is any one of the multipliers of a product. Factors are of two kinds, *numeral* and *literal*. Thus, in the expression, $5abc$, there are four factors: the *numeral* factor, 5, and the three *literal* factors, *a*, *b*, and *c*.

9. The SIGN OF DIVISION, ÷, is read, *divided by*. When written between two quantities, it indicates that the first is to be divided by the second.

---

7. How is the sign of multiplication read? When placed between two quantities, what does it indicate? In how many ways may multiplication be indicated?

8. What is a factor? How many kinds of factors are there? How many factors are there in $3abc$?

9. How is the sign of division read? When written between two quantities, what does it indicate? How many ways are there of indicating division?

There are three signs used to denote *division*. Thus,

$a \div b$ denotes that a is to be divided by b.

$\frac{a}{b}$ denotes that a is to be divided by b.

$a \,\underline{|\, b}$ denotes that a is to be divided by b.

10. The SIGN OF EQUALITY, $=$, is read, *equal to*. When written between two quantities, it indicates that they are equal to each other. Thus, the expression, $a + b = c$, indicates that the sum of a and b is equal to c. If a stands for 3, and b for 5, c will be equal to 8.

11. The SIGN OF INEQUALITY, $> <$, is read, *greater than*, or *less than*. When placed between two quantities, it indicates that they are unequal, the greater one being placed at the opening of the sign. Thus, the expression, $a > b$, indicates that a is greater than b; and the expression, $c < d$, indicates that c is less than d.

12. The sign $.\dot{}.$ means, *therefore*, or *consequently*.

13. A COEFFICIENT is a number written before a quantity, to show how many times it is taken. Thus,

$$a + a + a + a + a = 5a,$$

in which 5 is the coefficient of a.

A coefficient may be denoted either by a *number*, or a *letter*. Thus, $5x$ indicates that x is taken 5 times, and ax

---

10. What is the sign of equality? When placed between two quantities, what does it indicate?

11. How is the sign of inequality read? Which quantity is placed on the side of the opening?

12. What does $.\dot{}.$ indicate?

13. What is a coefficient? How many times is a taken in $5a$. By what may a coefficient be denoted? If no coefficient is written, what coefficient is understood? In $5ax$, how many times is ax taken? How many times is x taken?

indicates that x is taken a times. If no coefficient is written, the coefficient 1 is understood. Thus, a is the same as $1a$.

14. AN EXPONENT is a number written at the right and above a quantity, to indicate how many times it is taken as a factor. Thus,

$$\begin{array}{rcc} a \times a & \text{is written} & a^2, \\ a \times a \times a & \text{"} & a^3, \\ a \times a \times a \times a & \text{"} & a^4, \\ \&c., & & \&c., \end{array}$$

in which 2, 3, and 4, are *exponents*. The expressions are read, a square, a cube or a third, a fourth; and if we have a^m, in which a enters m times as a factor, it is read, a to the mth, or simply a, *mth*. The exponent 1 is generally omitted. Thus, a^1 is the same as a, each denoting that a enters but once as a factor.

15. A POWER is a product which arises from the multiplication of equal factors. Thus,

$a \times a = a^2$ is the square, or second power of a.
$a \times a \times a = a^3$ is the cube, or third power of a.
$a \times a \times a \times a = a^4$ is the fourth power of a.
$a \times a \times \ldots\ldots = a^m$ is the mth power of a.

16. A ROOT of a quantity is one of the equal factors. The radical sign, $\sqrt{}$, when placed over a quantity, indicates that a root of that quantity is to be extracted. The root is indicated by a number written over the radical sign,

---

14. What is an exponent? In a^3, how many times is a taken as a factor? When no exponent is written, what is understood?

15. What is a power of a quantity? What is the third power of 2? Of 4? Of 6?

16. What is the root of a quantity? What indicates a root? What indicates the kind of root? What is the index of the square root? Of the cube root? Of the *m*th root?

called an *index*. When the index is 2, it is generally omitted. Thus,

$\sqrt[2]{a}$, or $\sqrt{a}$, indicates the square root of a.

$\sqrt[3]{a}$ indicates the cube root of a.

$\sqrt[4]{a}$ indicates the fourth root of a.

$\sqrt[m]{a}$ indicates the mth root of a.

17. An ALGEBRAIC EXPRESSION is a quantity written in algebraic language. Thus,

$3a$ is the algebraic expression of three times the number denoted by a;

$5a^2$ is the algebraic expression of five times the square of a;

$7a^3b^2$ is the algebraic expression of seven times the the cube of a multiplied by the square of b;

$3a - 5b$ is the algebraic expression of the difference between three times a and five times b;

$2a^2 - 3ab + 4b^2$ is the algebraic expression of twice the square of a, diminished by three times the product of a by b, augmented by four times the square of b.

18 A TERM is an algebraic expression of a single quantity. Thus, $3a$, $2ab$, $-5a^2b^2$, are terms.

19. The DEGREE of a term is the number of its literal factors. Thus,

$3a$ is a term of the first degree, because it contains but one literal factor.

---

17. What is an algebraic expression

18. What is a term?

19. What is the degree of a term? What determines the degree of a term?

$5a^2$ is of the second degree, because it contains two literal factors.

$7a^3b$ is of the fourth degree, because it contains four literal factors. The degree of a term is determined by the sum of the exponents of all its letters.

20. A MONOMIAL is a single term, unconnected with any other by the signs $+$ or $-$; thus, $3a^2$, $3b^3a$, are monomials.

21. A POLYNOMIAL is a collection of terms connected by the signs $+$ or $-$; as,

$$3a - 5, \text{ or, } 2a^3 - 3b + 4b^2.$$

22. A BINOMIAL is a polynomial of two terms; as,

$$a + b,\ 3a^2 - c^2,\ 6ab - c^2.$$

23. A TRINOMIAL is a polynomial of three terms; as,

$$abc - a^3 + c^3,\ ab - gh - f.$$

24. HOMOGENEOUS TERMS are those which contain the same number of literal factors. Thus, the terms, abc, $-a^3$, $+c^3$, are homogeneous; as are the terms, ab, $-gh$.

25. A POLYNOMIAL IS HOMOGENEOUS, when all its terms are homogeneous. Thus, the polynomial, $abc - a^3 + c^3$, is homogeneous; but the polynomial, $ab - gh - f$ is not homogeneous.

26. SIMILAR TERMS are those which contain the same literal factors affected with the same exponents. Thus,

$$7ab + 3ab - 2ab,$$

---

20. What is a monomial?
21. What is a polynomial?
22. What is a binomial?
23. What is a trinomial?
24. What are homogeneous terms?
25. When is a polynomial homogeneous?
26. What are similar terms?

are similar terms; and so also are,

$$4a^2b^2 - 2a^2b^2 - 3a^2b^2;$$

but the terms of the first polynomial and of the last, are not similar.

27. THE VINCULUM, ———, the *Bar* |, the *Parenthesis*, ( ), and the *Brackets*, [ ], are each used to connect several quantities, which are to be operated upon in the same manner. Thus, each of the expressions,

$$\overline{a+b+c} \times x, \qquad \begin{array}{r|l} a & x \\ +b & \\ +c & \end{array} \qquad (a+b+c) \times x,$$

$$\text{and} \qquad [a+b+c] \times x,$$

indicates, that the sum of a, b, and c, is to be multiplied by x.

28. THE RECIPROCAL of a quantity is 1, divided by that quantity; thus,

$$\frac{1}{a}, \ \frac{1}{a+b}, \ \frac{c}{d},$$

are the reciprocals of

$$a, \ a+b, \ \frac{d}{c}.$$

29. THE NUMERICAL VALUE of an algebraic expression, is the result obtained by assigning a numerical value to each letter, and then performing the operations indicated. Thus, the numerical value of the expression,

$$ab + bc + d,$$

when, $a = 1$, $b = 2$, $c = 3$, and $d = 4$, is

$$1 \times 2 + 2 \times 3 + 4 = 12;$$

by performing the indicated operations.

---

27. For what is the vincular used? Point out the other ways in which this may be done?

28. What is the reciprocal of a quantity?

29. What is the numerical value of an algebraical expression?

EXAMPLES IN WRITING ALGEBRAIC EXPRESSIONS.

1. Write a added to b. *Ans.* $a + b$.

2. Write b subtracted from a. *Ans.* $a - b$.

Write the following:

3. Six times the square of a, minus twice the square of b.

4. Six times a multiplied by b, diminished by 5 times c cube multiplied by d.

5. Nine times a, multiplied by c plus d, diminished by 8 times b multiplied by d cube.

6. Five times a minus b, plus 6 times a cube into b cube.

7. Eight times a cube into d fourth, into c fourth, plus 9 times c cube into d fifth, minus 6 times a into b, into c square.

8. Fourteen times a plus b, multiplied by a minus b, plus 5 times a, into c plus d.

9. Six times a, into c plus d, minus 5 times b, into a plus c, minus 4 times a cube b square.

10. Write a, multiplied by c plus d, plus f minus g.

11. Write a divided by $b + c$. Three ways.

12. Write $a - b$ divided by $a + b$.

13. Write a polynomial of three terms; of four terms; of five, of six.

14. Write a homogeneous binomial of the first degree; of the second; of the third; 4th; 5th; 6th.

15. Write a homogeneous trinomial of the first degree; with its second and third terms negative; of the second degree; of the 3rd; of the 4th.

16. Write in the same column, on the slate, or black-board, a monomial, a binomial, a trinomial, a polynomial of four terms, of five terms, of six terms and of seven terms, and all of the *same degree*.

INTERPRETATION OF ALGEBRAIC LANGUAGE.

Find the numerial values of the following expressions, when,

$$a = 1,\quad b = 2,\quad c = 3,\quad d = 4.$$

1. $ab + bc$. *Ans.* 8.
2. $a + bc + d$. *Ans.* 11.
3. $ad + b - c$. *Ans.* 3.
4. $ab + bc - d$. *Ans.* 4.
5. $(a + b)\,c^2 - d$. *Ans.* 23.
6. $(a + b)\,(d - b.)$ *Ans.* 6.
7. $(ab + ad)\,c + d$. *Ans.* 22.
8. $(ab + c)\,(ad - a)$. *Ans.* 15.
9. $3a^2b^2 - 2(a + d + 1)$. *Ans.* 0.
10. $\frac{a + c}{2} \times (a + d)$. *Ans.* 10.
11. $\frac{a^2 + b^2 + c^2}{7} \times \frac{a^3 + b^3 + c^3 - d}{2}$. *Ans.* 32.
12. $\frac{ab^4 - c - a^3}{6} \times \frac{4a^2 - b + d^3}{33}$ *Ans.* 4.

Find the numerical values of the following expressions, when,

$$a = 4,\quad b = 3,\quad c = 2,\quad \text{and } d = 1.$$

13. $\frac{a}{2} - \frac{b}{3} + c - d$. *Ans.* 2.
14. $5\left(\frac{ab}{3} - \frac{a - d}{3}\right)$. *Ans.* 15.
15. $[(a^2b + 1)d] \div (a^2b + d)$. *Ans.* 1.
16. $4\left(abc - \frac{b^3}{9}\right) \times (30c^3 - ab^3d^3)$. *Ans.* 11088.
17. $\frac{a + b + c}{a - b + d} + \frac{abcd}{ab} + \frac{4a^2 + b^2 - d^2}{bc + b}$. *Ans.* 14$\frac{1}{2}$.
18. $\frac{15(a+d+b)}{3c^2} - \frac{a-c}{2} + \frac{3}{abd} \times a^3b^3c^3d^3$. *Ans.* 3465.

CHAPTER II.

FUNDAMENTAL OPERATIONS.

ADDITION.

30. Addition is the operation of finding the simplest equivalent expression for the aggregate of two or more algebraic quantities. Such expression is called their Sum.

When the terms are similar and have like signs.

31. 1. What is the sum of a, $2a$, $3a$, and $4a$? Take the sum of the coefficients, and annex the literal parts. The first term, a, has a coefficient, 1, understood (Art. 13).

$$\begin{array}{r} +\ a \\ +\ 2a \\ +\ 3a \\ +\ 4a \\ \hline +\ 10a \end{array}$$

2. What is the sum of $2ab$, $3ab$, $6ab$, and ab. When no sign is writtten, the sign $+$ is understood (Art 5).

$$\begin{array}{r} 2ab \\ 3ab \\ 6ab \\ ab \\ \hline 12ab \end{array}$$

Add the following:

$$\begin{array}{rrrr} (3.) & (4.) & (5.) & (6.) \\ a & 8ab & 7ac & +\ 4abc \\ a & 7ab & 5ac & 3abc \\ \hline +\ 2a & 15ab & 12ac & +\ 7abc \end{array}$$

---

30. What is addition?

31. What is the rule for addition when the terms are similar and have like signs?

$$\begin{array}{cccc}
(7.) & (8.) & (9.) & (10.) \\
\begin{array}{r} -3abc \\ -2abc \\ \hline -5abc \end{array} &
\begin{array}{r} -3ad \\ -2ad \\ \hline -5ad \end{array} &
\begin{array}{r} -2adf \\ -6adf \\ \hline -8adf \end{array} &
\begin{array}{r} -9abd \\ -15abd \\ \hline -24abd \end{array}
\end{array}$$

Hence, when the terms are similar and have like signs:

RULE.

Add the coefficients, and to their sum prefix the common sign; to this, annex the common literal part.

EXAMPLES.

$$\begin{array}{ccc}
(11.) & (12.) & (13.) \\
\begin{array}{r} 9ab + ax \\ 8ab + 3ax \\ 12ab + 4ax \\ \hline \end{array} &
\begin{array}{r} 8ac^2 - 3b^2 \\ 7ac^2 - 8b^2 \\ 3ac^2 - 9b^2 \\ \hline \end{array} &
\begin{array}{r} 15ab^3c^4 - 12abc^2 \\ 12ab^3c^4 - 15abc^2 \\ ab^3c^4 - abc^2 \\ \hline \end{array}
\end{array}$$

When the terms are similar and have unlike signs.

32. The signs, + and –, stand in direct opposition to each other.

If a merchant writes + before his gains and – before his losses, at the end of the year the sum of the plus numbers will denote the gains, and the sum of the minus numbers the losses. If the gains exceed the losses, the *difference*, which is called the *algebraic sum*, will be plus; but if the losses exceed the gains, the *algebraic sum* will be minus.

1. A merchant in trade gained $1500 in the first quarter of the year, $3000 in the second quarter, but lost $3000 in the third quarter, and $800 in the fourth: what was the result of the year's business?

$$\begin{array}{llr@{\qquad}llr}
\text{1st quarter,} & & +1500 & \text{3d quarter,} & & -3000 \\
\text{2d} & \text{“} & 3000 & \text{4th} & \text{“} & -800 \\
\cline{3-3}\cline{6-6}
 & & +4500 & & & -3800
\end{array}$$

$$+4500 - 3800 = +700, \text{ or } \$700 \text{ gain.}$$

---

32. What is the rule when the terms are similar and have unlike signs?

2. A merchant in trade gained \$1000 in the first quarter, and \$2000 the second quarter; in the third quarter he lost \$1500, and in the fourth quarter \$1800: what was the result of the year's business?

1st quarter,	$+ 1000$	3d quarter	$- 1500$
2d "	$+ 2000$	4th "	$- 1800$
	$+ 3000$		$- 3300$

$+ 3000 - 3300 = - 300$, or \$300 loss.

3. A merchant in the first half-year gained a dollars and lost b dollars; in the second half-year he lost a dollars and gained b dollars: what is the result of the year's business?

1st half-year,	$+ a$	$- b$
2d "	$- a$	$+ b$
Result,	0	0

Hence, *the algebraic sum of a positive and negative quantity is their arithmetical difference, with the sign of the greater prefixed.* Add the following:

$$\begin{array}{rcrcr} 8ab & \qquad & 4acb^2 & \qquad & -\ 4a^2b^2c^2 \\ 3ab & & -\ 8acb^2 & & +\ 6a^2b^2c^2 \\ -\ 6ab & & acb^2 & & -\ 2a^2b^2c^2 \\ \hline 5ab & & -\ 3acb^2 & & 0 \end{array}$$

Hence, when the terms are similar and have unlike signs:

I. *Write the similar terms in the same column:*

II. *Add the coefficients of the additive terms, and also the coefficients of the subtractive terms:*

III. *Take the difference of these sums, prefix the sign of the greater, and then annex the literal part.*

EXAMPLES.

1. What is the sum of

$$2a^2b^3 - 5a^2b^3 + 7a^2b^3 + 6a^2b^3 - 11a^2b^3?$$

Having written the similar terms in the same column, we find the sum of the positive coefficients to be 15, and the sum of the negative coefficients to be -16: their difference is -1; hence, the sum is $-a^2b^3$.

$$\begin{array}{r} 2a^2b^3 \\ -\ \ 5a^2b^3 \\ +\ \ 7a^2b^3 \\ +\ \ 6a^2b^3 \\ -\ 11a^2b^3 \\ \hline -\ \ \ a^2b^3 \end{array}$$

2. What is the sum of

$3a^2b + 5a^2b - 3a^2b + 4a^2b - 6a^2b - a^2b$? *Ans.* $2a^2b$.

3. What is the sum of

$12a^3bc^2 - 4a^3bc^2 + 6a^3bc^2 - 8a^3bc^2 + 11a^3bc^2$? *Ans.* $17a^3bc^2$.

4. What is the sum of

$4a^2b - 8a^2b - 9a^2b + 11a^2b$? *Ans.* $-2a^2b$

5. What is the sum of

$7abc^2 - abc^2 - 7abc^2 + 8abc^2 + 6abc^2$? *Ans.* $13abc^2$.

6. What is the sum of

$9cb^3 - 5cb^3 - 8ac^2 + 20cb^3 + 9ac^2 - 24cb^3$? *Ans.* $+ac$.

To add any Algebraic Quantities.

33. 1. What is the sum of $3a$, $5b$, and $-2c$? Write the quantities, thus,

$$3a + 5b - 2c;$$

which denotes their sum, as there are no *similar terms*.

2. Let it be required to find the sum of the quantities,

$$\begin{array}{rrr} 2a^2 & -\ 4ab & \\ 3a^2 & -\ 3ab & +\ \ b^2 \\ & 2ab & -\ 5b^2 \\ \hline 5a^2 & -\ 5ab & -\ 4b^2 \end{array}$$

---

33. What is the rule for the addition of any algebraic quantities?

From the preceding examples, we have, for the addition of algebraic quantities, the following

RULE.

I. *Write the quantities to be added, placing similar terms in the same column, and giving to each its proper sign:*

II. *Add up each column separately and then annex the dissimilar terms with their proper signs.*

EXAMPLES.

1. Add together the polynomials,

$3a^2 - 2b^3 - 4ab$, $5a^2 - b^2 + 2ab$, and $3ab - 3c^2 - 2b^3$.

The term $3a^2$ being similar to $5a^2$, we write $8a^2$ for the result of the reduction of these two terms, at the same time slightly crossing them, as in the first term.

$$\left\{\begin{array}{rrrr}
3\cancel{a}^2 & - 4\cancel{a}b & - 2\cancel{b}^2 & \\
5\cancel{a}^2 & + 2\cancel{a}b & - \cancel{b}^2 & \\
 & + 3\cancel{a}b & - 2\cancel{b}^2 & - 3\cancel{c}^2 \\
\hline
8a^2 & + ab & - 5b^2 & - 3c^3
\end{array}\right.$$

Passing then to the term $-4ab$, which is similar to $+2ab$ and $+3ab$, the three reduce to $+ab$, which is placed after $8a^2$, and the terms crossed like the first term. Passing then to the terms involving b^2, we find their sum to be $-5b^2$, after which we write $-3c^2$.

The marks are drawn across the terms, that none of them may be overlooked and omitted.

(2.)

$$\begin{array}{r}
7abc + 9ax \\
-3abc - 3ax \\
\hline
4abc + 6ax
\end{array}$$

(3.)

$$\begin{array}{r}
8ax + 3b \\
5ax - 9b \\
\hline
13ax - 6b
\end{array}$$

(4.)

$$\begin{array}{r}
12a - 6c \\
-3a - 9c \\
\hline
9a - 15c
\end{array}$$

NOTE.—If $a = 5$, $b = 4$, $c = 2$, $x = 1$, what are the numerical values of the several sums above found?

(5.)

$$\begin{array}{r} 9a + f \\ -6a + g \\ -2a - f \\ \hline \end{array}$$

(6.)

$$\begin{array}{r} 6ax - 8ac \\ -7ax - 9ac \\ ax + 17ac \\ \hline \end{array}$$

(7.)

$$\begin{array}{r} 3af + g + m \\ ag - 3af - m \\ ab - ag + 3g \\ \hline \end{array}$$

(8.)

$$\begin{array}{r} 7x + 3ab + 3c \\ -3x - 3ab - 5c \\ 5x - 9ab - 9c \\ \hline \end{array}$$

(9.)

$$\begin{array}{r} 8x^2 + 9acx + 13a^2b^2c^2 \\ -7x^2 - 13acx + 14a^2b^2c^2 \\ -4x^2 + 4acx - 20a^2b^2c^2 \\ \hline \end{array}$$

(10.)

$$\begin{array}{r} 22h - 3c - 7f + 3g \\ -3h + 8c - 2f - 9g + 5x \\ \hline \end{array}$$

(11.)

$$\begin{array}{r} 19ah^2 + 3a^3b^4 - 8ax^3 \\ -17ah^2 - 9a^3b^4 + 9ax^3 \\ \hline \end{array}$$

(12.)

$$\begin{array}{r} 7x - 9y + 5z + 3 - g \\ -x - 3y - 8 - g \\ -x + y - 3z + 1 + 7g \\ -2x + 6y + 3z - 1 - g \\ \hline \end{array}$$

(13.)

$$\begin{array}{r} 8a + b \\ 2a - b + c \\ -3a + b + 2d \\ -6b - 3c + 3d \\ \hline \end{array}$$

14. Add together $-b + 3c - d - 115e + 6f - 5g$, $3b - 2c - 3d - e + 27f$, $5c - 8d + 3f - 7g$, $-7b - 6c + 17d + 9e - 5f + 11g$, $-3b - 5d - 2e + 6f - 9g + h$.

Ans. $-8b - 109e + 37f - 10g + h$.

15. Add together the polynomials $7a^2b - 3abc - 8b^2c - 9c^3 + cd^2$, $8abc - 5a^2b + 3c^3 - 4b^2c + cd^2$, and $4a^2b - 8c^3 + 9b^2c - 3d^3$.

Ans. $6a^2b + 5abc - 3b^2c - 14c^3 + 2cd^2 - 3d^3$.

16. What is the sum of, $5a^2bc + 6bx - 4af$, $-3a^2bc - 6bx + 14af$, $-af + 9bx + 2a^2bc$, $+6af - 8bx + 6a^2bc$?

Ans. $10a^2bc + bx + 15af$.

17. What is the sum of $a^2n^2 + 3a^3m + b$, $-6a^2n^2 - 6a^3m - b$, $+9b - 9a^3m - 5a^2n^2$?

Ans. $-10a^2n^2 - 12a^3m + 9b$.

18. What is the sum of $4a^3b^2c - 16a^4x - 9ax^3d$, $+ 6a^3b^2c - 6ax^3d + 17a^4x$, $+ 16ax^3d - a^4x - 9a^3b^2c$? *Ans.* $a^3b^2c + ax^3d$.

19. What is the sum of $- 7g + 3b + 4g - 2b + 3g - 3b + 2b$? *Ans.* 0.

20. What is the sum of, $ab + 3xy - m - n$, $- 6xy - 3m + 11n + cd$, $+ 3xy + 4m - 10n + fg$? *Ans.* $ab + cd + fg$.

21. What is the sum of $4xy + n + 6ax + 9am$, $- 6xy + 6n - 6ax - 8am$, $2xy - 7n + ax - am$? *Ans.* $+ ax$.

$$\begin{array}{ccc}
(22.) & (23.) & (24.) \\
2(a + b) & 5(a^2 - c^2) & 9(c^3 - af^3) \\
3(a + b) & -4(a^2 - c^2) & 7(c^3 - af^3) \\
\underline{2(a + b)} & -\underline{1(a^2 - c^2)} & -\underline{10(c^3 - af^3)} \\
7(a + b) & & 6(c^3 - af^3)
\end{array}$$

NOTE. The quantity within the parenthesis must be regarded as a *single* quantity.

25. Add $3a(g^2 - h^2) - 2a(g^2 - h^2) + 4a(g^2 - h^2) + 8a(g^2 - h^2) - 2a(g^2 - h^2)$. *Ans.* $11a(g^2 - h^2)$.

26. Add $3c(a^2c - b^2) - 9c(a^2c - b^2) - 7c(a^2c - b^2) + 15c(a^2c - b^2) + c(a^2c - b^2)$. *Ans.* $3c(a^2c - b^2)$.

34. In algebra, the term *add* does not always, as in arithmetic, convey the idea of augmentation; nor the term *sum*, the idea of a number numerically greater than any of the numbers added. For, if to a we add $- b$, we have, $a - b$, which is, arithmetically speaking, a difference between the number of units expressed by a, and the number

---

34. Do the words *add* and *sum*, in Algebra, convey the same ideas as in Arithmetic. What is the algebraic sum of 9 and -4? Of 8 and -2? May an algebraic sum be negative? What is the sum of 5 and -10? How are such sums distinguished from arithmetical sums?

of units expressed by b. Consequently, this result is numerically less than a. To distinguish this sum from an arithmetical sum, it is called the *algebraic sum.*

SUBTRACTION.

35. Subtraction is the operation of finding the difference between two algebraic quantities.

36. The quantity to be subtracted is called the *Subtrahend;* and the quantity from which it is taken, is called the *Minuend.*

The *difference* of two quantities, is such a quantity as added to the subtrahend will give a sum equal to the minuend.

EXAMPLES.

1. From $17a$ take $6a$.

In this example, $17a$ is the minuend, and $6a$ the subtrahend: the difference is $11a$; *because,* $11a$, added to $6a$, gives $17a$.

OPERATION.

$$\begin{array}{r}17a\\6a\\\hline 11a\end{array}$$

The difference may be expressed by writing the quantities thus:

$$17a - 6a = 11a;$$

in which the sign of the subtrahend is changed from + to −.

2. From $15x$ take $-9x$.

The *difference,* or remainder, is such a quantity, as being added to the subtrahend, $-9x$, will give the minuend, $15x$. That quantity is $24x$, and may be found by simply *changing the sign*

OPERATION

$$\begin{array}{r}15x\\-\quad 9x\\\hline 24a\end{array}$$

of the subtrahend, and adding. Whence, we may write,

$$15x - (-9x) = 24x.$$

3. From $10ax$ take $a - b$.

The *difference*, or *remainder*, is such a quantity, as added to $a - b$, will give the minuend, $10ax$: what is that quantity?

If you change the signs of both terms of the subtrahend, and add, you have, $10ax - a + b$. Is this the true remainder? Certainly. For, if you add the remainder to the subtrahend, $a - b$, you obtain the minuend, $10ax$.

OPERATION.

$$\begin{array}{lrrr}
 & 10ax & & \\
 & & +a & -b \\
\hline
\text{Rem.} & 10ax & -a & +b \\
\text{add} & & +a & -b \\
\hline
 & 10ax & &
\end{array}$$

It is plain, that if you change the signs of all the terms of the subtrahend, and then add them to the minuend, and to this result add the given subtrahend, the last sum can be no other than the given minuend; hence, the *first* result is the true difference, or remainder (Art. 36).

Hence, for the subtraction of algebraic quantities, we have the following

RULE.

I. *Write the terms of the subtrahend under those of the minuend, placing similar terms in the same column:*

II. *Conceive the signs of all the terms of the subtrahend to be changed from + to −, or from − to +, and then proceed as in Addition.*

EXAMPLES OF MONOMIALS.

	(1.)	(2.)	(3.)
From	$3ab$	$6ax$	$9abc$
take	$2ab$	$3ax$	$7abc$
Rem.	ab	$3ax$	$2abc$

	(4.)	(5.)	(6.)
From	$16a^2b^2c$	$17a^3b^3c$	$24a^2b^2x$
take	$9a^2b^2c$	$3a^3b^3c$	$7a^2b^2x$
Rem.	$7a^2b^2c$	$14a^3b^3c$	$17a^2b^2x$

	(7.)	(8.)	(9.)
From	$3ax$	$4abx$	$2am$
take	$8c$	$9ac$	ax
Rem.	$3ax - 8c$	$4abx - 9ac$	$2am - ax$

10. From $9a^2b^2$ take $3a^2b^2$. *Ans.* $6a^2b^2$.
11. From $16a^2xy$ take $-15a^2xy$. *Ans.* $31a^2xy$.
12. From $12a^4y^3$ take $8a^4y^3$. *Ans.* $4a^4y^3$.
13. From $19a^5x^8y$ take $-18a^5x^8y$. *Ans.* $37a^5x^8y$.
14. From $3a^2b^3$ take $3a^3b^2$. *Ans.* $3a^2b^3 - 3a^3b^2$.
15. From $7a^2b^4$ take $6a^4b^2$. *Ans.* $7a^2b^4 - 6a^4b^2$.
16. From $3ab^2$ take a^2b^5. *Ans.* $3ab^2 - a^2b^5$.
17. From x^2y take y^2x. *Ans.* $x^2y - y^2x$.
18. From $3x^2y^3$ take xy. *Ans.* $3x^2y^3 - xy$.
19. From $8a^2y^3x$ take xyz. *Ans.* $8a^2y^3x - xyz$.
20. From $9a^2b^2$ take $-3a^2b^2$. *Ans.* $12a^2b^2$.
21. From $14a^2y^2$ take $-20a^2y^2$. *Ans.* $34a^2y^2$.
22. From $-24a^4b^5$ take $16a^4b^5$. *Ans.* $-40a^4b^5$.
23. From $-13x^2y^4$ take $-14x^2y^4$. *Ans.* x^2y^4.
24. From $-47a^3x^2y$ take $-5a^3x^2y$. *Ans.* $-42a^3x^2y$.
25. From $-94a^2x^2$ take $3a^2x^2$. *Ans.* $-97a^2x^2$
26. From $a + x^2$ take $-y^3$. *Ans.* $a + x^2 + y^3$.
27. From $a^3 + b^3$ take $-a^3 - b^3$. *Ans* $2a^3 + 2b^3$
28. From $-16a^2x^3y$ take $-19a^2x^3y$. *Ans.* $+3a^2x^3y$.
29. From $a^2 - x^2$ take $a^2 + x^2$. *Ans.* $-2x^2$.

GENERAL EXAMPLES.

(1.)

$$\begin{array}{ll} \text{From} & 6ac - 5ab + c^2 \\ \text{take} & 3ac + 3ab + 7c \\ \hline \text{Rem.} & 3ac - 8ab + c^2 - 7c. \end{array}$$

The same with the signs of the lower line changed.

(1.)

$$\begin{array}{l} 6ac - 5ab + c^2 \\ -3ac - 3ab - 7c \\ \hline 3ac - 8ab + c^2 - 7c \end{array}$$

(2.)

$$\begin{array}{ll} \text{From} & 6ax - a + 3b^2 \\ \text{take} & 9ax - x + b^2 \\ \hline \text{Rem.} & -3ax - a + x + 2b^2. \end{array}$$

(3.)

$$\begin{array}{l} 6yx - 3x^2 + 5b \\ yx - 3 + a \\ \hline 5yx - 3x^2 + 3 + 5b - a. \end{array}$$

(4.)

$$\begin{array}{ll} \text{From} & 5a^3 - 4a^2b + 3b^2c \\ \text{take} & -2a^3 + 3a^2b - 8b^2c \\ \hline \text{Rem.} & 7a^3 - 7a^2b + 11b^2c. \end{array}$$

(5.)

$$\begin{array}{l} 4ab - cd + 3a^2 \\ 5ab - 4cd + 3a^2 + 5b^2 \\ \hline -ab + 3cd - 5b^2. \end{array}$$

6. From $a + 8$ take $c - 5$. *Ans.* $a - c + 13$.
7. From $6a^2 - 15$ take $9a^2 + 30$. *Ans.* $-3a^2 - 45$.
8. From $6xy - 8a^2c^3$ take $-7xy - a^2c^3$. *Ans.* $13xy - 7a^2c^3$
9. From $a + c$ take $-a - c$. *Ans.* $2a + 2c$
10. From $4(a + b)$ take $2(a + b)$. *Ans.* $2(a + b)$
11. From $3(a + x)$ take $(a + x)$. *Ans.* $2(a + x)$
12. From $9(a^2 - x^2)$ take $-2(a^2 - x^2)$. *Ans.* $11(a^2 - x^2)$
13. From $6a^2 - 15b^2$ take $-3a^2 + 9b^2$. *Ans.* $9a^2 - 24b^2$
14. From $3a^m - 2b^n$ take $a^m - 2b^n$. *Ans.* $2a^m$.
15. From $9c^2m^2 - 4$ take $4 - 7c^2m^2$. *Ans.* $16c^2m^2 - 8$.
16. From $6am + y$ take $3am - x$ *Ans.* $3am + x + y$
17. From $3ax$ take $3ax - y$. *Ans.* $+y$.

18. From $-7f + 3m - 8x$ take $-6f - 5m - 2x + 3d + 8$. *Ans.* $-f + 8m - 6x - 3d - 8$.

19. From $-a - 5b + 7c + d$ take $4b - c + 2d + 2k$. *Ans.* $-a - 9b + 8c - d - 2k$.

20. From $-3a + b - 8c + 7e - 5f + 3h - 7x - 13y$ take $k + 2a - 9c + 8e - 7x + 7f - y - 3l - k$. *Ans.* $-5a + b + c - e - 12f + 3h - 12y + 3l$.

21. From $2x - 4a - 2b + 5$ take $8 - 5b + a + 6x$. *Ans.* $-4x - 5a + 3b - 3$.

22. From $3a + b + c - d - 10$ take $c + 2a - d$. *Ans.* $a + b - 10$.

23. From $3a + b + c - d - 10$ take $b - 19 + 3a$. *Ans.* $c - d + 9$.

24. From $a^3 + 3b^2c + ab^2 - abc$ take $b^3 + ab^2 - abc$. *Ans.* $a^3 + 3b^2c - b^3$.

25 From $12x + 6a - 4b + 40$ take $4b - 3a + 4x + 6d - 10$. *Ans.* $8x + 9a - 8b - 6d + 50$.

26. From $2x - 3a + 4b + 6c - 50$ take $9a + x + 6b - 6c - 40$. *Ans.* $x - 12a - 2b + 12c - 10$.

27. From $6a - 4b - 12c + 12x$ take $2x - 8a + 4b - 6c$. *Ans.* $14a - 8b - 6c + 10x$.

38. In Algebra, the term *difference* does not always, as in Arithmetic, denote a number less than the minuend. For, if from a we subtract $-b$, the remainder will be $a + b$; and this is numerically greater than a. We distinguish between the two cases by calling this result the *algebraic difference.*

---

38. In Algebra, as in Arithmetic, does the term *difference* denote a number less than the minuend? How are the results in the two cases, distinguished from each other?

39. When a polynomial is to be subtracted from an algebraic quantity, we inclose it in a parenthesis, place the minus sign before it, and then write it after the minuend. Thus, the expression,

$$6a^2 - (3ab - 2b^2 + 2bc),$$

indicates that the polynomial, $3ab - 2b^2 + 2bc$, is to be taken from $6a^2$. Performing the operations indicated, by the rule for subtraction, we have the equivalent expression:

$$6a^2 - 3ab + 2b^2 - 2bc.$$

The last expression may be changed to the former, by changing the signs of the last three terms, inclosing them in a parenthesis, and prefixing the sign —. Thus,

$$6a^2 - 3ab + 2b^2 - 2bc = 6a^2 - (3ab - 2b^2 + 2bc).$$

In like manner any polynomial may be transformed, as indicated below:

$$\begin{aligned} 7a^3 - 8a^2b - 4b^2c + 6b^3 &= 7a^3 - (8a^2b + 4b^2c - 6b^3) \\ &= 7a^3 - 8a^2b - (4b^2c - 6b^3). \\ 8a^3 - 7b^2 + c - d &= 8a^3 - (7b^2 - c + d) \\ &= 8a^3 - 7b^2 - (-c + d). \\ 9b^3 - a + 3a^2 - d &= 9b^3 - (a - 3a^2 + d) \\ &= 9b^3 - a - (-3a^2 + d). \end{aligned}$$

NOTE.—The sign of every quantity is changed when it is placed within a parenthesis, and also when it is brought out.

40. From the preceding principles, we have,

$$\begin{aligned} a - (+b) &= a - b; \text{ and} \\ a - (-b) &= a + b. \end{aligned}$$

---

39. How is the subtraction of a polynomial indicated? How is this indicated operation performed? How may the result be again put under the first form? What is the general rule in regard to the parenthesis?

40. What is the sign which immediately precedes a quantity called? What is the sign which precedes the parenthesis called? What is the

The sign immediately preceding b is called the *sign of the quantity;* the sign preceding the parenthesis is called the *sign of operation;* and the sign resulting from the combination of the signs,is called the *essential sign.*

When the sign of operation is different from the sign of the quantity, the essential sign will be $-$; when the sign of operation is the same as the sign of the quantity, the essential sign will be $+$.

MULTIPLICATION.

41. 1. If a man earns a dollars in 1 day, how much will he earn in 6 days?

ANALYSIS.—In 6 days he will earn six times as much as in 1 day. If he earns a dollars in 1 day, in 6 days he will earn $6a$ dollars.

2. If one hat costs d dollars, what will 9 hats cost?
Ans. $9d$ dollars.

3. If 1 yard of cloth costs c dollars, what will 10 yards cost? *Ans.* $10c$ dollars.

4. If 1 cravat costs b cents, what will 40 cost?
Ans. $40b$ cents.

5. If 1 pair of gloves costs b cents, what will a pairs cost?

ANALYSIS.—If 1 pair of gloves cost b cents, a pairs will cost as many times b cents as there are units in a: that is, b taken a times, or ab; which denotes the *product* of b by a, or of a by b.

---

resulting sign called? When the sign of operation is different from the sign of the quantity, what is the essential sign? When the sign of operation is the same as the sign of the quantity, what is the essential sign?

41. What is Multiplication? What is the quantity to be multiplied called? What is that called by which it is multiplied? What is the result called?

MULTIPLICATION *is the operation of finding the product of two quantities.*

The quantity to be multiplied is called the *Multiplicand;* that by which it is multiplied is called the *Multiplier;* and the result is called the *Product.* The Multiplier and Multiplicand are called *Factors* of the Product.

6. If a man's income is $3a$ dollars a week, how much will he receive in $4b$ weeks?

$$3a \times 4b = 12ab.$$

If we suppose $a = 4$ dollars, and $b = 3$ weeks, the product will be 144 dollars.

NOTE.—It is proved in Arithmetic (Davies' School, Art. 48. University, Art. 50), that the product is not altered by changing the arrangement of the factors; that is,

$$12ab = a \times b \times 12 = b \times a \times 12 = a \times 12 \times b.$$

MULTIPLICATION OF POSITIVE MONOMIALS.

42. Multiply $3a^2b^2$ by $2a^2b$. We write,

$$3a^2b^2 \times 2a^2b = 3 \times 2 \times a^2 \times a^2 \times b^2 \times b$$
$$= 3 \times 2\, a\, a\, a\, a\, b\, b\, b;$$

in which a is a factor 4 times, and b a factor 3 times; hence (Art. 14),

$$3a^2b^2 \times 2a^2b = 3 \times 2a^4b^3 = 6a^4b^3,$$

in which *we multiply the coefficients together, and add the exponents of the like letters.*

The product of any two positive monomials may be found in like manner; hence the

RULE.

I. *Multiply the coefficients together for a new coefficient:*

II. *Write after this coefficient all the letters in both mono-*

---

42. What is the rule for multiplying one monomial by another?

mials, giving to each letter an exponent equal to the sum of its exponents in the two factors.

EXAMPLES.

1. $8a^2bc^2 \times 7abd^2 = 56a^3b^2c^2d^2$.
2. $21a^3b^2cd \times 8abc^3 = 168a^4b^3c^4d$.
3 $4abc \times 7df = 28abcdf$.

	(4.)	(5.)	(6.)
Multiply	$3a^2b$	$12a^2x$	$6xyz$
by	$2a^2b$	$12x^2y$	ay^2z
	$6a^4b^2$	$144a^2x^3y$	$6axy^3z^2$

(7.)	(8.)	(9.)
a^2xy	$3ab^2c^3$	$87ax^2y$
$2xy^2$	$9a^2b^3c$	$3b^3x^4y^3$
$2a^2x^2y^3$	$27a^3b^5c^4$	$261ab^3x^6y^4$

10. Multiply $5a^3b^2x^2$ by $6c^5x^6$. *Ans.* $30a^3b^2c^5x^8$.
11. Multiply $10a^4b^5c^8$ by $7acd$. *Ans.* $70a^5b^5c^9d$.
12. Multiply $36a^8b^7c^6d^5$ by $20ab^2c^3d^4$. *Ans.* $720a^9b^9c^9d^9$
13. Multiply $5a^m$ by $3ab^n$. *Ans.* $15a^{m+1}b^n$.
14. Multiply $3a^mb^3$ by $6a^2b^n$. *Ans.* $18a^{m+2}b^{n+3}$.
15. Multiply $6a^mb^n$ by $9a^5b^7$. *Ans.* $54a^{m+5}b^{n+7}$.
16. Multiply $5a^mb^n$ by $2a^pb^q$. *Ans.* $10a^{m+p}b^{n+q}$.
17. Multiply $5a^mb^2c^2$ by $2ab^nc$. *Ans.* $10a^{m+1}b^{n+2}c^3$.
18. Multiply $6a^2b^mc^n$ by $3a^3b^2c^2$. *Ans.* $18a^5b^{m+2}c^{n+2}$.
19. Multiply $20a^5b^5cd$ by $12a^2x^2y$. *Ans.* $240a^7b^5cdx^2y$.
20. Multiply $14a^4b^6d^4y$ by $20a^3c^2x^2y$. *A.* $280a^7b^6c^2d^4x^2y^2$
21. Multiply $8a^3b^3y^4$ by $7a^4bxy^5$. *Ans.* $56a^7b^4xy^9$.
22. Multiply $75axyz$ by $5a^5bcdx^2y^2$. *Ans.* $375a^6bcdx^3y^3z$.

23. Multiply $64a^3m^5x^4yz$ by $8ab^2c^3$. *A.* $512a^4b^2c^3m^5x^4yz$.

24. Multiply $9a^2b^2c^2d^3$ by $12a^3b^4c^6$. *Ans.* $108a^5b^6c^8d^3$.

25. Multiply $216ab^7c^3d^8$ by $3a^3b^2c^5$. *Ans.* $648a^4b^9c^8d^8$

26. Multiply $70a^8b^7c^4d^2fx$ by $12a^7b^5c^3dx^2y^3$.
Ans. $840a^{15}b^{12}c^7d^3fx^3y^3$.

MULTIPLICATION OF POLYNOMIALS.

43. 1. Multiply $a - b$ by c.

It is required to take the *difference* between a and b, c times; or, to take c, $a - b$ times.

As we can not subtract b from c, we begin by taking a, c times, which is ac; but this product is too large by b taken c times, which is bc; hence, the true product is $ac - bc$.

$$\begin{array}{l} a - b \\ c \\ \hline ac - bc \end{array}$$

$$\begin{array}{rcrcr} 8 & - & 3 & = & 5 \\ 7 & . & . & . & 7 \\ \hline 56 & - & 21 & = & 35 \end{array}$$

If a, b, and c, denote numbers, as $a = 8$, $b = 3$, and $c = 7$, the operation may be written in figures.

Multiply $a - b$ by $c - d$.

It is required to take $a - b$ as many times as there are units in $c - d$.

If we take $a - b$, c times, we have $ac - bc$; but this product is too large by $a - b$ taken d times. But $a - b$ taken d times, is $ad - db$. Subtracting this product from the preceding, by changing the signs of its terms (Art. 37), and we have,

$$\begin{array}{l} a - b \\ c - d \\ \hline ac - bc \\ \quad - ad + bd \\ \hline ac - bc - ad + bd \end{array}$$

$$\begin{array}{rrrcr} 8 - & 3 & & = & 5 \\ 7 - & 2 & & = & 5 \\ \hline 56 - & 21 & & & \\ - & 16 & + 6 & & \\ \hline 56 - & 37 & + 6 & = & 25. \end{array}$$

$$(a - b) + (a - c) = ab - bc - ad + bd.$$

Hence, we have the following

RULE FOR THE SIGNS.

I. *When the factors have like signs, the sign of their product will be* + *:*

II. *When the factors have unlike signs, the sign of their product will be* − *:*

Therefore, we say in Algebraic language, that + multiplied by +, or − multiplied by −, gives +; − multiplied by + or + multiplied by −, gives −.

Hence, for the multiplication of polynomials, we have the following

RULE.

Multiply every term of the multiplicand by each term of the multiplier, observing that like signs give +, and unlike signs −; then reduce the result to its simplest form.

EXAMPLES IN WHICH ALL THE TERMS ARE PLUS.

$$
\begin{array}{ll}
\text{1. Multiply} \quad . \;\; . \;\; . \;\; . & 3a^2 + 4ab + b^2 \\
\qquad \text{by} \quad . \;\; . \;\; . \;\; . \;\; . & \underline{2a + 5b \qquad\qquad\quad} \\
 & 6a^3 + 8a^2b + 2ab^2 \\
\text{The product, after reducing,} & \underline{\qquad + 15a^2b + 20ab^2 + 5b^3} \\
\qquad \text{becomes} \quad . \;\; . \;\; . \;\; . & 6a^3 + 23a^2b + 22ab^2 + 5b^3.
\end{array}
$$

44. NOTE.—It will be found convenient to *arrange* the terms of the polynomials with reference to some letter; that is, to write them down, so that the highest power of that letter shall enter the first term; the next highest, the second term, and so on to the last term.

---

44. How are the terms of a polynomial arranged with reference to a particular letter? What is this letter called? It the leading letter in the multiplicand and multiplier is the same, which will be the leading letter in the product?

The letter with reference to which the arrangement is made, is called the *leading letter.* In the above example the leading letter is a. The leading letter of the product will always be the same as that of the factors.

2. Multiply $x^2 + 2ax + a^2$ by $x + a$.
Ans. $x^3 + 3ax^2 + 3a^2x + a^3$.

3. Multiply $x^3 + y^3$ by $x + y$.
Ans. $x^4 + xy^3 + x^3y + y^4$.

4. Multiply $3ab^2 + 6a^2c^2$ by $3ab^2 + 3a^2c^2$.
Ans. $9a^2b^4 + 27a^3b^2c^2 + 18a^4c^4$.

5. Multiply $a^2b^2 + c^2d$ by $a + b$.
Ans. $a^3b^2 + ac^2d + a^2b^3 + bc^2d$.

6. Multiply $3ax^2 + 9ab^3 + cd^5$ by $6a^2c^2$.
Ans. $18a^3c^2x^2 + 54a^3c^2b^3 + 6a^2c^3d^5$.

7. Multiply $64a^3x^3 + 27a^2x + 9ab$ by $8a^3cd$.
Ans. $512a^6cdx^3 + 216a^5cdx + 72a^4bcd$

8. Multiply $a^3 + 3a^2x + 3ax^2 + x^3$ by $a + x$.
Ans. $a^4 + 4a^3x + 6a^2x^2 + 4ax^3 + x^4$.

9. Multiply $x^2 + y^2$ by $x + y$.
Ans. $x^3 + xy^2 + x^2y + y^3$.

10. Multiply $x^5 + xy^6 + 7ax$ by $ax + 5ax$.
Ans. $6ax^6 + 6ax^2y^6 + 42a^2x^2$

11. Multiply $a^3 + 3a^2b + 3ab^2 + b^3$ by $a + b$.
Ans. $a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4$.

12. Multiply $x^3 + x^2y + xy^2 + y^3$ by $x + y$.
Ans. $x^4 + 2x^3y + 2x^2y^2 + 2xy^3 + y^4$.

13. Multiply $x^3 + 2x^2 + x + 3$ by $3x + 1$.
Ans. $3x^4 + 7x^3 + 5x^2 + 10x + 3$.

GENERAL EXAMPLES.

$$\begin{array}{lrrr}
\text{1. Multiply} \ . \ . \ . \ . \ . \ . \ . \ . & 2ax - & 3ab & \\
\quad\text{by} \ . \ . \ . \ . \ . \ . \ . \ . \ . & 3x - & b. & \\
\hline
\text{The product} \ . \ . \ . \ . \ . \ . \ . \ . & 6ax^2 - & 9abx & \\
\quad\text{becomes after} \ . \ . \ . \ . \ . \ . & - & 2abx & + 3ab^2 \\
\hline
\quad\text{reducing} \ . \ . \ . \ \ . \ . \ . \ . & 6ax^2 - & 11abx & + 3ab^2.
\end{array}$$

2. Multiply $a^4 - 2b^3$ by $a - b$.
Ans. $a^5 - 2ab^3 - a^4b + 2b^4$.

3. Multiply $x^2 - 3x - 7$ by $x - 2$.
Ans. $x^3 - 5x^2 - x + 14$.

4. Multiply $3a^2 - 5ab + 2b^2$ by $a^2 - 7ab$.
Ans. $3a^4 - 26a^3b + 37a^2b^2 - 14ab^3$.

5. Multiply $b^2 + b^4 + b^6$ by $b^2 - 1$. *Ans.* $b^8 - b^2$.

6. Multiply $x^4 - 2x^3y + 4x^2y^2 - 8xy^3 + 16y^4$ by $x + 2y$.
Ans. $x^5 + 32y^5$.

7. Multiply $4x^2 - 2y$ by $2y$. *Ans.* $8x^2y - 4y^2$.

8. Multiply $2x + 4y$ by $2x - 4y$. *Ans.* $4x^2 - 16y^2$.

9. Multiply $x^3 + x^2y + xy^2 + y^3$ by $x - y$.
Ans. $x^4 - y^4$.

10. Multiply $x^2 + xy + y^2$ by $x^2 - xy + y^2$.
Ans. $x^4 + x^2y^2 + y^4$.

11. Multiply $2a^2 - 3ax + 4x^2$ by $5a^2 - 6ax - 2x^2$.
Ans. $10a^4 - 27a^3x + 34a^2x^2 - 18ax^3 - 8x^4$.

12. Multiply $3x^2 - 2xy + 5$ by $x^2 + 2xy - 3$.
Ans. $3x^4 + 4x^3y - 4x^2 - 4x^2y^2 + 16xy - 15$.

13. Multiply $3x^3 + 2x^2y^2 + 3y^2$ by $2x^3 - 3x^2y^2 + 5y^3$.

$$\textit{Ans.} \left\{ \begin{array}{l} 6x^6 - 5x^5y^2 - 6x^4y^4 + 6x^3y^2 + \\ 15x^3y^3 - 9x^2y^4 + 10x^2y^5 + 15y^5. \end{array} \right.$$

14. Multiply $8ax - 6ab - c$ by $2ax + ab + c$.
Ans. $16a^2x^2 - 4a^2bx - 6a^2b^2 + 6acx - 7abc - c^2$.

15. Multiply $3a^2 - 5b^2 + 3c^2$ by $a^2 - b^2$.

Ans. $3a^4 - 8a^2b^2 + 3a^2c^2 + 5b^4 - 3b^2c^2$.

16.

$$\begin{array}{r} 3a^2 - 5bd + cf \\ -\ 5a^2 + 4bd - 8cf. \\ \hline \end{array}$$

Pro. red. $-15a^4 + 37a^2bd - 29a^2cf - 20b^2d^2 + 44bcdf - 8c^2f^2$

17. Multiply $a^mx - a^2b^2$ by a^2x^n.

Ans. $a^{m+2}x^{n+1} - a^4b^2x^n$.

18. Multiply $a^m + b^n$ by $a^m - b^n$. *Ans.* $a^{2m} - b^{2n}$.

19. Multiply $a^m + b^n$ by $a^m + b^n$.

Ans. $a^{2m} + 2a^mb^n + b^{2n}$.

DIVISION.

45. DIVISION is the operation of finding from two quantities a third, which being multiplied by the second, will produce the first.

The first is called the *Dividend*, the second the *Divisor*, and the third, the *Quotient*.

Division is the converse of Multiplication. In it, we have given the product and one factor, to find the other. The rules for Division are just the converse of those for Multiplication.

To divide one monomial by another.

46. Divide $72a^5$ by $8a^3$. The division is indicated, thus:

$$\frac{72a^5}{8a^3}.$$

The quotient must be such a monomial, as, being *multiplied by the divisor*, will give the dividend. Hence, the coefficient

---

45. What is division? What is the first quantity called? The second? The third? What is given in division? What is required?

46. What is the rule for the division of monomials?

of the quotient must be 9, and the literal part a^2; for these quantities multiplied by $8a^3$ will give $72a^5$. Hence,

$$\frac{72a^5}{8a^3} = 9a^2.$$

The coefficient 9 is obtained by dividing 72 by 8; and the literal part is found by giving to a, an exponent equal to 5 minus 3.

Hence, for dividing one monomial by another, we have the following

RULE.

I. *Divide the coefficient of the dividend by the coefficient of the divisor, for a new coefficient:*

II. *After this coefficient write all the letters of the dividend, giving to each an exponent equal to the excess of its exponent in the dividend over that in the divisor.*

SIGNS IN DIVISION.

47. Since the Quotient multiplied by the Divisor must produce the Dividend: and, since the product of two factors having the *same sign* will be + ; and the product of two factors having different signs will be − ; we conclude:

1. When the signs of the dividend and divisor are like, the sign of the quotient will be +.

2. When the signs of the dividend and divisor are unlike, the sign of the quotient will be −. Again, for brevity, we say,

+ divided by +, and − divided by −, give + ;
− divided by +, and + divided by −, give −.

$$\frac{+ab}{+a} = +b; \qquad \frac{-ab}{-b} = +a;$$

$$\frac{-ab}{+a} = -b; \qquad \frac{+ab}{-a} = -b.$$

---

47. What is the rule for the signs, in division?

EXAMPLES.

(1.)

$$\frac{+\ 18a^5b^2c}{+\ 9a^3bc} = +\ 2a^2b.$$

(2)

$$\frac{-\ 15a^3x^2y}{-\ 5a^2x} = +\ 3axy.$$

(3.)

$$\frac{-\ 24a^4bc}{+\ 3abc} = -\ 8a^3.$$

(4.)

$$\frac{32a^8b^3x^5}{-\ 8a^7b^3x} = -\ 4ax^4.$$

5. Divide $15ax^2y^3$ by $-\ 3ay$. *Ans.* $-\ 5x^2y^2$.
6. Divide $84ab^3x$ by $12b^2$. *Ans.* $7abx$.
7. Divide $-\ 36a^4b^5c^2$ by $9a^3b^2c$. *Ans.* $-\ 4ab^3c$.
8. Divide $-\ 99a^4b^4x^5$ by $11a^3b^2x^4$. *Ans.* $-\ 9ab^2x$.
9. Divide $108x^6y^5z^3$ by $54x^5z$. *Ans.* $2xy^5z^2$.
10. Divide $64x^7y^5z^6$ by $-\ 16x^6y^4z^5$. *Ans.* $-\ 4xyz$.
11. Divide $-\ 96a^7b^6c^5$ by $12a^2bc$. *Ans.* $-\ 8a^5b^5c^4$.
12. Divide $-\ 38a^4b^6d^4$ by $2a^3b^5d$. *Ans.* $-\ 19abd^3$.
13. Divide $-\ 64a^5b^4c^8$ by $32a^4bc$. *Ans.* $-\ 2ab^3c^7$.
14. Divide $128a^5x^6y^7$ by $16axy^4$. *Ans.* $8a^4x^5y^3$.
15. Divide $-\ 256a^4b^9c^8d^7$ by $16a^3bc^6$. *Ans.* $-\ 16ab^8c^2d^7$.
16. Divide $200a^8m^2n^2$ by $-\ 50a^7mn$. *Ans.* $-\ 4amn$.
17. Divide $300x^3y^4z^2$ by $60xy^2z$. *Ans.* $5x^2y^2z$.
18. Divide $27a^5b^2c^2$ by $-\ 9abc$. *Ans.* $-\ 3a^4bc$.
19. Divide $64a^3y^6z^8$ by $32ay^5z^7$. *Ans.* $2a^2yz$.
20. Divide $-\ 88a^5b^6c^8$ by $11a^3b^4c^6$. *Ans.* $-\ 8a^2b^2c^2$.
21. Divide $77a^4y^3z^4$ by $-\ 11a^4y^3z^4$. *Ans.* $-\ 7$.
22. Divide $84a^4b^2c^2d$ by $-\ 42a^4b^2c^2d$. *Ans.* $-\ 2$.
23. Divide $-\ 88a^6b^7c^6$ by $8a^5b^6c^6$. *Ans.* $-\ 11ab$.
24. Divide $16x^2$ by $-\ 8x$. *Ans.* $-\ 2x$.
25. Divide $-\ 88a^nb^2$ by $11a^mb$. *Ans.* $-\ 8a^{n-m}b$.

26. Divide $77a^mb^n$ by $-11a^2b^3$. *Ans.* $-7a^{m-2}b^{n-3}$.

27. Divide $84a^8b^m$ by $42a^nb^9$. *Ans.* $2a^{8-n}b^{m-9}$.

28. Divide $-88a^pb^q$ by $8a^nb^m$. *Ans.* $-11a^{p-n}b^{q-m}$.

29. Divide $96ab^p$ by $48a^nb^q$. *Ans.* $2a^{1-n}b^{p-q}$.

30. Divide $168x^ay^b$ by $12x^ny^m$. *Ans.* $14x^{a-n}y^{b-m}$.

31. Divide $256ab^3c^3$ by $16a^nb^mc^p$. *Ans.* $16a^{1-n}b^{3-m}c^{3-p}$.

MONOMIAL FRACTIONS.

48. It follows from the preceding rules, that the exact division of monomials will be impossible:

1st. When the coefficient of the dividend is not exactly divisible by that of the divisor.

2d. When the exponent of the same letter is greater in the divisor than in the dividend.

3d. When the divisor contains one or more letters not found in the dividend.

In either case, the quotient will be expressed by a fraction.

A fraction is said to be in its *simplest form*, when the numerator and denominator do not contain a common factor.

For example, $12a^4b^2cd$, divided by $8a^2bc^2$, gives

$$\frac{12a^4b^2d}{8a^2bc^2};$$

which may be reduced by dividing the numerator and denominator by the common factors, 4, a^2, b, and c, giving

$$\frac{12a^4b^2cd}{8a^2bc^2} = \frac{3a^2bd}{2c}.$$

Also, $$\frac{25a^5b^2d^3}{15a^4b^6d^4} = \frac{5a}{3b^4d}.$$

---

48. Under what circumstances will the division of monomials be impossible? How will the quantities then be expressed? How is a monomial fraction reduced to its simplest form?

Hence, for the reduction of a monomial fraction to its simplest form, we have the following

RULE.

Suppress every factor, whether numerical or literal, that is common to both terms of the fraction; the result will be the reduced fraction sought.

EXAMPLES.

(1.) $$\frac{48a^3b^2cd^3}{36a^2b^3c^2de} = \frac{4ad^2}{3bce}:$$ and (2.) $$\frac{37ab^3c^5d}{6a^3bc^4d^2} = \frac{37b^2c}{6a^2d};$$

also, (3.) $$\frac{7a^2b}{14a^3b^2} = \frac{1}{2ab}:$$ and (4.) $$\frac{4a^2b^2}{6ab^4} = \frac{2a}{3b^2}.$$

5. Divide $49a^2b^2c^6$ by $14a^3bc^4$. *Ans.* $\frac{7bc^2}{2a}$.

6. Divide $6amn$ by $3abc$. *Ans.* $\frac{2mn}{bc}$.

7. Divide $18a^2b^2mn^2$ by $12a^4b^4cd$. *Ans.* $\frac{3mn^2}{2a^2b^2cd}$.

8. Divide $28a^5b^6c^7d^8$ by $16ab^9cd^7m$. *Ans.* $\frac{7a^4c^6d}{4b^3m}$.

9. Divide $72a^3c^2b^2$ by $12a^5c^4b^3d$. *Ans.* $\frac{6}{a^2c^2bd}$.

10. Divide $100a^8b^5xmn$ by $25a^3b^4d$. *Ans.* $\frac{4a^5bxmn}{d}$.

11. Divide $96a^5b^8c^9df$ by $75a^2cxy$. *Ans.* $\frac{32a^3b^8c^8df}{25xy}$.

12. Divide $85m^2n^3fx^2y^3$ by $15am^4nf$. *Ans.* $\frac{17n^2x^2y^3}{3am^2}$.

13. Divide $127d^3x^2y^2$ by $16d^4x^4y^4$. *Ans.* $\frac{127}{16dx^2y^2}$.

49. In dividing monomials, it often happens that the exponents of the same letter, in the dividend and divisor, are equal; in which case that letter may not appear in the quotient. It might, however, be retained by giving to it the exponent 0.

If we have expressions of the form

$$\frac{a}{a}, \quad \frac{a^2}{a^2}, \quad \frac{a^3}{a^3}, \quad \frac{a^4}{a^4}, \quad \frac{a^5}{a^5}, \quad \&c.,$$

and apply the rule for the exponents, we shall have,

$$\frac{a}{a} = a^{1-1} = a^0, \quad \frac{a^2}{a^2} = a^{2-2} = a^0, \quad \frac{a^3}{a^3} = a^{3-3} = a^0, \quad \&c.$$

But since any quantity divided by itself is equal to 1, it follows that,

$$\frac{a}{a} = a^0 = 1, \quad \frac{a^2}{a^2} = a^{2-2} = a^0 = 1, \quad \&c.;$$

or, finally, if we designate the exponent by m, we have,

$$\frac{a^m}{a^m} = a^{m-m} = a^0 = 1; \text{ that is,}$$

The 0 *power of any quantity is equal to* 1 : therefore,

Any quantity may be retained in a term, or introduced into a term, by giving it the exponent 0.

EXAMPLES.

1. Divide $6a^2b^2c^4$ by $2a^2b^2$.

$$\frac{6a^2b^2c^4}{2a^2b^2} = 3a^{2-2}b^{2-2}c^4 = 3a^0b^0c^4 = 3c^4.$$

2. Divide $8a^4b^3c^5$ by $-4a^4b^3c$. *Ans.* $-2a^0b^0c^4 = -2c^4$.

3. Divide $-32m^3n^2x^2y^2$ by $4m^3n^2xy$.

Ans. $-8m^0n^0xy = -8xy$.

---

49. When the exponents of the same letter in the dividend and divisor are equal, what takes place? May the letter still be retained? With what exponent? What is the zero power of any quantity equal to?

4. Divide $-96a^4b^5c^n$ by $-24a^4b^5$. *Ans.* $4a^0b^0c^n = 4c^n$

5. Introduce a, as a factor, into $6b^5c^4$. *Ans.* $6a^0b^5c^4$.

6. Introduce ab, as factors, into $9c^5d^n$. *Ans.* $9a^0b^0c^5d^n$.

7. Introduce abc, as factors, into $8d^4f^m$. *A.* $8a^0b^0c^0d^4f^m$.

50. When the exponent of any letter is greater in the divisor than it is in the dividend, the exponent of that letter in the quotient may be written with a negative sign. Thus,

$$\frac{a^2}{a^5} = \frac{1}{a^3}; \text{ also, } \frac{a^2}{a^5} = a^{2-5} = a^{-3}, \text{ by the rule;}$$

hence,

$$a^{-3} = \frac{1}{a^3}.$$

Since, $a^{-3} = \frac{1}{a^3}$, we have, $b \times a^{-3} = \frac{b}{a^3}$;

that is, a in the numerator, with a negative exponent, is equal to a in the denominator, with an equal positive exponent; hence,

Any quantity having a negative exponent, is equal to the reciprocal of the same quantity with an equal positive exponent.

Hence, also,

Any factor may be transferred from the denominator to the numerator of a fraction, or the reverse, by changing the sign of its exponent.

EXAMPLES.

1. Divide $32a^2bc$ by $16a^5b^2$.

Ans. $\frac{32a^2bc}{16a^5b^2} = 2a^{-3}b^{-1}c = \frac{2c}{a^3b}$.

---

50. When the exponent of any letter in the divisor is greater than in the dividend, how may the exponent of that letter be written in the quotient? What is a quantity with a negative exponent equal to? How may a factor be transferred from the numerator to the denominator of a fraction?

2. $\frac{54a^2b^3c}{9a^4b^5} = 6a^{-2}b^{-2}c.$ *Ans.* $\frac{6c}{a^2b^2}$.

3. Reduce $\frac{17x^2y^3z}{51x^4y^3}$. *Ans.* $\frac{x^{-2}z}{3}$, or $\frac{z}{3x^2}$.

4. In $5ay^{-3}x^{-2}$, get rid of the negative exponents.

Ans. $\frac{5a}{y^3x^2}$.

5. In $\frac{4a^2b^3x^{-2}}{3a^{-3}b^{-5}}$, get rid of the negative exponents.

Ans. $\frac{4a^5b^8}{3x^2}$.

6. In $\frac{15a^{-3}c^{-4}d^{-5}}{45x^{-3}y^{-5}c^{-2}}$, get rid of the negative exponents.

Ans. $\frac{x^3y^5}{3a^3c^2d^5}$.

7. Reduce $\frac{-8a^{-3}b^5c}{14a^2b^{-5}c}$. *Ans.* $\frac{-4a^{-5}b^{10}c^0}{7}$, or $\frac{-4b^{10}}{7a^5}$.

8. Reduce $72a^5b^2 \div 8a^6b^3$. *Ans.* $9a^{-1}b^{-1}$, or $\frac{9}{ab}$.

9. In $\frac{15a^{-4}b^6c^{-1}}{5a^{-2}b^{-1}}$, get rid of the negative exponents.

Ans. $\frac{3b^7}{a^2c}$.

10. Reduce $\frac{-15a^{-5}b^{-5}c^2}{-5a^{-6}b^{-7}}$. *Ans.* $3ab^2c^2$.

To divide a polynomial by a monomial.

51. To divide a polynomial by a monomial:

Divide each term of the dividend, separately, by the divisor; the algebraic sum of the quotients will be the quotient sought.

EXAMPLES.

1. Divide $3a^2b^2 - a$ by a. *Ans.* $3ab^2 - 1$.

---

51. How do you divide a polynomial by a monomial?

2. Divide $5a^3b^2 - 25a^4b^2$ by $5a^3b^2$. *Ans.* $1 - 5a$.

3. Divide $35a^2b^2 - 25ab$ by $-5ab$. *Ans.* $-7ab + 5$

4. Divide $10ab - 15ac$ by $5a$. *Ans.* $2b - 3c$.

5. Divide $6ab - 8ax + 4a^2y$ by $2a$.
Ans. $3b - 4x + 2ay$.

6. Divide $-15ax^2 + 6x^3$ by $-3x$. *Ans.* $5ax - 2x^2$.

7. Divide $-21xy^2 + 35a^2b^2y - 7c^2y$ by $-7y$.
Ans. $3xy - 5a^2b^2 + c^2$.

8. Divide $40a^8b^4 + 8a^4b^7 - 32a^4b^4c^4$ by $8a^4b^4$.
Ans. $5a^4 + b^3 - 4c^4$

DIVISION OF POLYNOMIALS.

52. 1. Divide $-2a + 6a^2 - 8$ by $2 + 2a$.

$$\begin{array}{lll}
\textit{Dividend.} & \textit{Divisor.} & \\
6a^2 - 2a - 8 & \underline{|\ 2a + 2} & \\
\underline{6a^2 + 6a} & 3a - 4 & \textit{Quotient.} \\
\quad -8a - 8 & & \\
\quad \underline{-8a - 8} & & \\
\qquad 0 & \textit{Remainder.} &
\end{array}$$

We first *arrange* the dividend and divisor with reference to a (Art. 44), placing the divisor on the left of the dividend. Divide the first term of the dividend by the first term of the divisor; the result will be the first term of the quotient, which, for convenience, we place under the divisor. The product of the divisor by this term ($6a^2 + 6a$), being subtracted from the dividend, leaves a new dividend, which may be treated in the same way as the original one, and so on to the end of the operation.

---

52. What is the rule for dividing one polynomial by another? When is the division exact? When is it not exact?

Since all similar cases may be treated in the same way, we have, for the division of polynomials, the following

RULE.

I. *Arrange the dividend and divisor with reference to the same letter:*

II. *Divide the first term of the dividend by the first term of the divisor, for the first term of the quotient. Multiply the divisor by this term of the quotient, and subtract the product from the dividend:*

III. *Divide the first term of the remainder by the first term of the divisor, for the second term of the quotient. Multiply the divisor by this term, and subtract the product from the first remainder, and so on:*

IV. *Continue the operation, until a remainder is found equal to* 0, *or one whose first term is not divisible by that of the divisor.*

NOTE.—1. When a remainder is found equal to 0, the division is exact.

2. When a remainder is found whose first term is not divisible by the first term of the divisor, the exact division is impossible. In that case, write the last remainder after the quotient found, placing the divisor under it, in the form of a fraction.

SECOND EXAMPLE.

Let it be required to divide

$51a^2b^2 + 10a^4 - 48a^3b - 15b^4 + 4ab^3$ by $4ab - 5a^2 + 3b^2$.

We first arrange the dividend and divisor with reference to a.

$$\begin{array}{r|l}
\textit{Dividend.} & \textit{Divisor.} \\
10a^4 - 48a^3b + 51a^2b^2 + 4ab^3 - 15b^4 & -5a^2 + 4ab + 3b^2 \\ \cline{2-2}
+10a^4 - 8a^3b - 6a^2b^2 & -2a^2 + 8ab - 5b^2 \\ \cline{1-1}
-40a^3b + 57a^2b^2 + 4ab^3 - 15b^4 & \quad \textit{Quotient.} \\
-40a^3b + 32a^2b^2 + 24ab^3 & \\ \cline{1-1}
25a^2b^2 - 20ab^3 - 15b^4 & \\
25a^2b^2 - 20ab^3 - 15b^4 &
\end{array}$$

(3.)

$$\begin{array}{l|l}
x^4 + x^3y + x^2y + xy^2 - 2y & x + y \\ \cline{2-2}
x^4 + x^3y & x^3 + xy \qquad \dfrac{2y}{x + y}. \\ \cline{1-1}
\qquad + x^2y + xy^2 & \\
\qquad + x^2y + xy^2 & \\ \cline{1-1}
\qquad\qquad\qquad - 2y &
\end{array}$$

Here the division is not exact, and the quotient is fractional.

(4.)

$$\begin{array}{l|l}
1 + a & 1 - a \\ \cline{2-2}
1 - a & 1 + 2a + 2a^2 + 2a^3 +, \text{\&c.} \\ \cline{1-1}
\;\; + 2a & \\
\;\; + 2a - 2a^2 & \\ \cline{1-1}
\qquad\quad + 2a^2 & \\
\qquad\quad + 2a^2 - 2a^3 & \\ \cline{1-1}
\qquad\qquad\quad + 2a^3 &
\end{array}$$

In this example the operation does not terminate. It may be continued to any extent.

EXAMPLES.

1. Divide $a^2 + 2ax + x^2$ by $a + x$. *Ans.* $a + x$.

2. Divide $a^3 - 3a^2y + 3ay^2 - y^3$ by $a - y$. *Ans.* $a^2 - 2ay + y^2$.

3. Divide $24a^2b - 12a^3cb^2 - 6ab$ by $-6ab$.
Ans. $-4a + 2a^2cb + 1$.

4. Divide $6x^4 - 96$ by $3x - 6$.
Ans. $2x^3 + 4x^2 + 8x + 16$.

5. Divide $a^5 - 5a^4x + 10a^3x^2 - 10a^2x^3 + 5ax^4 - x^5$ by $a^2 - 2ax + x^2$.
Ans. $a^3 - 3a^2x + 3ax^2 - x^3$.

6. Divide $48x^3 - 76ax^2 - 64a^2x + 105a^3$ by $2x - 3a$.
Ans. $24x^2 - 2ax - 35a^2$.

7. Divide $y^6 - 3y^4x^2 + 3y^2x^4 - x^6$ by $y^3 - 3y^2x + 3yx^2 - x^3$.
Ans. $y^3 + 3y^2x + 3yx^2 + x^3$.

8. Divide $64a^4b^6 - 25a^2b^8$ by $8a^2b^3 + 5ab^4$.
Ans. $8a^2b^3 - 5ab^4$.

9. Divide $6a^3 + 23a^2b + 22ab^2 + 5b^3$ by $3a^2 + 4ab + b^2$.
Ans. $2a + 5b$.

10. Divide $6ax^6 + 6ax^2y^6 + 42a^2x^2$ by $ax + 5ax$.
Ans. $x^5 + xy^6 + 7ax$.

11. Divide $-15a^4 + 37a^2bd - 29a^2cf - 20b^2d^2 + 44bcdf - 8c^2f^2$ by $3a^2 - 5bd + cf$.
Ans. $-5a^2 + 4bd - 8cf$.

12. Divide $x^4 + x^2y^2 + y^4$ by $x^2 - xy + y^2$.
Ans. $x^2 + xy + y^2$.

13. Divide $x^4 - y^4$ by $x - y$.
Ans. $x^3 + x^2y + xy^2 + y^3$.

14. Divide $3a^4 - 8a^2b^2 + 3a^2c^2 + 5b^4 - 3b^2c^2$ by $a^2 - b^2$
Ans. $3a^2 - 5b^2 + 3c^2$.

15. Divide $6x^6 - 5x^5y^2 - 6x^4y^4 + 6x^3y^2 + 15x^3y^3 - 9x^2y^4 + 10x^2y^5 + 15y^5$ by $3x^3 + 2x^2y^2 + 3y^2$.
Ans. $2x^3 - 3x^2y^2 + 5y^3$.

16. Divide $-c^2 + 16a^2x^2 - 7abc - 4a^2bx - 6a^2b^2 + 6acx$ by $8ax - 6ab - c$.
Ans. $2ax + ab + c$.

17. Divide $3x^4 + 4x^3y - 4x^2 - 4x^2y^2 + 16xy - 15$ by $2xy + x^2 - 3$.
Ans. $3x^2 - 2xy + 5$.

18. Divide $x^5 + 32y^5$ by $x + 2y$.

Ans. $x^4 - 2x^3y + 4x^2y^2 - 8xy^3 + 16y^4$.

19. Divide $3a^4 - 26a^3b - 14ab^3 + 37a^2b^2$ by $2b^2 - 5ab + 3a^2$.

Ans. $a^2 - 7ab$.

20. Divide $a^4 - b^4$ by $a^3 + a^2b + ab^2 + b^3$.

Ans. $a - b$

21. Divide $x^3 - 3x^2y + y^3$ by $x + y$.

Ans. $x^2 - 4xy + 4y^2 - \frac{3y^3}{x + y}$.

22. Divide $1 + 2a$ by $1 - a - a^2$.

Ans. $1 + 3a + 4a^2 + 7a^3 +$, &c.

CHAPTER III.

USEFUL FORMULAS. FACTORING. GREATEST COMMON DIVISOR. LEAST COMMON MULTIPLE.

USEFUL FORMULAS.

53. A FORMULA is an algebraic expression of a general rule, or principle.

Formulas serve to shorten algebraic operations, and are also of much use in the operation of factoring. When translated into common language, they give rise to practical rules.

The verification of the following formulas affords additional exercises in Multiplication and Division.

(1.)

54. To form the square of $a + b$, we have,

$$(a + b)^2 = (a + b)(a + b) = a^2 + 2ab + b^2.$$

That is,

The square of the sum of any two quantities is equal to the square of the first, plus twice the product of the first by the second, plus the square of the second.

1. Find the square of $2a + 3b$. We have from the rule,

$$(2a + 3b)^2 = 4a^2 + 12ab + 9b^2.$$

---

53. What is a formula? What are the uses of formulas?

54. What is the square of the sum of two quantities equal to?

2. Find the square of $5ab + 3ac$.

Ans. $25a^2b^2 + 30a^2bc + 9a^2c^2$.

3. Find the square of $5a^2 + 8a^2b$.

Ans. $25a^4 + 80a^4b + 64a^4b^2$.

4. Find the square of $6ax + 9a^2x^2$.

Ans. $36a^2x^2 + 108a^3x^3 + 81a^4x^4$

(2.)

55. To form the square of a difference, $a - b$, we have,

$$(a - b)^2 = (a - b)(a - b) = a^2 - 2ab + b^2.$$

That is,

The square of the difference of any two quantities is equal to the square of the first, minus twice the product of the first by the second, plus the square of the second.

1. Find the square of $2a - b$. We have,

$$(2a - b)^2 = 4a^2 - 4ab + b^2.$$

2. Find the square of $4ac - bc$.

Ans. $16a^2c^2 - 8abc^2 + b^2c^2$.

3. Find the square of $7a^2b^2 - 12ab^3$.

Ans. $49a^4b^4 - 168a^3b^5 + 144a^2b^6$.

(3.)

56. Multiply $a + b$ by $a - b$. We have,

$$(a + b) \times (a - b) = a^2 - b^2.$$ Hence,

The sum of two quantities, multiplied by their difference, is equal to the difference of their squares.

1. Multiply $2c + b$ by $2c - b$. Ans. $4c^2 - b^2$

2. Multiply $9ac + 3bc$ by $9ac - 3bc$.

Ans. $81a^2c^2 - 9b^2c^2$

---

55. What is the square of the difference of two quantities equal to?

56. What is the sum of two quantities multiplied by their difference equal to?

3. Multiply $8a^3 + 7ab^2$ by $8a^3 - 7ab^2$.

Ans. $64a^6 - 49a^2b^4$.

(**4.**)

57 Multiply $a^2 + ab + b^2$ by $a - b$. We have,

$$(a^2 + ab + b^2)(a - b) = a^3 - b^3.$$

(**5.**)

58. Multiply $a^2 - ab + b^2$ by $a + b$. We have,

$$(a^2 - ab + b^2)(a + b) = a^3 + b^3.$$

(**6.**)

59. Multiply together, $a + b$, $a - b$, and $a^2 + b^2$. We have,

$$(a + b)(a - b)(a^2 + b^2) = a^4 - b^4.$$

60. Since every product is divisible by any of its factors, each formula establishes the principle set opposite its number.

1. *The sum of the squares of any two quantities, plus twice their product, is divisible by their sum.*

2. *The sum of the squares of any two quantities, minus twice their product, is divisible by the difference of the quantities.*

3. *The difference of the squares of any two quantities is divisible by the sum of the quantities, and also by their difference.*

4. *The difference of the cubes of any two quantities is divisible by the difference of the quantities; also, by the sum of their squares, plus their product.*

5. *The sum of the cubes of any two quantities is divisi-*

---

60. By what is any product divisible? By applying this principle, what follows from Formula (1)? What from (2)? What from (3)? What from (4)? What from (5)? What from (6)?

ble by the sum of the quantities; also, by the sum of their squares minus their product.

6. *The difference between the fourth powers of any two quantities is divisible by the sum of the quantities, by their difference, by the sum of their squares, and by the difference of their squares.*

FACTORING.

61. Factoring is the operation of resolving a quantity into factors. The principles employed are the converse of those of Multiplication. The operations of factoring are performed by inspection.

1. What are the factors of the polynomial

$$ac + ab + ad.$$

We see, by inspection, that a is a common factor of all the terms; hence, it may be placed without a parenthesis, and the other parts within; thus:

$$ac + ab + ad = a(c + b + d).$$

2. Find the factors of the polynomial $a^2b^2 + a^2d - a^2f$.
Ans. $a^2(b^2 + d - f)$.

3. Find the factors of the polynomial $3a^2b - 6a^2b^2 + b^2d$.
Ans. $b(3a^2 - 6a^2b + bd)$.

4. Find the factors of $3a^2b - 9a^2c - 18a^2xy$.
Ans. $3a^2(b - 3c - 6xy)$.

5. Find the factors of $8a^2cx - 18acx^2 + 2ac^5y - 30a^6c^9$.
Ans. $2ac(4ax - 9x^2 + c^4y - 15a^5c^8)$.

6. Factor $30a^4b^2c - 6a^3b^2d^3 + 18a^3b^2c^2$.
Ans. $6a^3b^2(5ac - d^3 + 3c^2)$.

7. Factor $12c^4bd^3 - 15c^3d^4 - 6c^2d^3f$.
Ans. $3c^2d^3(4c^2b - 5cd - 2f)$.

---

61. What is factoring?

8. Factor $15a^3bcf - 10abc^4 - 25abcd$.
Ans. $5abc(3a^2f - 2c^3 - 5d)$.

62. When two terms of a trinomial are squares, and positive, and the third term is equal to twice the product of their square roots, the trinomial may be resolved into factors by Formula (**1**).

1. Factor $a^2 + 2ab + b^2$ *Ans.* $(a + b)(a + b)$.

2. Factor $4a^2 + 12ab + 9b^2$.
Ans. $(2a + 3b)(2a + 3b)$.

3. Factor $9a^2 + 12ab + 4b^2$.
Ans. $(3a + 2b)(3a + 2b)$.

4. Factor $4x^2 + 8x + 4$. *Ans.* $(2x + 2)(2x + 2)$.

5. Factor $9a^2b^2 + 12abc + 4c^2$.
Ans. $(3ab + 2c)(3ab + 2c)$.

6. Factor $16x^2y^2 + 16xy^3 + 4y^4$.
Ans. $(4xy + 2y^2)(4xy + 2y^2)$.

63. When two terms of a trinomial are squares, and positive, and the third term is equal to *minus* twice their square roots, the trinomial may be factored by Formula (**2**).

1. Factor $a^2 - 2ab + b^2$. *Ans.* $(a - b)(a - b)$.

2. Factor $4a^2 - 4ab + b^2$. *Ans.* $(2a - b)(2a - b)$.

3. Factor $9a^2 - 6ac + c^2$. *Ans.* $(3a - c)(3a - c)$.

4. Factor $a^2x^2 - 4ax + 4$. *Ans.* $(ax - 2)(ax - 2)$

5. Factor $4x^2 - 4xy + y^2$. *Ans.* $(2x - y)(2x - y)$.

---

62. When may a trinomial be factored?

63. When may a trinomial be factored by this method?

6. Factor $36x^2 - 24xy + 4y^2$.
Ans. $(6x - 2y)(6x - 2y)$.

64. When the two terms of a binomial are squares and have contrary signs, the binomial may be factored by Formula (**3**).

1. Factor $4c^2 - b^2$. *Ans.* $(2c + b)(2c - b)$

2. Factor $81a^2c^2 - 9b^2c^2$.
Ans. $(9ac + 3bc)(9ac - 3bc)$.

3. Factor $64a^4b^4 - 25x^2y^2$.
Ans. $(8a^2b^2 + 5xy)(8a^2b^2 - 5xy)$.

4. Factor $25a^2c^2 - 9x^4y^2$.
Ans. $(5ac + 3x^2y)(5ac - 3x^2y)$.

5. Factor $36a^4b^4c^2 - 9x^6$.
Ans. $(6a^2b^2c + 3x^3)(6a^2b^2c - 3x^3)$.

6. Factor $49x^4 - 36y^4$. *Ans.* $(7x^2 + 6y^2)(7x^2 - 6y^2)$.

65. When the two terms of a binomial are cubes, and have contrary signs, the binomial may be factored by Formula (**4**).

1. Factor $8a^3 - c^3$. *Ans.* $(2a - c)(4a^2 + 2ac + c^2)$.

2. Factor $27a^3 - 64$.
Ans. $(3a - 4)(9a^2 + 12a + 16)$.

3. Factor $a^3 - 64b^3$.
Ans. $(a - 4b)(a^2 + 4ab + 16b^2)$.

4. Factor $a^3 - 27b^3$. *Ans.* $(a - 3b)(a^2 + 3ab + 9b^2)$.

---

64. When may a binomial be factored?

65. When may a binomial be factored by this method?

66. When the terms of a binomial are cubes and have like signs, the binomial may be factored by Formula (**5**).

1. Factor $8a^3 + c^3$. *Ans.* $(2a + c)(4a^2 - 2ac + c^2)$.

2. Factor $27a^3 + 64$.
Ans. $(3a + 4)(9a^2 - 12a + 16)$.

3. Factor $a^3 + 64b^3$.
Ans. $(a + 4b)(a^2 - 4ab + 16b^2)$.

4. Factor $a^3 + 27b^3$. *Ans.* $(a + 3b)(a^2 - 3ab + 9b^2)$.

67. When the terms of a binomial are 4th powers, and have contrary signs, the binomial may be factored by Formula (**6**).

1. What are the factors of $a^4 - b^4$?
Ans. $(a + b)(a - b)(a^2 + b^2)$.

2. What are the factors of $81a^4 - 16b^4$?
Ans. $(3a + 2b)(3a - 2b)(9a^2 + 4b^2)$.

3. What are the factors of $16a^4b^4 - 81c^4d^4$?
Ans. $(2ab + 3cd)(2ab - 3cd)(4a^2b^2 + 9c^2d^2)$.

GREATEST COMMON DIVISOR.

68. A COMMON DIVISOR of two quantities, is a quantity that will divide them both without a remainder. Thus, $3a^2b$, is a common divisor of $9a^2b^2c$ and $3a^2b^2 - 6a^3b^3$.

---

66. When may a binomial be factored by this method?
67. When may a binomial be factored by this method?
68. What is the common divisor of two quantities?

69. A SIMPLE or PRIME FACTOR is one that cannot be resolved into any other factors.

Every prime factor, common to two quantities, is a common divisor of those quantities. The continued product of any number of prime factors, common to two quantities, is also a common divisor of those quantities.

70. The GREATEST COMMON DIVISOR of two quantities, is the continued product of all the prime factors which are common to both.

71. When both quantities can be resolved into prime factors, by the method of factoring already given, the greatest common divisor may be found by the following

RULE.

I. *Resolve both quantities into their prime factors :*

II. *Find the continued product of all the factors which are common to both ; it will be the greatest common divisor required.*

EXAMPLES.

1. Required the greatest common divisor of $75a^2b^2c$ and $25abd$. Factoring, we have,

$$75a^2b^2c = 3 \times 5 \times 5aabbc$$
$$25abd = 5 \times 5abd.$$

The factors, 5, 5, a and b, are common; hence,

$$5 \times 5 \times a \times b = 25ab,$$

is the divisor sought.

---

69. What is a simple or prime factor? Is a prime factor, common to two quantities, a common divisor?

70. What is the greatest common divisor?

71. If both quantities can be resolved into prime factors, how do you find the greatest common divisor?

VERIFICATION.

$$75a^2b^2c \div 25ab = 3abc$$
$$25abd \div 25ab = d;$$

and since the quotients have no common factor, they cannot be further divided.

2. Required the greatest common divisor of $a^2 - 2ab + b^2$ and $a^2 - b^2$. *Ans.* $a - b$

3. Required the greatest common divisor of $a^2 + 2ab + b^2$ and $a + b$. *Ans.* $a + b$

4. Required the greatest common divisor of $a^2x^2 - 4ax + 4$ and $ax - 2$. *Ans.* $ax - 2$.

5. Find the greatest common divisor of $3a^2b - 9a^2c - 18a^2xy$ and $b^2c - 3bc^2 - 6bcxy$. *Ans.* $b - 3c - 6xy$.

6. Find the greatest common divisor of $4a^2c - 4acx$ and $3a^2g - 3agx$. *Ans.* $a(a - x)$, or $a^2 - ax$.

7. Find the greatest common divisor of $4c^2 - 12cx + 9x^2$ and $4c^2 - 9x^2$. *Ans.* $2c - 3x$.

8. Find the greatest common divisor of $x^3 - y^3$ and $x^2 - y^2$. *Ans.* $x - y$.

9. Find the greatest common divisor of $4c^2 + 4bc + b^2$ and $4c^2 - b^2$. *Ans.* $2c + b$.

10. Find the greatest common divisor of $25a^2c^3 - 9x^4y^4$ and $5acd^2 + 3d^2x^2y^2$. *Ans.* $5ac + 3x^2y^2$.

NOTE.—To find the greatest common divisor of three quantities. First find the greatest common divisor of two of them, and then the greatest common divisor between this result and the third.

1. What is the greatest common divisor of $4ax^2y$, $16abx^2$, and $24acx^2$? *Ans.* $4ax^2$.

2. Of $3x^2 - 6x$, $2x^3 - 4x^2$, and $x^2y - 2xy$? *Ans.* $x - 2x$.

---

72. When is one quantity a multiple of another?

LEAST COMMON MULTIPLE.

72. One quantity is a MULTIPLE of another, when it can be divided by that other without a remainder. Thus, $8a^2b$, is a multiple of 8, also of a^2, and of b.*

73. A quantity is a *Common Multiple* of two or more quantities, when it can be divided by each, separately, without a remainder. Thus, $24a^3x^3$, is a common multiple of $6ax$ and $4a^2x$.

74. The LEAST COMMON MULTIPLE of two or more quantities, is the simplest quantity that can be divided by each, without a remainder. Thus, $12a^2b^2x^2$, is the least common multiple of $2a^2x$, $4ab^2$, and $6a^2b^2x^2$.

75. Since the common multiple is a dividend of each of the quantities, and since the division is exact, the common multiple must contain every prime factor in all the quantities; and if the same factor enters more than once, it must enter an equal number of times into the common multiple.

When the given quantities can be factored, by any of the methods already given, the least common multiple may be found by the following

RULE.

I. *Resolve each of the quantities into its prime factors:*

II. *Take each factor as many times as it enters any one of the quantities, and form the continued product of these factors; it will be the least common multiple.*

---

73. When is a quantity a common multiple of several others?

74. What is the least common multiple of two or more quantities?

75. What does the common multiple of two or more quantities contain, as factors? How may the least common multiple be found?

* The *multiple* of a quantity, is simply a *dividend* which will give an exact quotient.

EXAMPLES.

1. Find the least common multiple of $12a^3b^2c^2$ and $8a^2b^3$.

$$12a^3b^2c^2 = 2.2.3.aaabbcc.$$
$$8a^2b^3 = 2.2.2.aabbb.$$

Now, since 2 enters 3 times as a factor, it must enter 3 times in the common multiple: 3 must enter once; a, 3 times; b, 3 times; and c, twice; hence,

$$2.2.2.3aaabbbcc = 24a^3b^3c^2,$$

is the least common multiple.

Find the least common multiples of the following:

2. $6a$, $5a^2b$, and $25abc^2$. *Ans.* $150a^2bc^2$

3. $3a^2b$, $9abc$, and $27a^2x^2$. *Ans.* $27a^2bcx^2$.

4. $4a^2x^2y^2$, $8a^3xy$, $16a^4y^3$, and $24a^5y^4x$. *Ans.* $48a^5x^2y^4$.

5. $ax - bx$, $ay - by$, and x^2y^2.
Ans. $(a - b)xx.yy = ax^2y^2 - bx^2y^2$.

6. $a + b$, $a^2 - b^2$, and $a^2 + 2ab + b^2$.
Ans. $(a + b)^2 (a - b)$.

7. $3a^3b^2$, $9a^2x^2$, $18a^4y^3$, $3a^2y^2$. *Ans.* $18a^4b^2x^2y^3$.

8. $8a^2(a \quad b)$, $15a^5(a - b)^2$, and $12a^3(a^2 - b^2)$.
Ans. $120a^5(a - b)^2 (a + b)$.

CHAPTER IV.

FRACTIONS.

76. If the unit 1 be divided into any number of equal parts, each part is called a FRACTIONAL UNIT. Thus, $\frac{1}{2}$, $\frac{1}{4}$, $\frac{1}{7}$, $\frac{1}{b}$, are fractional units.

77. A FRACTION is a fractional unit, or a collection of fractional units. Thus, $\frac{1}{2}$, $\frac{3}{4}$, $\frac{5}{7}$, $\frac{a}{b}$, are fractions.

78. Every fraction is composed of two parts, the Denominator and Numerator. The *Denominator* shows into how many equal parts the unit 1 is divided; and the *Numerator* how many of these parts are taken. Thus, in the fraction $\frac{a}{b}$, the denominator b, shows that 1 is divided into b equal parts, and the numerator a, shows that a of these parts are taken. The fractional unit, in all cases, is equal to the reciprocal of the denominator.

---

76. If 1 be divided into any number of equal parts, what is each part called?

77. What is a fraction?

78. Of how many parts is any fraction composed? What are they called? What does the denominator show? What the numerator? What is the fractional unit equal to?

79. An ENTIRE QUANTITY is one which contains no fractional part. Thus, 7, 11, a^3x, $4x^2 - 3y$, are entire quantities.

An entire quantity may be regarded as a fraction whose denominator is 1. Thus, $7 = \frac{7}{1}$, $ab = \frac{ab}{1}$.

80 A MIXED QUANTITY is a quantity containing both entire and fractional parts. Thus, $7\frac{4}{10}$, $8\frac{3}{7}$, $a + \frac{bx}{c}$, are mixed quantities.

81. Let $\frac{a}{b}$ denote any fraction, and q any quantity whatever. From the preceding definitions, $\frac{a}{b}$ denotes that $\frac{1}{b}$ is taken a times; also, $\frac{aq}{b}$ denotes that $\frac{1}{b}$ is taken aq times; that is,

$$\frac{aq}{b} = \frac{a}{b} \times q; \text{ hence,}$$

Multiplying the numerator of a fraction by any quantity, is equivalent to multiplying the fraction by that quantity.

We see, also, that *any quantity may be multiplied by a fraction, by multiplying it by the numerator, and then dividing the result by the denominator.*

82. It is a principle of Division, that the same result will be obtained if we divide the quantity a by the product of two factors, $p \times q$, as would be obtained by dividing it

---

79. What is an entire quantity? When may it be regarded as a fraction?

80. What is a mixed quantity?

81. How may a fraction be multiplied by any quantity?

82 How may a fraction be divided by any quantity?

first by one of the factors, p, and then dividing that result by the other factor, q. That is,

$$\frac{a}{pq} = \left(\frac{a}{p}\right) \div q; \text{ or, } \frac{a}{pq} = \left(\frac{a}{q}\right) \div p; \text{ hence,}$$

Multiplying the denominator of a fraction by any quantity, is equivalent to dividing the fraction by that quantity

83. Since the operations of Multiplication and Division are the converse of each other, it follows, from the preceding principles, that,

Dividing the numerator of a fraction by any quantity, is equivalent to dividing the fraction by that quantity; and,

Dividing the denominator of a fraction by any quantity, is equivalent to multiplying the fraction by that quantity.

84. Since a quantity may be multiplied, and the result divided by the same quantity, without altering the value, it follows that,

Both terms of a fraction may be multiplied by any quantity, or both divided by any quantity, without changing the value of the fraction.

TRANSFORMATION OF FRACTIONS.

85. The *transformation* of a quantity, is the operation of changing its form, without altering its value. The term *reduce* has a technical signification, and means, to *Transform.*

---

83. What follows from the preceding principles?

84. What operations may be performed without altering the value of a fraction?

85. What is the transformation of a quantity?

FIRST TRANSFORMATION.

To reduce an entire quantity to a fractional form having a given denominator.

86. Let a be the quantity, and b the given denominator. We have, evidently, $a = \frac{ab}{b}$; hence, the

RULE.

Multiply the quantity by the given denominator, and write the product over this given denominator.

SECOND TRANSFORMATION.

To reduce a fraction to its lowest terms.

87. A fraction is in its *lowest terms*, when the numerator and denominator contain no common factors.

It has been shown, that both terms of a fraction may be divided by the same quantity, without altering its value. Hence, if they have any common factors, we may strike them out.

RULE.

Resolve each term of the fraction into its prime factors; then strike out all that are common to both.

The same result is attained by dividing both terms of the fraction by any quantity that will divide them, without a remainder; or, by dividing them by their greatest common divisor.

---

86. How do you reduce an entire quantity to a fractional form having a given denominator?

87. How do you reduce a fraction to its lowest terms?

EXAMPLES.

1. Reduce $\frac{15a^2c^2}{25acd}$ to its lowest terms.

Factoring, $$\frac{15a^2c^2}{25acd} = \frac{3.5aacc}{5.5acd};$$

Canceling the common factors, 5, a, and c, we have,

$$\frac{15a^2c^2}{25acd} = \frac{3ac}{5d}. \quad \textit{Ans.}$$

2. Reduce $\frac{85b^7cd^5}{15b^7c^8d^5}$. *Ans.* $\frac{17}{3c^7}$.

3. Reduce $\frac{60c^6d^4f^5}{12c^5d^8f^9}$. *Ans.* $\frac{5c}{d^4f^4}$.

4. Reduce $\frac{ab - ac}{b - c}$. *Ans.* $\frac{a}{1} = a$.

5. Reduce $\frac{n^2 - 2n + 1}{n^2 - 1}$. *Ans.* $\frac{n - 1}{n + 1}$.

6. Reduce $\frac{x^3 - ax^2}{x^2 - 2ax + a^2}$. *Ans.* $\frac{x^2}{x - a}$.

7. Reduce $\frac{96a^3b^2c}{-12a^3b^2c}$. *Ans.* $-\frac{8}{1} = -8$.

8. Reduce $\frac{24b^5 - 36ab^4}{48a^4b^4 - 66a^5b^6}$. *Ans.* $\frac{4b - 6a}{8a^4 - 11a^5b^2}$.

9. Reduce $\frac{a^2 - b^2}{a^2 - 2ab + b^2}$. *Ans.* $\frac{a + b}{a - b}$.

10. Reduce $\frac{5a^3 - 10a^2b + 5ab^2}{8a^3 - 8a^2b}$. *Ans.* $\frac{5(a - b)}{8a}$.

11. Reduce $\frac{3a^2 + 6a^2b^2}{12a^4 + 6a^3c^2}$. *Ans.* $\frac{1 + 2b^2}{4a^2 + 2ac^2}$

12. Reduce $\frac{a^2 + 2ax + x^2}{3(a^2 - x^2)}$. *Ans.* $\frac{a + x}{3(a - x)}$.

THIRD TRANSFORMATION.

To reduce a fraction to a mixed quantity.

88. When any term of the numerator is divisible by any term of the denominator, the transformation can be effected by Division.

RULE.

Perform the indicated division, continuing the operation as far as possible; then write the remainder over the denominator, and annex the result to the quotient found.

EXAMPLES.

1. Reduce $\frac{ax - a^2}{x}$. *Ans.* $a - \frac{a^2}{x}$.

2. Reduce $\frac{ax - x^2}{x}$. *Ans.* $a - x$.

3. Reduce $\frac{ab - 2a^2}{b}$. *Ans.* $a - \frac{2a^2}{b}$.

4. Reduce $\frac{a^2 - x^2}{a - x}$. *Ans.* $a + x$.

5. Reduce $\frac{x^3 - y^3}{x - y}$. *Ans.* $x^2 + xy + y^2$.

6. Reduce $\frac{10x^2 - 5x + 3}{5x}$. *Ans.* $2x - 1 + \frac{3}{5x}$.

7. Reduce $\frac{36x^3 - 72x + 32a^2x^2}{9x}$ .. $4x^2 - 8 + \frac{32a^2x}{9}$.

8. Reduce $\frac{18acf - 6bdcf - 2ad}{3adf}$ .. $\frac{6c}{d} - \frac{2bc}{a} - \frac{2}{3f}$.

9. Reduce $\frac{x^2 + x - 4}{x + 2}$. *Ans.* $x - 1 - \frac{2}{x + 2}$.

---

88. How do you reduce a fraction to a mixed quantity?

10. Reduce $\dfrac{a^2+b^2}{a+b}$. *Ans.* $a-b+\dfrac{2b^2}{a+b}$.

11. Reduce $\dfrac{x^2+3x-25}{x-4}$. *Ans.* $x+7+\dfrac{3}{x-4}$

FOURTH TRANSFORMATION.

To reduce a mixed quantity to a fractional form.

89. This transformation is the converse of the preceding, and may be effected by the following

RULE.

Multiply the entire part by the denominator of the fraction, and add to the product the numerator; write the result over the denominator of the fraction.

EXAMPLES.

1. Reduce $6\frac{1}{7}$ to the form of a fraction.

$$6 \times 7 = 42;\quad 42 + 1 = 43;\ \text{hence,}\ 6\tfrac{1}{7} = \frac{43}{7}.$$

Reduce the following to fractional forms:

2. $x - \dfrac{a^2-x^2}{x} = \dfrac{x^2-(a^2-x^2)}{x}$. *Ans.* $\dfrac{2x^2-a^2}{x}$.

3. $x - \dfrac{ax+x^2}{2a}$. *Ans.* $\dfrac{ax-x^2}{2a}$.

4. $5 + \dfrac{2x-7}{3x}$. *Ans.* $\dfrac{17x-7}{3x}$.

5. $1 - \dfrac{x-a-1}{a}$. *Ans.* $\dfrac{2a-x+1}{a}$

6. $1 + 2x - \dfrac{x-3}{5x}$. *Ans.* $\dfrac{10x^2+4x+3}{5x}$

---

89. How do you reduce a mixed quantity to a fractional form?

7. $2a + b - \frac{3c+4}{8}$. *Ans.* $\frac{16a + 8b - 3c - 4}{8}$.

8. $6ax + b - \frac{6a^2x - ab}{4a}$. *Ans.* $\frac{18a^2x + 5ab}{4a}$.

9. $8 + 3ab - \frac{8 + 6a^2b^2x^4}{12abx^4}$.

Ans. $\frac{96abx^4 + 30a^2b^2x^4 - 8}{12abx^4}$

FIFTH TRANSFORMATION.

To reduce fractions having different denominators, to equivalent fractions having the least common denominator.

90. This transformation is effected by finding the least common multiple of the denominators.

1. Reduce $\frac{1}{3}$, $\frac{3}{4}$, and $\frac{5}{12}$, to their least common denominators.

The least common multiple of the denominators is 12, which is also the least common denominator of the required fractions. If each fraction be multiplied by 12, and the result divided by 12, the values of the fractions will not be changed.

$\frac{1}{3} \times 12 = 4$, 1st new numerator;

$\frac{3}{4} \times 12 = 9$, 2d new numerator;

$\frac{5}{12} \times 12 = 5$, 3rd new numerator; hence,

$\frac{4}{12}$, $\frac{9}{12}$, and $\frac{5}{12}$ are the new equivalent fractions.

---

90. How do you reduce fractions having different denominators, to equivalent fractions having the least common denominator? When the numerators have no common factor, how do you reduce them?

RULE.

I. *Find the least common multiple of the denominators:*

II. *Multiply each fraction by it, and cancel the denominator:*

III. *Write each product over the common multiple, and the results will be the required fractions.*

GENERAL RULE.

Multiply each numerator by all the denominators except its own, for the new numerators, and all the denominators together for a common denominator.

EXAMPLES.

1. Reduce $\frac{a}{a^2 - b^2}$ and $\frac{c}{a + b}$ to their least common denominator.

The least common multiple of the denominators is $(a + b)(a - b)$:

$$\frac{a}{a^2 - b^2} \times (a + b)(a - b) = a$$

$$\frac{c}{a + b} \times (a + b)(a - b) = c(a - b\text{; hence,}$$

$\frac{a}{(a + b)(a - b)}$ and $\frac{c(a - b)}{(a + b)(a - b)}$, are the required fractions.

Reduce the following to their least common denominators:

2. $\frac{3x}{4}$, $\frac{4}{6}$, and $\frac{12x^2}{15}$. *Ans.* $\frac{45x}{60}$, $\frac{40}{60}$, $\frac{48x^2}{60}$

3. a, $\frac{3b^2}{4}$, and $\frac{5c^3}{6}$. *Ans.* $\frac{12a}{12}$, $\frac{9b^2}{12}$, $\frac{10c^3}{12}$

4. $\frac{3x}{2a}$, $\frac{2b}{3c}$, and d. *Ans.* $\frac{9cx}{6ac}$, $\frac{4ab}{6ac}$, $\frac{6acd}{6ac}$

5. $\frac{3}{4}$, $\frac{2x}{3}$, $a + \frac{2x}{a}$. *Ans.* $\frac{9a}{12a}$, $\frac{8ax}{12a}$, $\frac{12a^2 + 24x}{12a}$.

6. $\frac{x}{1 - x}$, $\frac{x^2}{(1 - x)^2}$, and $\frac{x^3}{(1 - x)^3}$.

Ans. $\frac{x(1 - x)^2}{(1 - x)^3}$, $\frac{x^2(1 - x)}{(1 - x)^3}$, and $\frac{x^3}{(1 - x)^3}$

7. $\frac{c}{5a}$, $\frac{c - b}{c}$, and $\frac{c}{a + b}$.

$\frac{ac^2 + bc^2}{5a^2c + 5abc}$, $\frac{5a^2c - 5a^2b + 5abc - 5ab^2}{5a^2c + 5abc}$, $\frac{5ac^2}{5a^2c + 5abc}$.

8. $\frac{cx}{a - x}$, $\frac{dx^2}{a + x}$, and $\frac{x^3}{a + x}$.

Ans. $\frac{cx(a + x)}{a^2 - x^2}$, $\frac{dx^2(a - x)}{a^2 - x^2}$, and $\frac{x^3(a - x)}{a^2 - x^2}$

ADDITION OF FRACTIONS.

91. Fractions can only be added when they have a common unit, that is, when they have a common denominator. In that case, the sum of the numerators will indicate how many times that unit is taken in the entire collection. Hence, the

RULE.

I. *Reduce the fractions to be added, to a common denominator:*

II. *Add the numerators together for a new numerator, and write the sum over the common denominator.*

EXAMPLES.

1. Add $\frac{6}{2}$, $\frac{4}{3}$, and $\frac{2}{5}$, together.

---

91. What is the rule for adding fractions?

By reducing to a common denominator, we have,

$6 \times 3 \times 5 = 90$, 1st numerator.
$4 \times 2 \times 5 = 40$, 2d numerator.
$2 \times 3 \times 2 = 12$, 3d numerator.
$2 \times 3 \times 5 = 30$, the denominator.

Hence, the expression for the sum of the fractions becomes

$$\frac{90}{30} + \frac{40}{30} + \frac{12}{30} = \frac{142}{30};$$

which, being reduced to the simplest form, gives $4\frac{11}{15}$.

2. Find the sum of $\frac{a}{b}$, $\frac{c}{d}$, and $\frac{e}{f}$.

$$\text{Here,}\quad \left.\begin{aligned} a \times d \times f &= adf \\ c \times b \times f &= cbf \\ e \times b \times d &= ebd \end{aligned}\right\}\text{the new numerators.}$$

$$\text{and}\quad b \times d \times f = bdf \quad \text{the common denominator.}$$

Hence, $\frac{adf}{bdf} + \frac{cbf}{bdf} + \frac{ebd}{bdf} = \frac{adf + cbf + ebd}{bdf}$, the sum.

Add the following:

3. $a - \frac{3x^2}{b}$, and $b + \frac{2ax}{c}$. *Ans.* $a + b + \frac{2abx - 3cx^2}{bc}$.

4. $\frac{x}{2}$, $\frac{x}{3}$, and $\frac{x}{4}$. *Ans.* $x + \frac{x}{12}$.

5. $\frac{x - 2}{3}$ and $\frac{4x}{7}$. *Ans.* $\frac{19x - 14}{21}$.

6. $x + \frac{x - 2}{3}$ and $3x + \frac{2x - 3}{4}$. *Ans.* $4x + \frac{10x - 17}{12}$.

7. $4x$, $\frac{5x^2}{2a}$, and $\frac{x + a}{2x}$. *Ans.* $4x + \frac{5x^3 + ax + a^2}{2ax}$.

8. $\frac{2x}{3}$, $\frac{7x}{4}$, and $\frac{2x + 1}{5}$. *Ans.* $2x + \frac{49x + 12}{60}$.

9. $4x$, $\frac{7x}{9}$, and $2 + \frac{x}{5}$. *Ans.* $2 + 4x + \frac{44x}{45}$.

10. $3x + \frac{2x}{5}$ and $x - \frac{8x}{9}$. *Ans.* $3x + \frac{23x}{45}$

11. $ac - \frac{6b}{8a}$ and $1 - \frac{c}{d}$.

Ans. $1 + ac - \frac{6bd + 8ac}{8ad}$.

12. $\frac{3}{(x-1)^3}$, $\frac{3}{(x-1)^2}$, and $\frac{4}{x-1}$.

Ans. $\frac{4x^2 - 5x + 4}{(x-1)^3}$

13. $\frac{1}{4(1+a)}$, $\frac{1}{4(1-a)}$, and $\frac{1}{2(1-a^2)}$. *Ans.* $\frac{1}{1-a^2}$.

SUBTRACTION OF FRACTIONS.

92. Fractions can only be subtracted when they have the same unit; that is, a common denominator. In that case, the numerator of the minuend, *minus* that of the subtrahend, will indicate the number of times that the common unit is to be taken in the difference. Hence, the

RULE.

I. *Reduce the two fractions to a common denominator:*

II. *Then subtract the numerator of the subtrahend from that of the minuend for a new numerator, and write the remainder over the common denominator.*

EXAMPLES.

1. What is the difference between $\frac{3}{7}$ and $\frac{2}{8}$.

$$\frac{3}{7} - \frac{2}{8} = \frac{24}{56} - \frac{14}{56} = \frac{10}{56} = \frac{5}{28}.\ \textit{Ans.}$$

---

92. What is the rule for subtracting fractions

2. Find the difference of the fractions $\frac{x-a}{2b}$ and $\frac{2a-4x}{3c}$

$$\text{Here, } \left\{ \begin{array}{rcl} (x-a) \times 3c & = & 3cx - 3ac \\ (2a-4x) \times 2b & = & 4ab - 8bx \end{array} \right\} \text{the numerators,}$$

and, $2b \times 3c = 6bc$ the common denominator.

$$\text{Hence, } \frac{3cx-3ac}{6bc} - \frac{4ab-8bx}{6bc} = \frac{3cx-3ac-4ab+8bx}{6bc}. \ \textit{Ans.}$$

3. Required the difference of $\frac{12x}{7}$ and $\frac{3x}{5}$. *Ans.* $\frac{39x}{35}$.

4. Required the difference of $5y$ and $\frac{3y}{8}$. *Ans.* $\frac{37y}{8}$.

5. Required the difference of $\frac{3x}{7}$ and $\frac{2x}{9}$. *Ans.* $\frac{13x}{63}$.

6. From $\frac{x+y}{x-y}$ subtract $\frac{x-y}{x+y}$. *Ans.* $\frac{4xy}{x^2-y^2}$.

7. From $\frac{1}{y-z}$ subtract $\frac{1}{y^2-z^2}$. *Ans.* $\frac{y+z-1}{y^2-z^2}$.

Find the differences of the following:

8. $\frac{3x+a}{5b}$ and $\frac{2x+7}{8}$. *Ans.* $\frac{24x+8a-10bx-35b}{40b}$

9. $3x+\frac{x}{b}$ and $x-\frac{x-a}{c}$. *Ans.* $2x+\frac{cx+bx-ab}{bc}$,

10. $a+\frac{a-x}{a(a+x)}$ and $\frac{a+x}{a(a-x)}$. *Ans.* $a-\frac{4x}{a^2-x^2}$.

MULTIPLICATION OF FRACTIONS.

98. Let $\frac{a}{b}$ and $\frac{c}{d}$, represent any two fractions. It has been shown (Art. 81), that any quantity may be multiplied

---

98. What is the rule for the multiplication of fractions?

by a fraction, by first multiplying by the numerator, and then dividing the result by the denominator.

To multiply $\frac{a}{b}$ by $\frac{c}{d}$, we first multiply by c, giving $\frac{ac}{b}$ then, we divide this result by d, which is done by multiplying the denominator by d; this gives for the product, $\frac{ac}{bd}$; that is,

$$\frac{a}{b} \times \frac{c}{d} = \frac{ac}{bd}; \text{ hence,}$$

RULE.

I. *If there are mixed quantities, reduce them to a fractional form; then,*

II. *Multiply the numerators together for a new numerator, and the denominators for a new denominator.*

EXAMPLES.

1. Multiply $a + \frac{bx}{a}$ by $\frac{c}{d}$. First, $a + \frac{bx}{a} = \frac{a^2 + bx}{a}$,

hence, $$\frac{a^2 + bx}{a} \times \frac{c}{d} = \frac{a^2c + bcx}{ad}. \quad \textit{Ans.}$$

Find the products of the following quantities:

2. $\frac{2x}{a}$, $\frac{3ab}{c}$, and $\frac{3ac}{2b}$ *Ans.* $9ax$.

3. $b + \frac{bx}{a}$ and $\frac{a}{x}$. *Ans.* $\frac{ab + bx}{x}$.

4. $\frac{x^2 - b^2}{bc}$ and $\frac{x^2 + b^2}{b + c}$. *Ans.* $\frac{x^4 - b^4}{b^2c + bc^2}$.

5. $x + \frac{x + 1}{a}$, and $\frac{x - 1}{a + b}$. *Ans.* $\frac{ax^2 - ax + x^2 - 1}{a^2 + ab}$.

6. $a + \frac{ax}{a - x}$ and $\frac{a^2 - x^2}{x + x^2}$. *Ans.* $\frac{a^3 + a^2x}{x + x^2}$

7 Multiply $\frac{2a}{a-b}$ by $\frac{a^2-b^2}{3}$.

We have, by the rule,

$$\frac{2a}{a-b} \times \frac{a^2-b^2}{3} = \frac{2a(a^2-b^2)}{3(a-b)} = \frac{2a(a+b)\,(a-b)}{3(a-b)}$$
$$= \frac{2a}{3}(a+b).$$

After indicating the operation, we factored both numerator and denominator, and then canceled the common factors, before performing the multiplication. *This should be done, whenever there are common factors.*

8. $\frac{2}{x-y}$ by $\frac{x^2-y^2}{a}$. *Ans.* $\frac{2(x+y)}{a}$.

9. $\frac{x^2-4}{3}$ by $\frac{4x}{x+2}$. *Ans.* $\frac{4x(x-2)}{3}$.

10. $\frac{(a+b)^2}{2x}$ by $\frac{4x^2}{(a+b)}$. *Ans.* $2x(a+b)$.

11. $\frac{(x-1)^2}{y^3}$ by $\frac{(x+1)y^2}{x-1}$. *Ans.* $\frac{x^2-1}{y}$.

12. $\frac{(a^2-x^2)}{1-x^2}$ by $\frac{1+x}{a+x}$. *Ans.* $\frac{a-x}{1-x}$.

13. $x+\frac{2xy}{x-y}$ by $x-\frac{2xy}{x+y}$ *Ans.* x^2.

14. $\frac{2a-b}{4a}$ by $\frac{6a-2b}{b^2-2ab}$ *Ans.* $\frac{b-3a}{2ab}$.

14. $x-\frac{y^2}{x}$ by $\frac{x}{y}+\frac{y}{x}$. *Ans.* $\frac{x^4-y^4}{x^2y}$

DIVISION OF FRACTIONS.

94 Since $\frac{p}{q} = p \times \frac{1}{q}$, it follows that, dividing by a quantity is equivalent to multiplying by its reciprocal. But the reciprocal of a fraction, $\frac{c}{d}$, is $\frac{d}{c}$ (Art. 28); consequently, to divide any quantity by a fraction, we invert the terms of the divisor, and multiply by the resulting fraction. Hence,

$$\frac{a}{b} \div \frac{c}{d} = \frac{a}{b} \times \frac{d}{c} = \frac{ad}{bc}.$$

Whence, the following rule for dividing one fraction by another:

RULE.

I. *Reduce mixed quantities to fractional forms:*

II. *Invert the terms of the divisor, and multiply the dividend by the resulting fraction.*

NOTE.—The same remarks as were made on *factoring* and *reducing*, under the head of Multiplication, are applicable in Division.

EXAMPLES.

1. Divide $a - \frac{b}{2c}$ by $\frac{f}{g}$.

$$a - \frac{b}{2c} = \frac{2ac - b}{2c}$$

Hence, $a - \frac{b}{2c} \div \frac{f}{g} = \frac{2ac - b}{2c} \times \frac{g}{f} = \frac{2acg - bg}{2cf}$. *Ans*

---

94. What is the rule for the division of fractions?

2. Divide $\frac{2(x+y)}{a}$ by $\frac{x^2-y^2}{a}$.

$$\frac{2(x+y)}{a} \times \frac{a}{x^2-y^2} = \frac{2(x+y)}{a} \times \frac{a}{(x+y)(x-y)}$$
$$= \frac{2}{x-y}. \quad \textit{Ans.}$$

3. Let $\frac{7x}{5}$ be divided by $\frac{12}{13}$. *Ans.* $\frac{91x}{60}$.

4. Let $\frac{4x^2}{7}$ be divided by $5x$. *Ans.* $\frac{4x}{35}$.

5. Let $\frac{x+1}{6}$ be divided by $\frac{2x}{3}$. *Ans.* $\frac{x+1}{4x}$.

6. Let $\frac{x}{x-1}$ be divided by $\frac{x}{2}$. *Ans.* $\frac{2}{x-1}$.

7. Let $\frac{5x}{3}$ be divided by $\frac{2a}{3b}$. *Ans.* $\frac{5bx}{2a}$.

8. Let $\frac{x-b}{8cd}$ be divided by $\frac{3cx}{4d}$. *Ans.* $\frac{x-b}{6c^2x}$.

Divide the following fractions:

9. $\frac{4x^2-8x}{3}$ by $\frac{x^2-4}{3}$. *Ans.* $\frac{4x}{x+2}$.

10. $\frac{x^4-b^4}{x^2-2bx+b^2}$ by $\frac{x^2+bx}{x-b}$. *Ans.* $x+\frac{b^2}{x}$.

11. $2x(a+b)$ by $\frac{4x^2}{a+b}$. *Ans.* $\frac{(a+b)^2}{2x}$.

12. $\frac{x^2-1}{y}$ by $\frac{(x+1)y^2}{x-1}$. *Ans.* $\frac{(x-1)^2}{y^3}$.

13. $\frac{a^2-ax}{bc+bx}$ by $\frac{3(c-x)}{4(a+x)}$. *Ans.* $\frac{4a(a^2-x^2)}{3b(c^2-x^2)}$

14. $\frac{a-x}{1-x}$ by $\frac{1+x}{a+x}$. *Ans.* $\frac{a^2-x^2}{1-x^2}$.

15 x^2 by $x - \frac{2xy}{x+y}$. *Ans.* $\frac{x^2+xy}{x-y}$.

16. $\frac{b-3a}{2ab}$ by $\frac{6a-2b}{b^2-2ab}$. *Ans.* $\frac{2a-b}{4a}$.

17. $\frac{x^4-y^4}{x^2y}$ by $\frac{x}{y}+\frac{y}{x}$. *Ans.* $\frac{x^2-y^2}{x}$.

18. $m^2+1+\frac{1}{m^2}$ by $m+\frac{1}{m}+1$.

Ans. $m+\frac{1}{m}-1$.

19 $\left(x+\frac{y-x}{1+xy}\right)$ by $\left(1-x\frac{y-x}{1+xy}\right)$. *Ans.* y.

20. $\frac{x+2y}{x+y}+\frac{x}{y})$ by $\left(\frac{x+2y}{y}-\frac{x}{x+y}\right)$. *Ans* 1

CHAPTER V.

EQUATIONS OF THE FIRST DEGREE.

95. AN EQUATION is the expression of equality between two quantities. Thus,

$$x = b + c,$$

is an equation, expressing the fact that the quantity x, is equal to the sum of the quantities b and c.

96. Every equation is composed of two parts, connected by the sign of equality. These parts are called *members:* the part on the left of the sign of equality, is called the *first member;* that on the right, the *second member.* Thus, in the equation,

$$x + a = b - c,$$

$x + a$ is the first member, and $b - c$, the second member.

97. An equation of the *first degree* is one which involves only the first power of the unknown quantity; thus,

$$6x + 3x - 5 = 13; \quad (1)$$

and

$$ax + bx + c = d; \quad (2)$$

are equations of the first degree.

---

95. What is an equation?

96. Of how many parts is every equation composed? How are the parts connected? What are the parts called? What is the part on the left called? The part on the right?

97. What is an equation of the first degree

98. A NUMERICAL EQUATION is one in which the coefficients of the unknown quantity are denoted by numbers.

99. A LITERAL EQUATION is one in which the coefficients of the unknown quantity are denoted by letters.

Equation (1) is a numerical equation; Equation (2) is a literal equation.

EQUATIONS OF THE FIRST DEGREE CONTAINING BUT ONE UNKNOWN QUANTITY.

100. The TRANSFORMATION of an equation, is the operation of changing its form without destroying the equality of its members.

101. An AXIOM is a self-evident proposition.

102. The transformation of equations depends upon the following axioms:

1. *If equal quantities be added to both members of an equation, the equality will not be destroyed.*
2. *If equal quantities be subtracted from both members of an equation, the equality will not be destroyed.*
3. *If both members of an equation be multiplied by the same quantity, the equality will not be destroyed.*
4. *If both members of an equation be divided by the same quantity, the equality will not be destroyed.*
5. *Like powers of the two members of an equation are equal.*
6. *Like roots of the two members of an equation are equal.*

---

98. What is a numerical equation?

99 What is a literal equation?

100. What is the transformation of an equation?

101. What is an axiom?

102. Name the axioms on which the transformation of an equation depends.

103. Two principal transformations are employed in the solution of equations of the first degree: *Clearing of fractions*, and *Transposing.*

CLEARING OF FRACTIONS.

1. Take the equation,

$$\frac{2x}{3} - \frac{3x}{4} + \frac{x}{6} = 11.$$

The least common multiple of the denominators is 12. If we multiply both members of the equation by 12, each term will reduce to an entire form, giving,

$$8x - 9x + 2x = 132.$$

Any equation may be reduced to entire terms in the same manner.

104. Hence for clearing of fractions, we have the following

RULE.

I. *Find the least common multiple of the denominators:*

II. *Multiply both members of the equation by it, reducing the fractional to entire terms.*

NOTE.—1. The reduction will be effected, if we divide the least common multiple by each of the denominators, and then multiply the corresponding numerator, dropping the denominator.

2. The transformation may be effected by multiplying each numerator into the product of all the denominators except its own, omitting denominators.

---

103. How many transformations are employed in the solution of equations of the first degree? What are they?

104. Give the rule for clearing an equation of fractions? In what three ways may the reduction be effected?

3. The transformation may also be effected, by *multiplying both members of the equation by any multiple of the denominators.*

EXAMPLES.

Clear the following equations of fractions:

1. $\frac{x}{5} + \frac{x}{7} - 4 = 3.$ *Ans.* $7x + 5x - 140 = 105$

2. $\frac{x}{6} + \frac{x}{9} - \frac{x}{27} = 8.$ *Ans.* $9x + 6x - 2x = 432.$

3. $\frac{x}{2} + \frac{x}{3} - \frac{x}{9} + \frac{x}{12} = 20.$
Ans. $18x + 12x - 4x + 3x = 720.$

4. $\frac{x}{5} + \frac{x}{7} - \frac{x}{2} = 4.$ *Ans.* $14x + 10x - 35x = 280.$

5. $\frac{x}{4} - \frac{x}{5} + \frac{x}{6} = 15.$ *Ans.* $15x - 12x + 10x = 900$

6. $-\frac{x-4}{3} - \frac{x-2}{6} = \frac{5}{3}.$
Ans. $-2x + 8 - x + 2 = 10.$

7. $\frac{x}{3-x} + 4 = \frac{3}{5}.$ *Ans.* $5x + 60 - 20x = 9 - 3x.$

8. $\frac{x}{4} - \frac{x}{6} + \frac{x}{8} + \frac{x}{9} = 12.$
Ans. $18x - 12x + 9x + 8x = 864.$

9. $\frac{a}{b} - \frac{c}{d} + f = g.$ *Ans.* $ad - bc + bdf = bdg$

10. $\frac{ax}{b} - \frac{2c^2x}{ab} + 4a = \frac{4bc^2x}{a^3} - \frac{5a^3}{b^2} + \frac{2c^2}{a} - 3b.$

The least common multiple of the denominators is a^3b^2

$$a^4bx - 2a^2bc^2x + 4a^4b^2 = 4b^3c^2x - 5a^6 + 2a^2b^2c^2 - 3a^3b^3.$$

TRANSPOSING.

105. **Transposition** is the operation of changing a term from one member to the other, without destroying the equality of the members.

1. Take, for example, the equation,

$$5x - 6 = 8 + 2x.$$

If, in the first place, we subtract $2x$ from both members the equality will not be destroyed, and we have,

$$5x - 6 - 2x = 8.$$

Whence we see, that the term $2x$, which was additive in the second member, becomes subtractive by passing into the first.

In the second place, if we add 6 to both members of the last equation, the equality will still exist, and we have,

$$5x - 6 - 2x + 6 = 8 + 6,$$

or, since -6 and $+6$ cancel each other, we have,

$$5x - 2x = 8 + 6.$$

Hence, the term which was subtractive in the first member, passes into the second member with the sign of addition.

106. Therefore, for the transposition of the terms, we have the following

RULE.

Any term may be transposed from one member of an equation to the other, if the sign be changed.

---

105. What is transposition?

106. What is the rule for the transposition of the terms of an equation?

EXAMPLES.

Transpose the unknown terms to the first member, and the known terms to the second, in the following:

1. $3x + 6 - 5 = 2x - 7.$ *Ans.* $3x - 2x = -7 - 6 + 5.$
2. $ax + b = d - cx.$ *Ans.* $ax + cx = d - b.$
3. $4x - 3 = 2x + 5.$ *Ans.* $4x - 2x = 5 + 3.$
4. $9x + c = cx - d.$ *Ans.* $9x - cx = -d - c.$
5. $ax + f = dx + b.$ *Ans.* $ax - dx = b - f.$
6. $6x - c = -ax + b.$ *Ans.* $6x + ax = b + c.$

SOLUTION OF EQUATIONS.

107. The SOLUTION of an equation is the operation of finding such a value for the unknown quantity, as will *satisfy* the equation; that is, such a value as, being substituted for the unknown quantity, will render the two members equal. This is called a ROOT of the equation.

A *Root* of an equation is said to be *verified*, when being substituted for the unknown quantity in the given equation, the two members are found equal to each other.

1. Take the equation,

$$\frac{3x}{2} - 4 = \frac{4(x-2)}{8} + 3.$$

Clearing of fractions (Art. 104), and performing the operations indicated, we have,

$$12x - 32 = 4x - 8 + 24.$$

---

107. When is the solution of an equation? What is the found value of the unknown quantity called? When is a root of an equation said to be verified.

Transposing all the unknown terms to the first member, and the known terms to the second (Art. 106), we have,

$$12x - 4x = -8 + 24 + 32.$$

Reducing the terms in the two members,

$$8x = 48.$$

Dividing both members by the coefficient of x,

$$x = \frac{48}{8} = 6.$$

VERIFICATION.

$$\frac{3 \times 6}{2} - 4 = \frac{4(6 - 2)}{8} + 3; \text{ or,}$$

$$+9 - 4 = 2 + 3 = 5.$$

Hence, 6 satisfies the equation, and therefore, is a root.

108. By processes similar to the above, all equations of the first degree, containing but one unknown quantity, may be solved.

RULE.

I. *Clear the equation of fractions, and perform all the indicated operations:*

II. *Transpose all the unknown terms to the first member, and all the known terms to the second member:*

III. *Reduce all the terms in the first member to a single term, one factor of which will be the unknown quantity, and the other factor will be the algebraic sum of its coefficients:*

IV. *Divide both members by the coefficient of the unknown quantity: the second member will then be the value of the unknown quantity.*

---

108. Give the rule for solving equations of the first degree with one unknown quantity.

EXAMPLES.

1. Solve the equation,

$$\frac{5x}{12} - \frac{4x}{3} - 13 = \frac{7}{8} - \frac{13x}{6}.$$

Clearing of fractions,

$$10x - 32x - 312 = 21 - 52x.$$

By transposing,

$$10x - 32x + 52x = 21 + 312.$$

By reducing, $30x = 333;$

hence, $x = \frac{333}{30} = \frac{111}{10} = 11.1;$

a result which may be verified by substituting it for x in the given equation.

2. Solve the equation,

$$(3a - x)(a - b) + 2ax = 4b(x + a).$$

Performing the indicated operations, we have,

$$3a^2 - ax - 3ab + bx + 2ax = 4bx + 4ab.$$

By transposing,

$$- ax + bx + 2ax - 4bx = 4ab + 3ab - 3a^2.$$

By reducing, $ax - 3bx = 7ab - 3a^2;$

Factoring, $(a - 3b)x = 7ab - 3a^2.$

Dividing both members by the coefficient of x,

$$x = \frac{7ab - 3a^2}{a - 3b}.$$

3. Given $3x - 2 + 24 = 31$ to find x. *Ans.* $x = 3$.

4. Given $x + 18 = 3x - 5$ to find x *Ans.* $x = 11\frac{1}{2}$

5. Given $6 - 2x + 10 = 20 - 3x - 2$, to find x.
Ans. $x = 2$.

6. Given $x + \frac{1}{2}x + \frac{1}{3}x = 11$, to find x. *Ans.* $x = 6$.

7. Given $2x - \frac{1}{2}x + 1 = 5x - 2$, to find x.
Ans. $x = \frac{6}{7}$.

Solve the following equations:

8. $3ax + \frac{a}{2} - 3 = bx - a$. *Ans.* $x = \frac{6 - 3a}{6a - 2b}$.

9. $\frac{x - 3}{2} + \frac{x}{3} = 20 - \frac{x - 19}{2}$. *Ans.* $x = 23\frac{1}{4}$.

10. $\frac{x + 3}{2} + \frac{x}{3} = 4 - \frac{x - 5}{4}$. *Ans.* $x = 3\frac{6}{13}$.

11. $\frac{x}{4} - \frac{3x}{2} + x = \frac{4x}{8} - 3$. *Ans.* $x = 4$.

12. $\frac{3ax}{c} - \frac{2bx}{d} - 4 = f$. *Ans.* $x = \frac{cdf + 4cd}{3ad - 2bc}$

13. $\frac{x - a}{3} - \frac{2x - 3b}{5} - \frac{a - x}{2} = 10a + 11b$.
Ans. $x = 25a + 24b$.

14. $\frac{x}{12} - \frac{8 - x}{8} - \frac{5 + x}{4} + \frac{11}{4} = 0$. *Ans.* $x = 12$.

15. $\frac{a + c}{a + x} + \frac{a - c}{a - x} = \frac{2b^2}{a^2 - x^2}$ *Ans.* $x = \frac{a^2 - b^2}{c}$

16. $\frac{8ax - b}{7} - \frac{3b - c}{2} = 4 - b$.
Ans $x = \frac{56 + 9b - 7c}{16a}$

17 $\frac{x}{5} - \frac{x - 2}{3} + \frac{x}{2} = \frac{13}{3}$. *Ans.* $x = 10$.

18. $\frac{x}{a} - \frac{x}{b} + \frac{x}{c} - \frac{x}{d} = f.$

Ans. $x = \frac{abcdf}{bcd - acd + abd - abc}.$

Note.—What is the numerical value of x, when $a = 1$, $b = 2$, $c = 3$, $d = 4$, and $f = 6$?

19. $\frac{x}{7} - \frac{8x}{9} - \frac{x-3}{5} = -12\frac{29}{45}.$ *Ans.* $x = 14.$

20. $x - \frac{3x-5}{13} + \frac{4x-2}{11} = x + 1.$ *Ans.* $x = 6.$

21. $x + \frac{x}{4} + \frac{x}{5} - \frac{x}{6} = 2x - 43.$ *Ans.* $x = 60.$

22. $2x - \frac{4x-2}{5} = \frac{3x-1}{2}.$ *Ans.* $x = 3.$

23. $3x + \frac{bx-d}{3} = x + a.$ *Ans.* $x = \frac{3a+d}{6+b}.$

24. $\frac{ax-b}{4} + \frac{a}{3} = \frac{bx}{2} - \frac{bx-a}{3}.$

Ans. $x = \frac{3b}{3a-2b}.$

25. $\frac{4x}{5-x} - \frac{20-4x}{x} = \frac{15}{x}.$ *Ans.* $x = 3\frac{2}{11}.$

26. $\frac{2x+1}{29} - \frac{402-3x}{12} = 9 - \frac{471-6x}{2}.$

Ans. $x = 72.$

27. $\frac{(a+b)(x-b)}{a-b} - 3a = \frac{4ab-b^2}{a+b} - 2x + \frac{a^2-bx}{b}.$

Ans. $x = \frac{a^4 + 3a^3b + 4a^2b^2 - 6ab^3 + 2b^4}{2b(2a^2 + ab - b^2)}.$

PROBLEMS.

109. A PROBLEM is a question proposed, requiring a solution.

The SOLUTION of a problem is the operation of finding a quantity, or quantities, that will satisfy the given conditions.

The solution of a problem consists of two parts:

I. The STATEMENT, *which consists in expressing, algebraically, the relation between the known and the required quantities.*

II. The SOLUTION, *which consists in finding the values of the unknown quantites, in terms of those which are known.*

The statement is made by representing the unknown quantities of the problem by some of the final letters of the alphabet, and then operating upon these so as to comply with the conditions of the problem. The method of stating problems is best learned by practical examples.

1. What number is that to which if 5 be added, the sum will be equal to 9?

Denote the number by x. Then, by the conditions,

$$x + 5 = 9.$$

This is the *statement* of the problem.

To find the value of x, transpose 5 to the second member; then,

$$x = 9 - 5 = 4.$$

This is the *solution* of the equation.

VERIFICATION.

$$x + 5 = 9.$$

---

109. What is a problem? What is the solution of a problem? Of how many parts does it consist? What are they? What is the statement? What is the solution?

2. Find a number such that the sum of one-half, one-third, and one-fourth of it, augmented by 45, shall be equal to 448

Let the required number be denoted by x.

Then, one-half of it will be denoted by $\frac{x}{2}$,

one-third " " by $\frac{x}{3}$,

one-fourth " " by $\frac{x}{4}$;

and, by the conditions,

$$\frac{x}{2}+\frac{x}{3}+\frac{x}{4}+45 = 448.$$

This is the statement of the problem.

Clearing of fractions,

$$6x + 4x + 3x + 540 = 5376,$$

Transposing and collecting the unknown terms,

$$13x = 4836;$$

hence,
$$x = \frac{4836}{13} = 372.$$

VERIFICATION.

$$\frac{372}{2}+\frac{372}{3}+\frac{372}{4}+45 = 186 + 124 + 93 + 45 = 448.$$

3. What number is that whose third part exceeds its fourth by 16?

Let the required number be denoted by x. Then,

$$\frac{1}{3}x = \text{the third part,}$$

$$\frac{1}{4}x = \text{the fourth part}$$

and, by the conditions of the problem,

$$\frac{1}{3}x - \frac{1}{4}x = 16.$$

This is the statement. Clearing of fractions,

$$4x - 3x = 192,$$

and hence, $x = 192.$

VERIFICATION.

$$\frac{192}{3} - \frac{192}{4} = 64 - 48 = 16.$$

4. Divide \$1000 between *A*, *B*, and *C*, so that *A* shall have \$72 more than *B*, and *C* \$100 more than *A*.

Let x denote the number of dollars which *B* received.

Then, $x = $ *B's* number,

$x + 72 = $ *A's* number,

and, $x + 172 = $ *C's* number;

and their sum, $3x + 244 = 1000$, the number of dollars.

This is the statement. By transposing,

$$3x = 1000 - 244 = 756;$$

and, $x = \frac{756}{3} = 252 = $ *B's* share.

Hence, $x + 72 = 252 + 72 = 324 = $ *A's* share,

and, $x + 172 = 252 + 172 = 424 = $ *C's* share.

VERIFICATION.

$$252 + 324 + 424 = 1000.$$

5. Out of a cask of wine which had leaked away a third part, 21 gallons were afterwards drawn, and the cask being then guaged, appeared to be half full: how much did it hold?

Let x denote the number of gallons.

Then, $\frac{x}{3}$ = the number that had leaked away.

and, $\frac{x}{3} + 21$ = what had leaked and been drawn.

Hence, by the conditions, $\frac{x}{3} + 21 = \frac{x}{2}$.

This is the statement. Clearing of fractions,

$$2x + 126 = 3x,$$

and,

$$-x = -126;$$

and by changing the signs of both members, which does not destroy their equality (since it is equivalent to multiplying both members by -1), we have,

$$x = 126.$$

VERIFICATION.

$$\frac{126}{3} + 21 = 42 + 21 = 63 = \frac{126}{2}.$$

6. A fish was caught whose tail weighed 9 lbs., his head weighed as much as his tail and half his body, and his body weighed as much as his head and tail together: what was the weight of the fish?

Let $2x$ = the weight of the body, in pounds.

Then, $9 + x$ = weight of the head;

and since the body weighed as much as both head and tail,

$$2x = 9 + 9 + x,$$

which is the statement. Then,

$$2x - x = 18, \text{ and } x = 18.$$

Hence, we have,

$$\begin{aligned} 2x &= 36lb. = \text{weight of the body,} \\ 9 + x &= 27lb. = \text{weight of the head,} \\ & \quad \underline{9lb.} = \text{weight of the tail;} \end{aligned}$$

hence, $72lb. = $ weight of the fish.

7. The sum of two numbers is 67, and their difference 19. what are the two numbers?

Let x denote the less number.
Then, $x + 19 =$ the greater; and, by the conditions,

$$2x + 19 = 67.$$

This is the statement. Transposing,

$$2x = 67 - 19 = 48;$$

hence, $x = \frac{48}{2} = 24$, and $x + 19 = 43$.

VERIFICATION.

$$43 + 24 = 67, \text{ and } 43 - 24 = 19.$$

ANOTHER SOLUTION.

Let x denote the greater number.
Then, $x - 19$ will represent the less,
and, $2x - 19 = 67$; whence $2x = 67 + 19$.

Therefore, $x = \frac{86}{2} = 43;$

and, consequently, $x - 19 = 43 - 19 = 24$.

GENERAL SOLUTION OF THIS PROBLEM.

The sum of two numbers is s, their difference is d: what are the two numbers?

Let x denote the *less* number.
Then, $x + d$ will denote the greater,
and $2x + d = s$, their sum. Whence,

$$x = \frac{s - d}{2} = \frac{s}{2} - \frac{d}{2};$$

and, consequently,

$$x + d = \frac{s}{2} - \frac{d}{2} + d = \frac{s}{2} + \frac{d}{2}.$$

As these two results are not dependent on particular values attributed to s or d, it follows that:

1. *The greater of two numbers is equal to half their sum plus half their difference:*

2. *The less is equal to half their sum, minus half their difference.*

Thus, if the sum of two numbers is 32, and their difference 16,

the greater is, $\frac{32}{2} + \frac{16}{2} = 16 + 8 = 24$; and

the less, $\frac{32}{2} - \frac{16}{2} = 16 - 8 = 8$.

VERIFICATION.

$$24 + 8 = 32; \text{ and } 24 - 8 = 16.$$

8. A person engaged a workman for 48 days. For each day that he labored he received 24 cents, and for each day that he was idle, he paid 12 cents for his board. At the end of the 48 days, the account was settled, when the laborer received 504 cents. *Required, the number of working days, and the number of days he was idle.*

If the number of working days, and the number of idle days, were known, and the first multiplied by 24, and the second by 12, the difference of these products would be 504. Let us indicate these operations by means of algebraic signs.

Let x denote the number of working days.

Then, $48 - x =$ the number of idle days,

$24 \times x =$ the amount earned,

and, $12(48 - x) =$ the amount paid for board.

Then,
$$24x - 12(48 - x) = 504,$$
what was received, which is the statement.

Then, performing the operations indicated,

$$24x - 576 + 12x = 504,$$

or,
$$36x = 504 + 576 = 1080,$$

and,
$$x = \frac{1080}{36} = 30, \text{ the number of working days;}$$

whence,
$$48 - 30 = 18, \text{ the number of idle days.}$$

VERIFICATION.

Thirty days' labor, at 24 cents a day, amounts to . . . . . $30 \times 24 = 720$ cents.

And 18 days' board, at 12 cents a day, amounts to . . . . . $18 \times 12 = \underline{216}$ cents.

The difference is the amount received . . . . 504 cents.

GENERAL SOLUTION.

This problem may be made general, by denoting the whole number of working and idle days, by n;

The amount received for each day's work, by a;

The amount paid for board, for each idle day, by b;

And what was due the laborer, or the balance of the account, by c.

As before, let the number of working days be denoted by x.

The number of idle days will then be denoted by $n - x$.

Hence, what is earned will be expressed by ax, and the sum to be deducted, on account of board, by $b(n - x)$.

The statement of the problem, therefore, is,

$$ax - b(n - x) = c.$$

Performing indicated operations,

$$ax - bn + bx = c, \quad \text{or,} \quad (a + b)x = c + bn;$$

whence, $x = \dfrac{c + bn}{a + b}$ = number of working days;

and, $n - x = n - \dfrac{c + bn}{a + b} = \dfrac{an + bn - c - bn}{a + b}$,

or, $n - x = \dfrac{an - c}{a + b}$ = number of idle days.

Let us suppose $n = 48$, $a = 24$, $b = 12$, and $c = 504$; these numbers will give for x the same value as before found.

9. A person dying leaves half of his property to his wife, one-sixth to each of two daughters, one-twelfth to a servant, and the remaining \$600 to the poor; what was the amount of the property?

Let x denote the amount, in dollars,

Then, $\dfrac{x}{2}$ = what he left to his wife,

$\dfrac{x}{6}$ = what he left to one daughter,

and, $\dfrac{2x}{6} = \dfrac{x}{3}$ what he left to both daughters,

also, $\dfrac{x}{12}$ = what he left to his servant,

and, \$600 = what he left to the poor.

Then, by the conditions,

$$\frac{x}{2} + \frac{x}{3} + \frac{x}{12} + 600 = x, \text{ the amount of the property,}$$

which gives, $x = \$7200.$

10. A and B play together at cards. A sits down with \$84, and B with \$48. Each loses and wins in turn, when it appears that A has five times as much as B. How much did A win?

Let x denote the number of dollars A won.

Then, A rose with $84 + x$ dollars,
and B rose with $48 - x$ dollars.

But, by the conditions, we have,

$$84 + x = 5(48 - x),$$

hence, $84 + x = 240 - 5x;$

and, $6x = 156,$

consequently, $x = 26$; or A won \$26.

VERIFICATION.

$$84 + 26 = 110; \quad 48 - 26 = 22;$$
$$110 = 5(22) = 110.$$

11. A can do a piece of work alone in 10 days, B in 13 days; in what time can they do it if they work together?

Denote the time by x, and the work to be done, by 1. Then, in

1 day, A can do $\frac{1}{10}$ of the work, and

B can do $\frac{1}{13}$ of the work; and in

x days, A can do $\frac{x}{10}$ of the work, and

B can do $\frac{x}{13}$ of the work.

Hence, by the conditions,

$$\frac{x}{10} + \frac{x}{13} = 1 \quad \text{which gives,} \quad 13x + 10x = 130;$$

hence, $23x = 130, \quad x = \frac{130}{23} = 5\frac{15}{23}$ days.

12 A fox, pursued by a hound, has a start of 60 of his own leaps. Three leaps of the hound are equivalent to 7 of the fox; but while the hound makes 6 leaps, the fox makes 9: how many leaps must the hound make to overtake the fox?

There is some difficulty in this problem, arising from the different units which enter into it.

Since 3 leaps of the hound are equal to 7 leaps of the fox, 1 leap of the hound is equal to $\frac{7}{3}$ fox leaps.

Since, while the hound makes 6 leaps, the fox makes 9, while the hound makes 1 leap, the fox will make $\frac{9}{6}$, or $\frac{3}{2}$ leaps.

Let x denote the *number* of leaps which the hound makes before he overtakes the fox; and let 1 fox leap denote the *unit of distance.*

Since 1 leap of the hound is equal to $\frac{7}{3}$ of a fox leap, x leaps will be equal to $\frac{7}{3}x$ fox leaps; and this will denote the distance passed over by the hound, in fox leaps.

Since, while the hound makes 1 leap, the fox makes $\frac{3}{2}$ leaps, while the hound makes x leaps, the fox makes $\frac{3}{2}x$ leaps; and this added to 60, his distance ahead, will give $\frac{3}{2}x + 60$, for the whole distance passed over by the fox.

Hence, from the conditions,

$$\frac{7}{3}x = \frac{3}{2}x + 60; \text{ wnence,}$$
$$14x = 9x + 360;$$
$$x = 72.$$

The hound, therefore, makes 72 leaps before overtaking the fox; in the same time, the fox makes $72 \times \frac{3}{2} = 108$ leaps.

VERIFICATION.

$$108 + 60 = 168, \text{ whole number of fox leaps,}$$
$$72 \times \frac{7}{3} = 168.$$

13. A father leaves his property, amounting to $2520, to four sons, *A*, *B*, *C*, and *D*. *C* is to have $360, *B* as much as *C* and *D* together, and *A* twice as much as *B*, less $1000: how much do *A*, *B*, and *D* receive?

Ans. *A*, $760; *B*, $880; *D*, $520.

14. An estate of $7500 is to be divided among a widow, two sons, and three daughters, so that each son shall receive twice as much as each daughter, and the widow herself $500 more than all the children: what was her share, and what the share of each child?

Ans.		
	Widow's share,	$4000.
	Each son's,	1000.
	Each daughter's,	500.

15. A company of 180 persons consists of men, women, and children. The men are 8 more in number than the women, and the children 20 more than the men and women together: how many of each sort in the company?

Ans. 44 men, 36 women, 100 children.

16. A father divides $2000 among five sons, so that each elder should receive $40 more than his next younger brother: what is the share of the youngest? *Ans.* $320.

17. A purse of $2850 is to be divided among three persons, *A*, *B*, and *C*. *A's* share is to be to *B's* as 6 to 11, and *C* is to have $300 more than *A* and *B* together: what is each one's share? *A's*, $450; *B's*, $825; *C's*, $1575.

18. Two pedestrians start from the same point and travel in the same direction; the first steps twice as far as the second, but the second makes 5 steps while the first makes but one. At the end of a certain time they are 300 feet apart. Now, allowing each of the longer paces to be 3 feet, how far will each have traveled?

Ans. 1st, 200 feet; 2d, 500.

19. Two carpenters, 24 journeymen, and 8 apprentices received at the end of a certain time $144. The carpenters received $1 per day, each journeyman, half a dollar, and each apprentice, 25 cents: how many days were they employed? *Ans.* 9 days.

20. A capitalist receives a yearly income of $2940; four-fifths of his money bears an interest of 4 per cent., and the remainder of 5 per cent.: how much has he at interest?

Ans. $70000.

21. A cistern containing 60 gallons of water has three unequal cocks for discharging it; the largest will empty it in one hour, the second in two hours, and the third, in three: in what time will the cistern be emptied if they all run together? *Ans.* $32\frac{8}{11}$ min.

22. In a certain orchard, one-half are apple trees, one-fourth peach trees, one-sixth plum trees; there are also, 120 cherry trees, and 80 pear trees: how many trees in the orchard? *Ans.* 2400.

23. A farmer being asked how many sheep he had,

answered, that he had them in five fields; in the 1st he had $\frac{1}{4}$, in the 2d, $\frac{1}{6}$, in the 3d, $\frac{1}{8}$, and in the 4th, $\frac{1}{12}$, and in the 5th, 450: how many had he? *Ans.* 1200.

24. My horse and saddle together are worth $132, and the horse is worth ten times as much as the saddle: what is the value of the horse? *Ans.* $120.

25. The rent of an estate is this year 8 per cent. greater than it was last. This year it is $1890: what was it last year? *Ans.* $1750.

26. What number is that, from which if 5 be subtracted, $\frac{2}{3}$ of the remainder will be 40? *Ans.* 65.

27. A post is $\frac{1}{4}$ in the mud, $\frac{1}{3}$ in the water, and 10 feet above the water: what is the whole length of the post? *Ans.* 24 feet.

28. After paying $\frac{1}{4}$ and $\frac{1}{5}$ of my money, I had 66 guineas left in my purse: how many guineas were in it at first? *Ans.* 120.

29. A person was desirous of giving 3 pence apiece to some beggars, but found he had not money enough in his pocket by 8 pence; he therefore gave them each 2 pence and had 3 pence remaining: required the number of beggars. *Ans.* 11.

30. A person, in play, lost $\frac{1}{4}$ of his money, and then won 3 shillings; after which he lost $\frac{1}{3}$ of what he then had; and this done, found that he had but 12 shillings remaining: what had he at first? *Ans.* 20*s.*

31. Two persons, *A* and *B*, lay out equal sums of money in trade; *A* gains $126, and *B* loses $87, and *A's* money is then double of *B's*: what did each lay out? *Ans.* $300.

32. A person goes to a tavern with a certain sum of money in his pocket, where he spends 2 shillings: he then borrows as much money as he had left, and going to another tavern, he there spends 2 shillings also; then borrowing

again as much money as was left, he went to a third tavern, where likewise he spent 2 shillings, and borrowed as much as he had left: and again spending 2 shillings at a fourth tavern, he then had nothing remaining. What had he at first? *Ans.* 3*s.* 9*d.*

33. A tailor cut 19 yards from each of three equal pieces of cloth, and 17 yards from another of the same length, and found that the four remnants were together equal to 142 yards. How many yards in each piece? *Ans.* 54.

34. A fortress is garrisoned by 2600 men, consisting of infantry, artillery, and cavalry. Now, there are nine times as many infantry, and three times as many artillery soldiers as there are cavalry. How many are there of each corps?
Ans. 200 cavalry; 600 artillery; 1800 infantry.

35. All the journeyings of an individual amounted to 2970 miles. Of these he traveled $3\frac{1}{2}$ times as many by water as on horseback, and $2\frac{1}{4}$ times as many on foot as by water. How many miles did he travel in each way?
Ans. 240 miles; 840 m.; 1890 m.

36. A sum of money was divided between two persons, *A* and *B*. *A's* share was to *B's* in the proportion of 5 to 3, and exceeded five-ninths of the entire sum by 50. What was the share of each? *Ans.* *A's* share, 450; *B's*, 270.

37. Divide a number a into three such parts that the second shall be n times the first, and the third m times as great as the first.

$$1\text{st}, \frac{a}{1+m+n}; \quad 2\text{d}, \frac{na}{1+m+n}; \quad 3\text{d}, \frac{ma}{1+m+n}.$$

38. A father directs that \$1170 shall be divided among his three sons, in proportion to their ages. The oldest is twice as old as the youngest, and the second is one-third older than the youngest. How much was each to receive?
Ans. \$270, youngest; \$360, second; \$540, oldest.

39. Three regiments are to furnish 594 men, and each to furnish in proportion to its strength. Now, the strength of the first is to the second as 3 to 5; and that of the second to the third as 8 to 7. How many must each furnish?

Ans. 1st, 144 men; 2d, 240; 3d, 210

40. Five heirs, *A*, *B*, *C*, *D*, and *E*, are to divide an inheritance of $5600. *B* is to receive twice as much as *A*, and $200 more; *C* three times as much as *A*, less $400; *D* the half of what *B* and *C* receive together, and 150 more; and *E* the fourth part of what the four others get, plus $475. How much did each receive?

A's, $500; *B's*, 1200; *C's*, 1100; *D's*, 1300; *E's*, 1500.

41. A person has four casks, the second of which being filled from the first, leaves the first four-sevenths full. The third being filled from the second, leaves it one-fourth full, and when the third is emptied into the fourth, it is found to fill only nine-sixteenths of it. But the first will fill the third and fourth, and leave 15 quarts remaining. How many gallons does each hold?

Ans. 1st, 35 gal.; 2d, 15 gal.; 3d, 11¼ gal.; 4th, 20 gal.

42. A courier having started from a place, is pursued by a second after the lapse of 10 days. The first travels 4 miles a day, the other 9. How many days before the second will overtake the first? *Ans.* 8.

43. A courier goes 31½ miles every five hours, and is followed by another after he had been gone eight hours. The second travels 22½ miles every three hours. How many hours before he will overtake the first? *Ans.* 42.

44. Two places are eighty miles apart, and a person leaves one of them and travels towards the other at the rate of 3½ miles per hour. Eight hours after, a person departs from

the second place, and travels at the rate of $5\frac{1}{4}$ miles per hour How long before they will be together?

Ans. 6 hours.

EQUATIONS CONTAINING TWO UNKNOWN QUANTITIES.

110. If we have a single equation, as,

$$2x + 3y = 21,$$

containing two unknown quantities, x and y, we may find the value of one of them in terms of the other, as,

$$x = \frac{21 - 3y}{2} \quad . \; . \; . \; . \; . \; . \quad (1.)$$

Now, if the value of y is unknown, that of x will also be unknown. Hence, from a *single* equation, containing two unknown quantities, the value of x cannot be determined.

If we have a second equation, as,

$$5x + 4y = 35,$$

we may, as before, find the value of x in terms of y, giving,

$$x = \frac{35 - 4y}{5} \quad . \; . \; . \; . \; . \; . \quad (2.)$$

Now, if the values of x and y are the same in Equations (1) and (2), the second members may be placed equal to each other, giving,

$$\frac{21 - 3y}{2} = \frac{35 - 4y}{5}, \quad \text{or} \quad 105 - 15y = 70 - 8y;$$

from which we find, $y = 5.$

---

110. In one equation containing two unknown quantities, can you find the value of either? If you have a second equation involving the same two unknown quantities, can you find their values? What are such equations called?

Subtituting this value for y in Equations (1) or (2), we find $x = 3$. Such equations are called *Simultaneous equations.* Hence,

111. SIMULTANEOUS EQUATIONS are those in which the values of the unknown quantity are the same in both.

ELIMINATION.

112. ELIMINATION is the operation of combining two equations, containing two unknown quantities, and deducing therefrom a single equation, containing but one.

There are three principal methods of elimination:

1st. By addition or subtraction.
2d. By substitution.
3d. By comparison.

We shall consider these methods separately.

Elimination by Addition or Subtraction.

1. Take the two equations,

$$\begin{aligned} 3x - 2y &= 7, \\ 8x + 2y &= 48. \end{aligned}$$

If we add these two equations, member to member, we obtain,

$$11x = 55;$$

which gives, by dividing by 11,

$$x = 5;$$

and substituting this value in either of the given equations, we find,

$$y = 4.$$

---

111. What are simultaneous equations?

112. What is elimination? How many methods of elimination are there? What are they?

2. Again, take the equations,

$$8x + 2y = 48,$$
$$3x + 2y = 23.$$

If we subtract the 2d equation from the 1st, we obtain,

$$5x = 25;$$

which gives, by dividing by 5,

$$x = 5;$$

and by substituting this value, we find,

$$y = 4.$$

3. Given the sum of two numbers equal to s, and their difference equal to d, to find the numbers.

Let $x =$ the greater, and y the less number.

Then, by the conditions, ... $x + y = s$.
and, ... $x - y = d$.
By adding (Art. 102, Ax. 1), ... $2x = s + d$.
By subtracting (Art. 102, Ax. 2), . . . $2y = s - d$.

Each of these equations contains but one unknown quantity.

From the first, we obtain, ... $x = \frac{s + d}{2}$,

and from the second, ... $y = \frac{s - d}{2}$.

These are the same values as were found in Prob. 7, page 120.

4. A person engaged a workman for 48 days. For each day that he labored he was to receive 24 cents, and for each day that he was idle he was to pay 12 cents for his board. At the end of the 48 days the account was settled, when the laborer received 504 cents. Required the number of working days, and the number of days he was idle.

Let $x =$ the number of working days,

$y =$ the number of idle days.

Then, $24x =$ what he earned,

and, $12y =$ what he paid for his board.

Then, by the conditions of the question, we have,

$$x + y = 48,$$

and, $$24x - 12y = 504.$$

This is the statement of the problem.

It has already been shown (Art. 102, Ax. 3), that the two members of an equation may be multiplied by the same number, without destroying the equality. Let, then, the first equation be multiplied by 24, the coefficient of x in the second; we shall then have,

$$\begin{array}{r} 24x + 24y = 1152 \\ \underline{24x - 12y = 504} \end{array}$$

and by subtracting, $$36y = 648$$

$$\therefore \quad y = \frac{648}{36} = 18.$$

Substituting this value of y in the equation,

$24x - 12y = 504$, we have, $24x - 216 = 504$;

which gives,

$$24x = 504 + 216 = 720, \text{ and } x = \frac{720}{24} = 30.$$

VERIFICATION.

$x + y = 48$ gives $30 + 18 = 48.$

$24x - 12y = 504$ gives $24 \times 30 - 12 \times 18 = 504.$

113. In a similar manner, either unknown quantity may be eliminated from either equation; hence, the following

RULE.

I. *Prepare the equations so that the coefficients of the quantity to be eliminated shall be numerically equal:*

II. *If the signs are unlike, add the equations, member to member; if alike, subtract them, member from member.*

EXAMPLES.

Find the values of x and y, by addition or subtraction, in the following simultaneous equations:

5. $\left\{\begin{array}{r} 3x - y = 8 \\ y + 2x = 7 \end{array}\right\}$ *Ans.* $x = 2,\ y = 3.$

6. $\left\{\begin{array}{r} 4x - 7y = -22 \\ 5x + 2y = 37 \end{array}\right\}$ *Ans.* $x = 5,\ y = 6.$

7. $\left\{\begin{array}{r} 2x + 6y = 42 \\ 8x - 6y = 3 \end{array}\right\}$ *Ans.* $x = 4\frac{1}{2},\ y = 5\frac{1}{2}.$

8. $\left\{\begin{array}{r} 8x - 9y = 1 \\ 6x - 3y = 4x \end{array}\right\}$ *Ans.* $x = \frac{1}{2},\ y = \frac{1}{3}.$

9. $\left\{\begin{array}{r} 14x - 15y = 12 \\ 7x + 8y = 37 \end{array}\right\}$ *Ans.* $x = 3,\ y = 2.$

10. $\left\{\begin{array}{r} \frac{1}{2}x + \frac{1}{3}y = 6 \\ \frac{1}{3}x + \frac{1}{2}y = 6\frac{1}{2} \end{array}\right\}$ *Ans.* $\left\{ x = 6,\ y = 9. \right.$

11. $\left\{\begin{array}{r} \frac{1}{7}x + \frac{1}{8}y = 4 \\ x - y = -2 \end{array}\right\}$ *Ans.* $\left\{ x = 14,\ y = 16. \right.$

---

113. What is the rule for elimination by addition or subtraction?

12. Says *A* to *B*, you give me \$40 of your money, and I shall then have five times as much as you will have left. Now they both had \$120: how much had each?

Ans. Each had \$60.

13 A father says to his son, "twenty years ago, my age was four times yours; now it is just double:" what were their ages?

Ans. { Father's, 60 years.
Son's, 30 years.

14. A father divided his property between his two sons. At the end of the first year the elder had spent one-quarter of his, and the younger had made \$1000, and their property was then equal. After this the elder spent \$500, and the younger made \$2000, when it appeared that the younger had just double the elder: what had each from the father?

Ans. { Elder, \$4000.
Younger, \$2000.

15. If John give Charles 15 apples, they will have the same number; but if Charles give 15 to John, John will have 15 times as many, wanting 10, as Charles will have left. How many has each?

Ans. { John, 50.
Charles, 20.

16. Two clerks, *A* and *B*, have salaries which are together equal to \$900. *A* spends $\frac{1}{10}$ per year of what he receives, and *B* adds as much to his as *A* spends. At the end of the year they have equal sums: what was the salary of each?

Ans. { *A's* = \$500.
B's = \$400.

Elimination by Substitution.

114. Let us again take the equations,

$$5x + 7y = 43, \qquad (1.)$$
$$11x + 9y = 69. \qquad (2.)$$

---

114. Give the rule for elimination by substitution. When is this method used to the greatest advantage?

Find the value of x in the first equation, which gives,

$$x = \frac{43 - 7y}{5}.$$

Substitute this value of x in the second equation, and we have,

$$11 \times \frac{43 - 7y}{5} + 9y = 69;$$

or, $$473 - 77y + 45y = 345;$$

or, $$-32y = -128.$$

Here, x has been eliminated by *substitution.*

In a similar manner, we can eliminate any unknown quantity; hence, the

RULE.

I. *Find from either equation the value of the unknown quantity to be eliminated:*

II. *Substitute this value for that quantity in the other equation.*

NOTE.—This method of elimination is used to great advantage when the coefficient of either of the unknown quantities is 1.

EXAMPLES.

Find, by the last method, the values of x and y in the following equations:

1. $3x - y = 1$, and $3y - 2x = 4$.
Ans. $x = 1$, $y = 2$

2. $5y - 4x = -22$, and $3y + 4x = 38$.
Ans. $x = 8$, $y = 2$

3. $x + 8y = 18$, and $y - 3x = -29$.
Ans. $x = 10$, $y = 1$

4. $5x - y = 13$, and $8x + \frac{2}{9}y = 29$.

Ans. $x = 3\frac{1}{2}$, $y = 4\frac{1}{2}$.

5. $10x - \frac{y}{5} = 69$, and $10y - \frac{x}{7} = 49$.

Ans. $x = 7$, $y = 5$.

6. $x + \frac{1}{2}x - \frac{y}{5} = 10$, and $\frac{x}{8} + \frac{y}{10} = 2$.

Ans. $x = 8$, $y = 10$.

7. $\frac{y}{7} - \frac{x}{3} + 5 = 2$, $x + \frac{y}{5} = 17\frac{4}{5}$.

Ans. $x = 15$, $y = 14$.

8. $\frac{y}{2} + \frac{x}{3} + 3 = 6\frac{1}{6}$, and $\frac{y}{4} - \frac{x}{7} = \frac{1}{2}$.

Ans. $x = 3\frac{1}{2}$, $y = 4$.

9. $\frac{y}{8} - \frac{x}{4} + 6 = 5$, and $\frac{x}{12} - \frac{y}{16} = 0$.

Ans. $x = 12$, $y = 16$.

10. $\frac{y}{7} - \frac{3x}{2} - 1 = -9$, and $5x - \frac{7y}{49} = 29$.

Ans. $x = 6$, $y = 7$.

11. Two misers, *A* and *B*, sit down to count over their money. They both have \$20000, and *B* has three times as much as *A*: how much has each?

Ans. { *A*, \$5000.
B, \$15000.

12. A person has two purses. If he puts \$7 into the first, the whole is worth three times as much as the second purse: but if he puts \$7 into the second, the whole is worth five times as much as the first: what is the value of each purse?

Ans. 1st, \$2; 2d, \$3.

13. Two numbers have the following relations: if the first be multiplied by 6, the product will be equal to the second multiplied by 5; and 1 subtracted from the first leaves the same remainder as 2 subtracted from the second: what are the numbers? *Ans.* 5 and 6.

14. Find two numbers with the following relations: the first increased by 2 is $3\frac{1}{4}$ times as great as the second; and the second increased by 4 gives a number equal to half the first: what are the numbers? *Ans.* 24 and 8.

15. A father says to his son, "twelve years ago, I was twice as old as you are now: four times your age at that time, plus twelve years, will express my age twelve years hence:" what were their ages?

Ans. { Father, 72 years.
Son, 30 "

Elimination by Comparison.

115. Take the same equations,

$$5x + 7y = 43$$
$$11x + 9y = 69.$$

Finding the value of x from the first equation, we have,

$$x = \frac{43 - 7y}{5};$$

and finding the value of x from the second, we obtain,

$$x = \frac{69 - 9y}{11}.$$

---

115. Give the rule for elimination by comparison.

Let these two values of x be placed equal to each other, and we have,

$$\frac{43 - 7y}{5} = \frac{69 - 9y}{11}.$$

Or, $$473 - 77y = 345 - 45y;$$

or, $$-32y = -128.$$

Hence, $$y = 4.$$

And, $$x = \frac{69 - 36}{11} = 3.$$

This method of elimination is called the method by *comparison*, for which we have the following

RULE.

I. *Find, from each equation, the value of the same unknown quantity to be eliminated:*

II. *Place these values equal to each other.*

EXAMPLES.

Find, by the last rule, the values of x and y, from the following equations,

1. $3x + \frac{y}{5} + 6 = 42$, and $y - \frac{x}{22} = 14\frac{1}{2}$.
Ans. $x = 11$, $y = 15$.

2. $\frac{y}{4} - \frac{x}{7} + 5 = 6$, and $\frac{y}{5} + 4 = \frac{x}{14} + 6$.
Ans. $x = 28$, $y = 20$.

3. $\frac{y}{10} - \frac{x}{4} + \frac{22}{8} = 1$, and $3y - x = 6$.
Ans. $x = 9$, $y = 5$.

4. $y - 3 = \frac{1}{2}x + 5$ and $\frac{x + y}{2} = y - 3\frac{1}{2}$.
Ans. $x = 2$, $y = 9$.

5. $\frac{y-x}{3}+\frac{x}{2}=y-2$, and $\frac{x}{8}+\frac{y}{7}=x-13$.

Ans. $x=16,\ y=7$

6. $\frac{y+x}{2}+\frac{y-x}{2}=x-\frac{2y}{3}$, and $x+y=16$.

Ans. $x=10,\ y=6$

7. $\frac{2x-3y}{5}=x-2\frac{2}{5}$, $x-\frac{y-1}{2}=0$.

Ans. $x=1,\ y=3$.

8. $2y+3x=y+43$, $y-\frac{x-4}{3}=y-\frac{x}{5}$.

Ans. $x=10,\ y=13$.

9. $4y-\frac{x-y}{2}=x+18$, and $27-y=x+y+4$.

Ans. $x=9,\ y=7$.

10. $1-\frac{y-x}{6}+4=y-16\frac{2}{3}$, $\frac{y}{5}-2=\frac{x}{5}$.

Ans. $x=10,\ y=20$.

116. Having explained the principal methods of elimination, we shall add a few examples which may be solved by any one of them; and often indeed, it may be advantageous to employ them all, even in the same example.

GENERAL EXAMPLES.

Find the values of x and y in the following simultaneous equations:

1. $2x+3y=16$, and $3x-2y=11$.

Ans. $x=5,\ y=2$.

2. $\frac{2x}{5} + \frac{3y}{4} = \frac{9}{20}$, and $\frac{3x}{4} + \frac{2y}{5} = \frac{61}{120}$.

Ans. $x = \frac{1}{2}$, $y = \frac{1}{3}$.

3. $\frac{x}{7} + 7y = 99$, and $\frac{y}{7} + 7x = 51$.

Ans. $x = 7$, $y = 14$.

4. $\frac{x}{2} - 12 = \frac{y}{4} + 8$, $\frac{x+y}{5} + \frac{x}{3} - 8 = \frac{2y-x}{4} + 27$.

Ans. $x = 60$, $y = 40$.

5. $$\begin{cases} x - \frac{1}{2}y + \frac{4x}{5} = 6\frac{4}{5} \\ \frac{x-y}{2} + 7x = 41 \end{cases}$$ *Ans.* $\begin{cases} x = 6. \\ y = 8. \end{cases}$

6. $$\begin{cases} \frac{x-y}{4} + \frac{x+y}{5} = 2\frac{1}{10} \\ \frac{1}{2}x - y + 4\frac{1}{4}y = 12\frac{1}{4} \end{cases}$$ *Ans.* $\begin{cases} x = 5. \\ y = 3. \end{cases}$

7. $$\begin{cases} \frac{3y-x}{6} + \frac{2x-y}{4} = 5 \\ 6x - y + \frac{8-2x}{4} = 43\frac{1}{2} \end{cases}$$ *Ans.* $\begin{cases} x = 9. \\ y = 8. \end{cases}$

8. $$\begin{cases} \frac{3x-8}{4} + \frac{y-6}{5} + y = 18\frac{7}{10} \\ 8x - 3 - \frac{6-y}{3} = 79 \end{cases}$$ *Ans.* $\begin{cases} x = 10. \\ y = 12. \end{cases}$

9. $$\begin{cases} \frac{4x-4}{3} - \frac{y-5}{4} + 6 = 12\frac{2}{3} \\ \frac{1}{2}x - \frac{1}{3}y + \frac{y-4}{3} = \frac{5}{3} \end{cases}$$ *Ans.* $\begin{cases} x = 6. \\ y = 5. \end{cases}$

$$10.\ \left\{\begin{array}{l} ax - by = c \\ a - y + x = d \end{array}\right\} \quad Ans. \left\{\begin{array}{l} x = \dfrac{c + ab - bd}{a - b}. \\ y = \dfrac{a^2 + c - ad}{a - b}. \end{array}\right.$$

$$11.\ \left\{\begin{array}{l} 13x + 7y - 341 = 7\tfrac{1}{2}y + 43\tfrac{1}{2}x \\ 2x + \tfrac{1}{2}y = 1 \end{array}\right\} \quad Ans. \left\{\begin{array}{l} x = -12. \\ y = 50. \end{array}\right.$$

$$12.\ \left\{\begin{array}{l} (x+5)(y+7) = (x+1)(y-9) + 112 \\ 2x + 10 = 3y + 1 \end{array}\right\} \quad Ans. \left\{\begin{array}{l} x = 3. \\ y = 5. \end{array}\right.$$

$$13.\ \left\{\begin{array}{l} ax = by \\ x + y = c \end{array}\right\} \quad Ans. \left\{\begin{array}{l} x = \dfrac{bc}{a + b}. \\ y = \dfrac{ac}{a + b}. \end{array}\right.$$

$$14.\ \left\{\begin{array}{l} ax + by = c \\ fx + gy = h \end{array}\right\} \quad Ans. \left\{\begin{array}{l} x = \dfrac{cg - bh}{ag - bf} \\ y = \dfrac{ah - cf}{ag - bf}. \end{array}\right.$$

$$15.\ \left\{\begin{array}{l} \dfrac{a}{b + y} = \dfrac{b}{3a + x} \\ ax + 2by = d \end{array}\right\} \quad Ans. \left\{\begin{array}{l} x = \dfrac{2b^2 - 6a^2 + d}{3a}. \\ y = \dfrac{3a^2 - b^2 + d}{3b}. \end{array}\right.$$

$$16.\ \left\{\begin{array}{l} bcx = cy - 2b \\ b^2y + \dfrac{a(c^3 - b^3)}{bc} = \dfrac{2b^3}{c} + c^3x \end{array}\right\} \quad Ans. \left\{\begin{array}{l} x = \dfrac{a}{bc}. \\ y = \dfrac{a + 2b}{c} \end{array}\right.$$

$$17.\ \left\{\begin{array}{l} 3x + 5y = \dfrac{(8b - 2f)bf}{b^2 - f^2} \\ y - x = \dfrac{-2bf^2}{b^2 - f^2} \end{array}\right\} \quad Ans. \left\{\begin{array}{l} x = \dfrac{bf}{b - f} \\ y = \dfrac{bf}{b + f}. \end{array}\right.$$

PROBLEMS.

1. What fraction is that, to the numerator of which if 1 be added, the value will be $\frac{1}{3}$, but if 1 be added to its denominator, the value will be $\frac{1}{4}$?

Let the fraction be denoted by $\frac{x}{y}$.

Then, by the conditions,

$$\frac{x+1}{y} = \frac{1}{3}, \text{ and, } \frac{x}{y+1} = \frac{1}{4}.$$

whence, $3x + 3 = y$, and $4x = y + 1$.

Therefore, by subtracting,

$$x - 3 = 1, \text{ and } x = 4.$$

Hence, $12 + 3 = y$;

$$\therefore\ y = 15.$$

2. A market-woman bought a certain number of eggs at 2 for a penny, and as many others at 3 for a penny; and having sold them all together, at the rate of 5 for 2*d*, found that she had lost 4*d*: how many of both kinds did she buy?

Let $2x$ denote the whole number of eggs.

Then, $x =$ the number of eggs of each sort.

Then will, $\frac{1}{2}x =$ the cost of the first sort,

and, $\frac{1}{3}x =$ the cost of the second sort.

But, by the conditions of the question,

$$5 : 2x :: 2 : \frac{4x}{5};$$

hence, $\frac{4x}{5}$ will denote the amount for which the eggs were sold.

But, by the conditions,

$$\frac{1}{2}x + \frac{1}{3}x - \frac{4x}{5} = 4;$$

therefore, $$15x + 10x - 24x = 120;$$

$$\therefore\ x = 120;\ \text{the number of eggs of each sort.}$$

3. A person possessed a capital of 30,000 dollars, for which he received a certain interest; but he owed the sum of 20,000 dollars, for which he paid a certain annual interest. The interest that he received exceeded that which he paid by 800 dollars. Another person possessed 35,000 dollars, for which he received interest at the second of the above rates; but he owed 24,000 dollars, for which he paid interest at the first of the above rates. The interest that he received, annually, exceeded that which he paid, by 310 dollars. Required the two rates of interest.

Let x denote the number of units in the first rate of interest, and y the unit in the second rate. Then each may be regarded as denoting the interest on $100 for 1 year.

To obtain the interest of $30,000 at the first rate, denoted by x, we form the proportion,

$$100 : 30{,}000 :: x : \frac{30{,}000x}{100},\ \text{or } 300x.$$

And for the interest of $20,000, the rate being y,

$$100 : 20{,}000 :: y : \frac{20{,}000y}{100},\ \text{or } 200y.$$

But, by the conditions, the difference between these two amounts is equal to 800 dollars.

We have, then, for the first equation of the problem,

$$300x - 200y = 800$$

By expressing, algebraically, the second condition of the problem, we obtain a second equation,

$$350y - 240x = 310.$$

Both members of the first equation being divisible by 100 and those of the second by 10, we have,

$$3x - 2y = 8, \qquad 35y - 24x = 31.$$

To eliminate x, multiply the first equation by 8, and then add the result to the second; there results,

$$19y = 95, \quad \text{whence,} \quad y = 5.$$

Substituting for y, in the first equation, this value, and that equation becomes,

$$3x - 10 = 8, \quad \text{whence,} \quad x = 6.$$

Therefore, the first rate is 6 per cent, and the second 5.

VERIFICATION.

$30,000,	at 6 per cent, gives	30,000 × .06	= $1800.
$20,000,	5 " "	20,000 × .05	= $1000.

And we have, $1800 - 1000 = 800.$

The second condition can be verified in the same manner.

4. What two numbers are those, whose difference is 7, and sum 33? *Ans.* 13 and 20.

5. Divide the number 75 into two such parts, that three times the greater may exceed seven times the less by 15. *Ans.* 54 and 21.

6. In a mixture of wine and cider, $\frac{1}{2}$ of the whole plus 25 gallons was wine, and $\frac{1}{3}$ part minus 5 gallons was cider: how many gallons were there of each? *Ans.* 85 of wine, and 35 of cider.

7. A bill of £120 was paid in guineas and moidores, and the number of pieces used, of both sorts, was just 100. If the guinea be estimated at 21*s*, and the moidore at 27*s*, how many pieces were there of each sort? *Ans.* 50.

8. Two travelers set out at the same time from London and York, whose distance apart is 150 miles. One of them travels 8 miles a day, and the other 7: in what time will they meet? *Ans.* In 10 days.

9. At a certain election, 375 persons voted for two candidates, and the candidate chosen had a majority of 91: how many voted for each?

Ans. 233 for one, and 142 for the other.

10. A person has two horses, and a saddle worth £50. Now, if the saddle be put on the back of the first horse, it makes their joint value double that of the second horse; but if it be put on the back of the second, it makes their joint value triple that of the first: what is the value of each horse? *Ans.* One £30, and the other £40.

11. The hour and minute hands of a clock are exactly together at 12 o'clock: when will they be again together?

Ans. 1h. $5\frac{5}{11}$m.

12. A man and his wife usually drank out a cask of beer in 12 days; but when the man was from home, it lasted the woman 30 days: how many days would the man alone be in drinking it? *Ans.* 20 days.

13. If 32 pounds of sea-water contain 1 pound of salt, how much fresh water must be added to these 32 pounds, in order that the quantity of salt contained in 32 pounds of the new mixture shall be reduced to 2 ounces, or $\frac{1}{8}$ of a pound?

Ans. 224 lbs.

14. A person who possessed 100,000 dollars, placed the greater part of it out at 5 per cent interest, and the other

at 4 per cent. The interest which he received for the whole, amounted to 4640 dollars. Required the two parts.

Ans. $64,000 and $36,000.

15. At the close of an election, the successful candidate had a majority of 1500 votes. Had a fourth of the votes of the unsuccessful candidate been also given to him, he would have received three times as many as his competitor, wanting three thousand five hundred: how many votes did each receive?

Ans. { 1st, 6500.
2d, 5000.

16. A gentleman bought a gold and a silver watch, and a chain worth $25. When he put the chain on the gold watch, it and the chain became worth three and a half times more than the silver watch; but when he put the chain on the silver watch, they became worth one-half the gold watch and 15 dollars over: what was the value of each watch?

Ans. { Gold watch, $80.
Silver " $30.

17. There is a certain number expressed by two figures, which figures are called digits. The sum of the digits is 11, and if 13 be added to the first digit the sum will be three times the second: what is the number? *Ans.* 56.

18. From a company of ladies and gentlemen 15 ladies retire; there are then left two gentlemen to each lady. After which 45 gentlemen depart, when there are left 5 ladies to each gentleman: how many were there of each at first?

Ans. { 50 gentlemen.
40 ladies.

19. A person wishes to dispose of his horse by lottery. If he sells the tickets at $2 each, he will lose $30 on his horse; but if he sells them at $3 each, he will receive $30

more than his horse cost him. What is the value of the horse, and number of tickets? *Ans.* $\left\{\begin{array}{ll}\text{Horse,} & \$150.\\ \text{No. of tickets,} & 60.\end{array}\right.$

20. A person purchases a lot of wheat at $1, and a lot of rye at 75 cents per bushel; the whole costing him $117.50. He then sells $\frac{1}{4}$ of his wheat and $\frac{1}{5}$ of his rye at the same rate, and realizes $27.50. How much did he buy of each?

Ans. $\left\{\begin{array}{l}\text{80 bush. of wheat.}\\ \text{50 bush. of rye.}\end{array}\right.$

21. There are 52 pieces of money in each of two bags. *A* takes from one, and *B* from the other. *A* takes twice as much as *B* left, and *B* takes 7 times as much as *A* left. How much did each take? *Ans.* $\left\{\begin{array}{l}A\text{, 48 pieces.}\\ B\text{, 28 pieces.}\end{array}\right.$

22. Two persons, *A* and *B*, purchase a house together, worth $1200. Says *A* to *B*, give me two-thirds of your money and I can purchase it alone; but, says *B* to *A*, if you will give me three-fourths of your money I shall be able to purchase it alone. How much had each?

Ans. *A*, $800; *B*, $600.

23. A grocer finds that if he mixes sherry and brandy in the proportion of 2 to 1, the mixture will be worth 78*s.* per dozen; but if he mixes them in the proportion of 7 to 2, he can get 79*s.* a dozen. What is the price of each liquor per dozen? *Ans.* Sherry, 81*s.*; brandy, 72*s.*

Equations containing three or more unknown quantities.

117. Let us now consider equations involving three or more unknown quantities.

Take the group of simultaneous equations,

---

117. Give the rule for solving any group of simultaneous equations?

$$5x - 6y + 4z = 15, \quad . \ . \quad (1.)$$
$$7x + 4y - 3z = 19, \quad . \ . \quad (2.)$$
$$2x + y + 6z = 46. \quad . \ . \ . \quad (3.)$$

To eliminate z by means of the first two equations, multiply the first by 3, and the second by 4; then, since the coefficients of z have contrary signs, add the two results together. This gives a new equation:

$$43x - 2y = 121 \ . \ . \ . \ . \ . \quad (4.)$$

Multiplying the second equation by 2 (a factor of the coefficient of z in the third equation), and adding the result to the third equation, we have,

$$16x + 9y = 84 \ . \ . \ . \ . \ . \quad (5.)$$

The question is then reduced to finding the values of x and y, which will satisfy the new Equations (4) and (5).

Now, if the first be multiplied by 9, the second by 2, and the results added together, we find,

$$419x = 1257; \text{ whence, } x = 3.$$

We might, by means of Equations (4) and (5) determine y in the same way that we have determined x; but the value of y may be determined more simply, by substituting the value of x in Equation (5); thus,

$$48 + 9y = 84. \qquad \therefore \ y = \frac{84 - 48}{9} = 4.$$

In the same manner, the first of the three given equations becomes, by substituting the values of x and y,

$$15 - 24 + 4z = 15 \qquad \therefore \ z = \frac{24}{4} = 6.$$

In the same way, any group of simultaneous equations may be solved Hence, the

RULE.

I. *Combine one equation of the group with each of the others, by eliminating one unknown quantity; there will result a new group containing one equation less than the original group:*

II. *Combine one equation of this new group with each of the others, by eliminating a second unknown quantity; there will result a new group containing two equations less than the original group:*

III. *Continue the operation until a single equation is found, containing but one unknown quantity:*

IV. *Find the value of this unknown quantity by the preceding rules; substitute this in one of the group of two equations, and find the value of a second unknown quantity; substitute these in either of the group of three, finding a third unknown quantity; and so on, till the values of all are found.*

NOTES.—1. In order that the value of the unknown quantities may be determined, there must be just as many independent equations of condition as there are unknown quantities. If there are fewer equations than unknown quantities, the resulting equation will contain at least two unknown quantities, and hence, their values cannot be found (Art. 110). If there are more equations than unknown quantities, the conditions may be contradictory, and the equations impossible.

2. It often happens that each of the proposed equations does not contain all the unknown quantities. In this case, with a little address, the elimination is very quickly performed.

Take the four equations involving four unknown quantities:

$$2x - 3y + 2z = 13. \quad (1.) \qquad 4y + 2z = 14. \quad (3.)$$

$$4u - 2x = 30. \quad (2.) \qquad 5y + 3u = 32. \quad (4.)$$

By inspecting these equations, we see that the elimination of z in the two Equations, (1) and (3), will give an equation involving x and y; and if we eliminate u in Equations (2) and (4), we shall obtain a second equation, involving x and y. These last two unknown quantities may therefore be easily determined. In the first place, the elimination of z from (1) and (3) gives,

$$7y - 2x = 1;$$

That of u from (2) and (4) gives,

$$20y + 6x = 38.$$

Multiplying the first of these equations by 3, and adding,

$$41y = 41;$$

Whence, $y = 1.$

Substituting this value in $7y - 2x = 1$, we find,

$$x = 3.$$

Substituting for x its value in Equation (2), it becomes

$$4u - 6 = 30.$$

Whence, $u = 9.$

And substituting for y its value in Equation (3), there results,

$$z = 5.$$

EXAMPLES.

1. Given $\left\{\begin{array}{r} x + y + z = 29 \\ x + 2y + 3z = 62 \\ \frac{1}{2}x + \frac{1}{3}y + \frac{1}{4}z = 10 \end{array}\right\}$ to find x, y, and z.

Ans. $x = 8$, $y = 9$, $z = 12$.

2. Given $\left.\begin{array}{r}2x + 4y - 3z = 22\\ 4x - 2y + 5z = 18\\ 6x + 7y - z = 63\end{array}\right\}$ to find x, y, and z.

Ans. $x = 3,\ y = 7,\ z = 4.$

3. Given $\left.\begin{array}{r}x + \frac{1}{2}y + \frac{1}{3}z = 32\\ \frac{1}{3}x + \frac{1}{4}y + \frac{1}{5}z = 15\\ \frac{1}{4}x + \frac{1}{5}y + \frac{1}{6}z = 12\end{array}\right\}$ to find x, y, and z.

Ans. $x = 12,\ y = 20,\ z = 30.$

4. Given $\left.\begin{array}{r}x + y + z = 29\frac{1}{4}\\ x + y - z = 18\frac{1}{4}\\ x - y + z = 13\frac{3}{4}\end{array}\right\}$ to find x, y, and z.

Ans. $x = 16,\ y = 7\frac{3}{4},\ z = 5\frac{1}{2}.$

5. Given $\left.\begin{array}{l}3x + 5y = 161\\ 7x + 2z = 209\\ 2y + z = 89\end{array}\right\}$ to find x, y, and z.

Ans. $x = 17,\ y = 22,\ z = 45.$

6. Given $\left.\begin{array}{l}\frac{1}{x} + \frac{1}{y} = a\\ \frac{1}{x} + \frac{1}{z} = b\\ \frac{1}{y} + \frac{1}{z} = c\end{array}\right\}$ to find x, y, and z.

$$x = \frac{2}{a + b - c},\quad y = \frac{2}{a + c - b},\quad z = \frac{2}{b + c - a}.$$

NOTE.—In this example we should not proceed to clear the equation of fractions; but subtract immediately the second equation from the first, and then add the third: we thus find the value of y.

PROBLEMS.

1. Divide the number 90 into four such parts, that the first increased by 2, the second diminished by 2, the third multiplied by 2, and the fourth divided by 2, shall be equal each to each.

This problem may be easily solved by introducing a new unknown quantity.

Let x, y, z, and u, denote the required parts, and designate by m the several equal quantities which arise from the conditions. We shall then have,

$$x + 2 = m, \quad y - 2 = m, \quad 2z = m, \quad \frac{u}{2} = m.$$

From which we find,

$$x = m - 2, \quad y = m + 2, \quad z = \frac{m}{2}, \quad u = 2m.$$

And, by adding the equations,

$$x + y + z + u = m + m + \frac{m}{2} + 2m = 4\tfrac{1}{2}m.$$

And since, by the conditions of the problem, the first member is equal to 90, we have,

$$4\tfrac{1}{2}m = 90, \quad \text{or} \quad \tfrac{9}{2}m = 90;$$

hence, $$m = 20.$$

Having the value of m, we easily find the other values; viz.:

$$x = 18, \quad y = 22, \quad z = 10, \quad u = 40.$$

2. There are three ingots, composed of different metals mixed together. A pound of the first contains 7 ounces of silver, 3 ounces of copper, and 6 of pewter. A pound of the second contains 12 ounces of silver, 3 ounces of copper, and 1 of pewter. A pound of the third contains 4 ounces of silver, 7 ounces of copper, and 5 of pewter. It is required

to find how much it will take of each of the three ingots to form a fourth, which shall contain in a pound, 8 ounces of silver, $3\frac{3}{4}$ of copper, and $4\frac{1}{4}$ of pewter.

Let x, y, and z, denote the number of ounces which it is necessary to take from the three ingots respectively, in order to form a pound of the required ingot. Since ther are 7 ounces of silver in a pound, or 16 ounces, of the first ingot, it follows that one ounce of it contains $\frac{7}{16}$ of an ounce of silver, and, consequently, in a number of ounces denoted by x, there is $\frac{7x}{16}$ ounces of silver. In the same manner, we find that, $\frac{12y}{16}$, and $\frac{4z}{16}$, denote the number of ounces of silver taken from the second and third; but, from the enunciation, one pound of the fourth ingot contains 8 ounces of silver. We have, then, for the first equation,

$$\frac{7x}{16} + \frac{12y}{16} + \frac{4z}{16} = 8;$$

or, clearing fractions,

$$7x + 12y + 4z = 128.$$

As respects the copper, we should find,

$$3x + 3y + 7z = 60;$$

and with reference to the pewter,

$$6x + y + 5z = 68.$$

As the coefficients of y in these three equations are the most simple, it is convenient to eliminate this unknown quantity first.

Multiplying the second equation by 4, and subtracting the first from it, member from member, we have,

$$5x + 24z = 112.$$

Multiplying the third equation by 3, and subtracting the second from the resulting equation, we have,

$$15x + 8z = 144.$$

Multiplying this last equation by 3, and subtracting the preceding one, we obtain,

$$40x = 320;$$

whence, $$x = 8.$$

Substitute this value for x in the equation,

$$15x + 8z = 144;$$

it becomes, $$120 + 8z = 144,$$

whence, $$z = 3.$$

Lastly, the two values, $x = 8$, $z = 3$, being substituted in the equation,

$$6x + y + 5z = 68,$$

give, $$48 + y + 15 = 68,$$

whence, $$y = 5.$$

Therefore, in order to form a pound of the fourth ingot, we must take 8 ounces of the first, 5 ounces of the second, and 3 of the third.

VERIFICATION.

If there be 7 ounces of silver in 16 ounces of the first ingot, in eight ounces of it there should be a number of ounces of silver expressed by

$$\frac{7 \times 8}{16}.$$

In like manner,

$$\frac{12 \times 5}{16}, \text{ and } \frac{4 \times 3}{16},$$

will express the quantity of silver contained in 5 ounces of the second ingot. and 3 ounces of the third.

Now, we have,

$$\frac{7 \times 8}{16} + \frac{12 \times 5}{16} + \frac{4 \times 3}{16} = \frac{128}{16} = 8;$$

therefore, a pound of the fourth ingot contains 8 ounces of silver, as required by the enunciation. The same conditions may be verified with respect to the copper and pewter.

3. *A's* age is double *B's*, and *B's* is triple of *C's*, and the sum of all their ages is 140: what is the age of each?

Ans. *A's* = 84; *B's* = 42; and *C's* = 14.

4. A person bought a chaise, horse, and harness, for £60; the horse came to twice the price of the harness, and the chaise to twice the cost of the horse and harness: what did he give for each?

$$\textit{Ans.} \left\{ \begin{array}{ll} \pounds 13\ \ 6s.\ 8d. & \text{for the horse.} \\ \pounds 6\ \ 13s.\ 4d. & \text{for the harness.} \\ \pounds 40 & \text{for the chaise.} \end{array} \right.$$

5. Divide the number 36 into three such parts that $\frac{1}{2}$ of the first, $\frac{1}{3}$ of the second, and $\frac{1}{4}$ of the third, may be all equal to each other. *Ans.* 8, 12, and 16.

6. If *A* and *B* together can do a piece of work in 8 days, *A* and *C* together in 9 days, and *B* and *C* in ten days, how many days would it take each to perform the same work alone? *Ans.* *A*, $14\frac{34}{49}$; *B*, $17\frac{23}{41}$; *C*, $23\frac{7}{31}$.

7. Three persons, *A*, *B*, and *C*, begin to play together, having among them all $600. At the end of the first game *A* has won one-half of *B's* money, which, added to his own, makes double the amount *B* had at first. In the second game, *A* loses and *B* wins just as much as *C* had at the beginning, when *A* leaves off with exactly what he had at first: now much had each at the beginning?

Ans. *A*, $300; *B*, $200; *C* $100.

8. Three persons, *A*, *B*, and *C*, together possess $3640.

If *B* gives *A* $400 of his money, then *A* will have $320 more than *B*; but if *B* takes $140 of *C's* money, then *B* and *C* will have equal sums: how much has each?

Ans. *A*, $800; *B*, $1280; *C*, $1560.

9. Three persons have a bill to pay, which neither alone is able to discharge. *A* says to *B*, "Give me the 4th of your money, and then I can pay the bill." *B* says to *C* "Give me the 8th of yours, and I can pay it." But *C* says to *A*, "You must give me the half of yours before I can pay it, as I have but $8": what was the amount of their bill, and how much money had *A* and *B*?

Ans. { Amount of the bill, $13.
A had $10, and *B* $12.

10. A person possessed a certain capital, which he placed out at a certain interest. Another person, who possessed 10000 dollars more than the first, and who put out his capital 1 per cent. more advantageously, had an annual income greater by 800 dollars. A third person, who possessed 15000 dollars more than the first, putting out his capital 2 per cent. more advantageously, had an annual income greater by 1500 dollars. Required, the capitals of the three persons, and the rates of interest.

Ans.				
	Sums at interest,	$30000,	$40000,	$45000.
	Rates of interest,	4	5	6 pr. ct.

11. A widow receives an estate of $15000 from her deceased husband, with directions to divide it among two sons and three daughters, so that each son may receive twice as much as each daughter, and she herself to receive $1000 more than all the children together: what was her share, and what the share of each child?

Ans.		
	The widow's share,	$8000
	Each son's,	$2000
	Each daughter's,	$1000

12. A certain sum of money is to be divided between three persons, *A*, *B*, and *C*. *A* is to receive $3000 less than half of it, *B* $1000 less than one-third part, and *C* to receive $800 more than the fourth part of the whole: what is the sum to be divided, and what does each receive?

Ans. Sum, $38400.
A receives $16200
B " $11800.
C " $10400.

13. A person has three horses, and a saddle which is worth $220. If the saddle be put on the back of the first horse, it will make his value equal to that of the second and third; if it be put on the back of the second, it will make his value double that of the first and third; if it be put on the back of the third, it will make his value triple that of the first and second: what is the value of each horse?

Ans. 1st, $20; 2d, $100; 3d, $140.

14. The crew of a ship consisted of her complement of sailors, and a number of soldiers. There were 22 sailors to every three guns, and 10 over; also, the whole number of hands was five times the number of soldiers and guns together. But after an engagement, in which the slain were one-fourth of the survivors, there wanted 5 men to make 13 men to every two guns: required, the number of guns, soldiers and sailors.

Ans. 90 guns, 55 soldiers, and 670 sailors.

15. Three persons have $96, which they wish to divide equally between them. In order to do this, *A*, who has the most, gives to *B* and *C* as much as they have already; then *B* divides with *A* and *C* in the same manner, that is, by giving to each as much as he had after *A* had divided with them. *C* then makes a division with *A* and *B*, when it is

found that they all have equal sums: how much had each at first? *Ans.* 1st, \$52; 2d, \$28; 3d, \$16.

16. Divide the number a into three such parts, that the first shall be to the second as m to n, and the second to the third as p to q.

$$a = \frac{amp}{mp+np+nq}, \quad y = \frac{anp}{mp+np+nq}, \quad z = \frac{anq}{mp+np+nq}$$

17. Three masons, *A*, *B*, and *C*, are to build a wall. *A* and *B* together can do it in 12 days; *B* and *C* in 20 days; and *A* and *C* in 15 days: in what time can each do it alone, and in what time can they all do it if they work together? *Ans.* *A*, in 20 days; *B*, in 30; and *C*, in 60; all, in 10.

CHAPTER VI.

FORMATION OF POWERS

118. A **Power** of a quantity is the product obtained by taking that quantity any number of times as a factor.

If the quantity be taken once as a factor, we have the first power; if taken twice, we have the second power; if three times, the third power; if n times, the n^{th} power, n being any whole number whatever.

A power is indicated by means of the exponential sign thus,

$a = a^1$ denotes first power of a.*
$a \times a = a^2$ " square, or 2d power of a.
$a \times a \times a = a^3$ " cube, or third power of a.
$a \times a \times a \times a = a^4$ " fourth power of a.
$a \times a \times a \times a \times a = a^5$ " fifth power of a.
$a \times a \times a \times a .... = a^m$ " m^{th} power of a.

In every power there are three things to be considered:

1st. The quantity which enters as a factor, and which is called the first power.

2d. The small figure which is placed at the right, and a little above the letter, is called the *exponent* of the

---

[*] Since $a^0 = 1$ (Art. 49), $a^0 \times a = 1 \times a = a^1$; so that the two factors of a^1, are 1 and a.

---

118. What is a power of a quantity? What is the power when the quantity is taken once as a factor? When taken twice? Three times? n times? How is a power indicated? In every power, how many things are considered? Name them.

power, and shows how many times the letter enters as a factor.

3d. The power itself, which is the final product, or result of the multiplications.

POWERS OF MONOMIALS.

119. Let it be required to raise the monomial $2a^3b^2$ to the fourth power. We have,

$$(2a^3b^2)^4 = 2a^3b^2 \times 2a^3b^2 \times 2a^3b^2 \times 2a^3b^2,$$

which merely expresses that the fourth power is equal to the product which arises from taking the quantity four times as a factor. By the rules for multiplication, this product is

$$(2a^3b^2)^4 = 2^4a^{3+3+3+3}b^{2+2+2+2} = 2^4a^{12}b^8;$$

from which we see,

1st. That the coefficient 2 must be raised to the 4th power; and,

2d. That the exponent of each letter must be multiplied by 4, the exponent of the power.

As the same reasoning applies to every example, we have, for the raising of monomials to any power, the following

RULE.

I. *Raise the coefficient to the required power:*

II. *Multiply the exponent of each letter by the exponent of the power.*

EXAMPLES.

1. What is the square of $3a^2y^3$? *Ans.* $9a^4y^6$

---

119. What is the rule for raising a monomial to any power? When the monomial is positive, what will be the sign of its powers? When negative, what powers will be plus? what minus?

2. What is the cube of $6a^5y^2x$? *Ans.* $216a^{15}y^6x^3$.

3. What is the fourth power of $2a^3y^3b^5$? $16a^{12}y^{12}b^{20}$.

4. What is the square of $a^2b^5y^3$? *Ans.* $a^4b^{10}y^6$.

5. What is the seventh power of a^2bcd^3?
Ans. $a^{14}b^7c^7d^{21}$

6. What is the sixth power of $a^2b^3c^2d$?
Ans. $a^{12}b^{18}c^{2}d^6$.

7. What is the square and cube of $-2a^2b^2$?

Square.	*Cube.*
$-2a^2b^2$	$-2a^2b^2$
$-2a^2b^2$	$-2a^2b^2$
$+4a^4b^4$.	$+4a^4b^4$
	$-2a^2b^2$
	$-8a^6b^6$.

By observing the way in which the powers are formed, we may conclude,

1st. *When the monomial is positive, all the powers will be positive.*

2d. *When the monomial is negative, all even powers will be positive, and all odd will be negative.*

8. What is the square of $-2a^4b^5$? *Ans.* $4a^8b^{10}$.

9. What is the cube of $-5a^nb^2$? *Ans.* $-125a^{3n}b^6$.

10. What is the eighth power of $-a^3xy^2$?
Ans. $+a^{24}x^8y^{16}$.

11. What is the seventh power of $-a^mb^nc$?
Ans. $-a^{7m}b^{7n}c^7$.

12. What is the sixth power of $2ab^6y^5$?
Ans. $64a^6b^{36}y^{30}$.

13. What is the ninth power of $-a^nbc^2$? *Ans.* $-a^{9n}b^9c^{18}$.

14. What is the sixth power of $-3ab^2d$? *Ans.* $729a^6b^{12}d^6$

15. What is the square of $-10a^mb^nc^3$? *Ans.* $100a^{2m}b^{2n}c^6$

16. What is the cube of $-9a^mb^nd^3f^2$? *Ans.* $-729a^{3m}b^{3n}d^9f^6$.

17. What is the fourth power of $-4a^5b^3c^4d^5$? *Ans.* $256a^{20}b^{12}c^{16}d^{20}$

18. What is the cube of $-4a^{2m}b^{2n}c^3d$? *Ans.* $-64a^{6m}b^{6n}c^9d^3$

19. What is the fifth power of $2a^3b^2xy$? *Ans.* $32a^{15}b^{10}x^5y^5$.

20. What is the square of $20x^ny^mc^5$? *Ans.* $400x^{2n}y^{2m}c^{10}$.

21. What is the fourth power of $3a^nb^{2n}c^3$? *Ans.* $81a^{4n}b^{8n}c^{12}$.

22. What is the fifth power of $-c^nd^{3m}x^2y^2$? *Ans.* $-c^{5n}d^{15m}x^{10}y^{10}$.

23. What is the sixth power of $-a^nb^{2n}c^m$? *Ans.* $a^{6n}b^{12n}c^{6m}$.

24. What is the fourth power of $-2a^2c^2d^3$. *Ans.* $16a^8c^8d^{12}$.

POWERS OF FRACTIONS.

120. From the definition of a power, and the rule for the multiplication of fractions, the cube of the fraction $\frac{a}{b}$, is written,

$$\left(\frac{a}{b}\right)^3 = \frac{a}{b} \times \frac{a}{b} \times \frac{a}{b} = \frac{a^3}{b^3};$$

---

120. What is the rule for raising a fraction to any power?

and since any fraction raised to any power, may be written under the same form, we find any power of a fraction by the following

RULE.

Raise the numerator to the required power for a new numerator, and the denominator to the required power for a new denominator.

The rule for signs is the same as in the last article.

EXAMPLES

Find the powers of the following fractions:

1. $\left(\frac{a-c}{b+c}\right)^2$. *Ans.* $\frac{a^2-2ac+c^2}{b^2+2bc+c^2}$.

2. $\left(\frac{xy}{3bc}\right)^3$. *Ans.* $\frac{x^3y^3}{27b^3c^3}$.

3. $\left(\frac{-x^2y}{2ab}\right)^4$. *Ans.* $\frac{x^8y^4}{16a^4b^4}$.

4. $\left(\frac{2ax^2y}{3bc^2}\right)^2$. *Ans.* $\frac{4a^2x^4y^2}{9b^2c^4}$.

5. $\left(-\frac{dx}{3y^2}\right)^3$. *Ans.* $-\frac{d^3x^3}{27y^6}$.

6. $\left(\frac{axy^3}{2bz^2}\right)^3$. *Ans.* $\frac{a^3x^3y^9}{8b^3z^6}$.

7. $\left(-\frac{3ay^4}{2b^2x}\right)^4$. *Ans.* $\frac{81a^4y^{16}}{16b^8x^4}$.

8. Fourth power of $\frac{ab^2c}{2x^2y^2}$ *Ans.* $\frac{a^4b^8c^4}{16x^8y^8}$.

9. Cube of $\frac{x-y}{x+y}$. *Ans.* $\frac{x^3-3x^2y+3xy^2-y^3}{x^3+3x^2y+3xy^2+y^3}$.

10. Fourth power of $\frac{2a^mx^n}{4a^py^q}$. *Ans.* $\frac{a^{4m}x^{4n}}{16a^{4p}y^{4q}}$.

11. Fifth power of $-\frac{9bc^nx^m}{18y^pz^q}$. *Ans.* $-\frac{b^5c^{5n}x^{5m}}{32y^{5p}z^{5q}}$.

POWERS OF BINOMIALS.

121. A Binomial, like a monomial, may be raised to any power by the process of continued multiplication.

1. Find the fifth power of the binomial $a + b$.

$$
\begin{array}{llllllll}
a & +\ b & & & & & \ldots\ldots\ldots\ldots & \text{1st power.} \\
a & +\ b & & & & & & \\
\hline
a^2 & +\ ab & & & & & & \\
 & +\ ab & +\ b^2 & & & & & \\
\hline
a^2 & +\ 2ab & +\ b^2 & & & & \ldots\ldots\ldots & \text{2d power.} \\
a & +\ b & & & & & & \\
\hline
a^3 & +\ 2a^2b & +\ ab^2 & & & & & \\
 & +\ a^2b & +\ 2ab^2 & +\ b^3 & & & & \\
\hline
a^3 & +\ 3a^2b & +\ 3ab^2 & +\ b^3 & & & \ldots\ldots & \text{3d power.} \\
a & +\ b & & & & & & \\
\hline
a^4 & +\ 3a^3b & +\ 3a^2b^2 & +\ ab^3 & & & & \\
 & +\ a^3b & +\ 3a^2b^2 & +\ 3ab^3 & +\ b^4 & & & \\
\hline
a^4 & +\ 4a^3b & +\ 6a^2b^2 & +\ 4ab^3 & +\ b^4 & & & \text{4th power.} \\
a & +\ b & & & & & & \\
\hline
a^5 & +\ 4a^4b & +\ 6a^3b^2 & +\ 4a^2b^3 & +\ ab^4 & & & \\
 & +\ a^4b & +\ 4a^3b^2 & +\ 6a^2b^3 & +\ 4ab^4 & +\ b^5 & & \\
\hline
a^5 & +\ 5a^4b & +\ 10a^3b^2 & +\ 10a^2b^3 & +\ 5ab^4 & +\ b^5 & & \textit{Ans.}
\end{array}
$$

---

121. How may a binomial be raised to any power?

122. How does the number of multiplications compare with the exponent of the power? If the exponent is 4, what is the number of multiplications? How many when it is *m*? How many things are considered in the raising of powers? Name them.

NOTE.—**122.** It will be observed that the number of multiplications is always 1 less than the units in the exponent of the power. Thus, if the exponent is 1, no multiplication is necessary. If it is 2, we multiply once; if it is 3, twice; if 4, three times, &c. The powers of polynomials may be expressed by means of an exponent. Thus, to express that $a + b$ is to be raised to the 5th power, we write

$$(a + b)^5;$$

if to the mth power, we write

$$(a + b)^m.$$

2. Find the 5th power of the binomial $a - b$.

$$\begin{array}{llllll l}
a & -\ b & & & & & \text{. . . . . 1st power.} \\
a & -\ b & & & & & \\ \hline
a^2 & -\ ab & & & & & \\
 & -\ ab & +\ b^2 & & & & \\ \hline
a^2 & -\ 2ab & +\ b^2 & & & & \text{. . . . . 2d power.} \\
a & -\ b & & & & & \\ \hline
a^3 & -\ 2a^2b & +\ ab^2 & & & & \\
 & -\ a^2b & +\ 2ab^2 & -\ b^3 & & & \\ \hline
a^3 & -\ 3a^2b & +\ 3ab^2 & -\ b^3 & & & \text{. . . . 3d power.} \\
a & -\ b & & & & & \\ \hline
a^4 & -\ 3a^3b & +\ 3a^2b^2 & -\ ab^3 & & & \\
 & -\ a^3b & +\ 3a^2b^2 & -\ 3ab^3 & +\ b^4 & & \\ \hline
a^4 & -\ 4a^3b & +\ 6a^2b^2 & -\ 4ab^3 & +\ b^4 & & \text{. 4th power.} \\
a & -\ b & & & & & \\ \hline
a^5 & -\ 4a^4b & +\ 6a^3b^2 & -\ 4a^2b^3 & +\ ab^4 & & \\
 & -\ a^4b & +\ 4a^3b^2 & -\ 6a^2b^3 & +\ 4ab^4 & -\ b^5 & \\ \hline
a^5 & -\ 5a^4b & +\ 10a^3b^2 & -\ 10a^2b^3 & +\ 5ab^4 & -\ b^5 & \textit{Ans.}
\end{array}$$

In the same way the higher powers may be obtained. By examining the powers of these binomials, it is plain that four things must be considered:

1st. The number of terms of the power.
2d. The signs of the terms.
3d. The exponents of the letters.
4th. The coefficients of the terms.

Let us see according to what laws these are formed.

Of the Terms.

123. By examining the several multiplications, we shall observe that the first power of a binomial contains two terms; the second power, three terms; the third power, four terms; the fourth power, five; the fifth power, six, &c.; and hence we may conclude:

That the number of terms in any power of a binomial, is greater by one than the exponent of the power.

Of the Signs of the Terms.

124. It is evident that when both terms of the given binomial are plus, *all the terms of the power will be plus.*

If the second term of the binomial is negative, then *all the odd terms, counted from the left, will be positive, and all the even terms negative.*

---

123. How many terms does the first power of a binomial contain? The second? The third? The nth power?

124. If both terms of a binomial are positive, what will be the signs of the terms of the power? If the second term is negative, how are the signs of the terms?

Of the Exponents.

125. The letter which occupies the first place in a binomial, is called the *leading letter*. Thus, a is the leading letter in the binomials $a + b$, and $a - b$.

1st. It is evident that the exponent of the leading letter in the first term, will be the same as the exponent of the power; and that this exponent will diminish by one in each term to the right, until we reach the last term, when it will be 0 (Art. 49).

2d. The exponent of the second letter is 0 in the first term, and increases by one in each term to the right, to the last term, when the exponent is the same as that of the given power.

3d. The sum of the exponents of the two letters, in any term, is equal to the exponent of the given power. This last remark will enable us to verify any result obtained by means of the binomial formula.

Let us now apply these principles in the two following examples, in which the coefficients are omitted:

$$(a + b)^6 \ . \ . \ . \ a^6 + a^5b + a^4b^2 + a^3b^3 + a^2b^4 + ab^5 + b^6,$$
$$(a - b)^6 \ . \ . \ . \ a^6 - a^5b + a^4b^2 - a^3b^3 + a^2b^4 - ab^5 + b^6.$$

As the pupil should be practised in writing the terms with their proper signs, without the coefficients, we will add a few more examples.

---

125. Which is the leading letter of a binomial? What is the exponent of this letter in the first term? How does it change in the terms towards the right? What is the exponent of the second letter in the second term? How does it change in the terms towards the right? What is it in the last term? What is the sum of the exponents in any term equal to?

1. $(a+b)^3$ . . $a^3+a^2b+ab^2 + b^3$.
2. $(a-b)^4$ . . $a^4-a^3b+a^2b^2- ab^3 + b^4$.
3. $(a+b)^5$ . . $a^5+a^4b+a^3b^2+a^2b^3+ab^4 + b^5$.
4. $(a-b)^7$ . . $a^7-a^6b+a^5b^2-a^4b^3+a^3b^4-a^2b^5+ab^6-b^7$.

Of the Coefficients.

126. The coefficient of the first term is 1. The coefficient of the second term is the same as the exponent of the given power. The coefficient of the third term is found by multiplying the coefficient of the second term by the exponent of the leading letter in that term, and dividing the product by 2. And finally:

If the coefficient of any term be multiplied by the exponent of the leading letter in that term, and the product divided by the number which marks the place of the term from the left, the quotient will be the coefficient of the next term.

Thus, to find the coefficients in the example,

$$(a-b)^7 \; . \; . \; . \; a^7-a^6b+a^5b^2-a^4b^3+a^3b^4-a^2b^5+ab^6-b^7,$$

we first place the exponent 7 as a coefficient of the second term. Then, to find the coefficient of the third term, we multiply 7 by 6, the exponent of a, and divide by 2. The quotient, 21, is the coefficient of the third term. To find the coefficient of the fourth, we multiply 21 by 5, and divide the product by 3; this gives 35. To find the coefficient of the fifth term, we multiply 35 by 4, and divide the product by 4; this gives 35. The coefficient of the sixth term, found

---

126. What is the coefficient of the first term? What is the coefficient of the second term? How do you find the coefficient of the third term How do you find the coefficient of any term? What are the coefficients of the first and last terms? How are the coefficients of the exponents of any two terms equally distant from the two extremes?

in the same way, is 21; that of the seventh, 7; and that of the eighth, 1. Collecting these coefficients,

$$(a - b)^7 =$$
$$a^7 - 7a^6b + 21a^5b^2 - 35a^4b^3 + 35a^3b^4 - 21a^2b^5 + 7ab^6 - b^7.$$

NOTE.—We see, in examining this last result, that *the coefficients of the extreme terms are each* 1, *and that the coefficients of terms equally distant from the extreme terms are equal.* It will, therefore, be sufficient to find the coefficients of the first half of the terms, and from these the others may be immediately written.

EXAMPLES

1. Find the fourth power of $a + b$.
Ans. $a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4$.

2. Find the fourth power of $a - b$.
Ans. $a^4 - 4a^3b + 6a^2b^2 - 4ab^3 + b^4$.

3. Find the fifth power of $a + b$.
Ans. $a^5 + 5a^4b + 10a^3b^2 + 10a^2b^3 + 5ab^4 + b^5$.

4. Find the fifth power of $a - b$.
Ans. $a^5 - 5a^4b + 10a^3b^2 - 10a^2b^3 + 5ab^4 - b^5$.

5. Find the sixth power of $a + b$.
$a^6 + 6a^5b + 15a^4b^2 + 20a^3b^3 + 15a^2b^4 + 6ab^5 + b^6$

6. Find the sixth power of $a - b$.
$a^6 - 6a^5b + 15a^4b^2 - 20a^3b^3 + 15a^2b^4 - 6ab^5 + b^6$.

127. When the terms of the binomial have coefficients, we may still write out any power of it by means of the Binomial Formula.

7. Let it be required to find the cube of $2c + 3d$.

$$(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3.$$

Here, $2c$ takes the place of a in the formula, and $3d$ the place of b. Hence, we have,

$$(2c+3d)^3 = (2c)^3+3.(2c)^2.3d+3(2c)(3d)^2+(3d)^3 \quad . \quad (1.)$$

and by performing the indicated operations, we have,

$$(2c + 3d)^3 = 8c^3 + 36c^2d + 54cd^2 + 27d^3.$$

If we examine the second member of Equation (1), we see that each term is made up of three factors: 1st, the numerical factor; 2d, some power of $2c$; and 3d, some power of $3d$. The powers of $2c$ are arranged in descending order towards the right, the last term involving the 0 power of $2c$ or 1; the powers of $3d$ are arranged in ascending order from the first term, where the 0 power enters, to the last term.

The operation of raising a binomial involving coefficients, is most readily effected by writing the three factors of each term in a vertical column, and then performing the multiplications as indicated below.

Find, by this method, the cube of $2c + 3d$.

OPERATION.

$$\begin{array}{lllllllll}
 & 1 & + & 3 & + & 3 & + & 1 & \text{Coefficients.} \\
 & 8c^3 & + & 4c^2 & + & 2c & + & 1 & \text{Powers of } 2c \\
 & 1 & + & 3d & + & 9d^2 & + & 27d^3 & \text{Powers of } 3d \\
\hline
(2c + d)^3 = & 8c^3 & + & 36c^2d & + & 54cd^2 & + & 27d^3 &
\end{array}$$

The preceding operation hardly requires explanation. In the first line, write the numerical coefficients corresponding to the particular power; in the second line, write the descending powers of the leading term to the 0 power; in the third line, write the ascending powers of the following term from the 0 power upwards. It will be easiest to commence the second line on the right hand. The multiplication should be performed from above, downwards.

8. Find the 4th power of $3a^2c - 2bd$.

$$(a + b)^4 = a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4.$$

$$\begin{array}{llllllllll}
1 & + & 4 & + & 6 & + & 4 & + & 1 \\
81a^8c^4 & + & 27a^6c^3 & + & 9a^4c^2 & + & 3a^2c & + & 1 \\
1 & - & 2bd & + & 4b^2d^2 & - & 8b^3d^3 & + & 16b^4d\,. \\
\hline
\end{array}$$

$$81a^8c^4 - 216a^6c^2bd + 216a^4c^2b^2d^2 - 96a^2cb^3d^3 + 16b^4d^4.^*$$

9. What is the cube of $3x - 6y$?

Ans. $27x^3 - 162x^2y + 324xy^2 - 216y^3$.

10. What is the fourth power of $a - 3b$?

Ans. $a^4 - 12a^3b + 54a^2b^2 - 108ab^3 + 81b^4$.

11. What is the fifth power of $c - 2d$?

Ans. $c^5 - 10c^4d + 40c^3d^2 - 80c^2d^3 + 80cd^4 - 32d^5$.

12. What is the cube of $5a - 3d$?

Ans. $125a^3 - 225a^2d + 135ad^2 - 27d^3$.

---

[*] This ingenious method of writing the development of a binomial is due to Professor WILLIAM G. PECK, of Columbia College.

CHAPTER VII.

SQUARE ROOT. RADICALS OF THE SECOND DEGREE.

128. THE SQUARE ROOT of a number is one of its two equal factors. Thus, $6 \times 6 = 36$; therefore, 6 is the square root of 36.

The symbol for the square root, is $\sqrt{\ }$, or the fractional exponent $\frac{1}{2}$; thus,

$$\sqrt{a}, \quad \text{or} \quad a^{\frac{1}{2}},$$

indicates the square root of a, or that one of the two equal factors of a is to be found. The operation of finding such factor is called, *Extracting the Square Root.*

129. Any number which can be resolved into *two equal integral factors*, is called a *perfect square.*

The following Table, verified by actual multiplication, indicates all the perfect squares between 1 and 100.

TABLE.

1,	4,	9,	16,	25,	36,	49,	64,	81,	100,	squares.
1,	2,	3,	4,	5,	6,	7,	8,	9,	10,	roots.

---

128. What is the square root of a number? Wha is the operation of finding the equal factor called?

129. What is a perfect square? How many perfect squares are there between 1 and 100, including both numbers? What are they?

We may employ this table for finding the square root of any perfect square between 1 and 100.

Look for the number in the first line; if it is found there, its square root will be found immediately under it.

If the given number is less than 100, and not a perfect square, *it will fall between two numbers of the upper line, and its square root will be found between the two numbers directly below; the lesser of the two will be the entire part of the root, and will be the true root to within less than* 1.

Thus, if the given number is 55, it is found between the perfect squares 49 and 64, and its root is 7 and a decimal fraction.

NOTE.—There are ten perfect squares between 1 and 100, if we include both numbers; and eight, if we exclude both.

If a number is greater than 100, its square root will be greater than 10, that is, it will contain *tens* and *units*. Let N denote such a number, x the tens of its square root, and y the units; then will,

$$N = (x + y)^2 = x^2 + 2xy + y^2 = x^2 + (2x + y)y.$$

That is, the number is equal to the *square of the tens* in its roots, plus *twice the product of the tens by the units*, plus *the square of the units.*

EXAMPLE.

1. Extract the square root of 6084.

Since this number is composed of more than two places of figures, its root will contain more than one. But since it is less than 10000, which is the square of 100, the root will contain but two figures; that is, units and tens.

$\dot{6}0\ \dot{8}4$

Now, the square of the tens must be found in the two left-hand figures, which we will separate from the other two by putting a point over the place of units, and a second over the place of hundreds. These parts, of two figures each, are called *periods*. The part 60 is comprised between the two squares 49 and 64, of which the roots are 7 and 8; hence, *7 expresses the number of tens sought;* and the required root is composed of 7 tens and a certain number of units.

The figure 7 being found, we write it on the right of the given number, from which we separate it by a vertical line: then we subtract its square, 49, from 60, which leaves a remainder of 11, to which we bring down the two next figures, 84. The result of this operation, 1184, contains *twice the product of the tens by the units, plus the square of the units.*

$$
\begin{array}{r|r|l}
 & 6\dot{0}\ 8\dot{4} & 78 \\
 & 49 & \\ \cline{2-2}
7 \times 2 = 14\ 8 & 118\ 4 & \\
 & 118\ 4 & \\ \cline{2-2}
 & 0 &
\end{array}
$$

But since tens multiplied by units cannot give a product of a less unit than tens, it follows that the last figure, 4, can form no part of the double product of the tens by the units, this double product is therefore found in the part 118, which we separate from the units' place, 4.

Now if we double the tens, which gives 14, and then divide 118 by 14, the quotient 8 *will express the units*, or a number greater than the units. This quotient can never be too small, since the part 118 will be at least equal to twice the product of the tens by the units; but it may be too large, for the 118, besides the double product of the tens by the units, may likewise contain tens arising from the square of the units. To ascertain if the quotient 8 expresses the right number of units, we write the 8 on the right of the 14, which gives 148, and then we multiply 148 by 8. This multiplication being effected, gives for a product, 1184. a number equal to the result of the first operation. Having subtracted the product, we find the remainder equal to 0; hence, 78 is the root required. In this operation, we form, 1st, the square of the tens; 2nd, the double product of the tens by the units; and 3d, the square of the units.

Indeed, in the operations, we have merely subtracted from the given number 6084: 1st, the square of 7 tens, or of 70; 2d, twice the product of 70 by 8; and, 3d, the square of 8; that is, the three parts which enter into the composition of the square, 70 + 8, or 78 and since the result of the subtraction is 0, it follows that 78 is the square root of 6084.

130. The operations in the last example have been performed on but two periods, but it is plain that the same methods of reasoning are equally applicable to larger numbers, for by changing the order of the units, we do not change the relation in which they stand to each other.

Thus, in the number 60 84 95, the two periods 60 84, have the same relation to each other as in the number 60 84; and hence the methods used in the last example are equally applicable to larger numbers.

131. Hence, for the extraction of the square root of numbers, we have the following

RULE.

I. *Point off the given number into periods of two figures each, beginning at the right hand:*

II. *Note the greatest perfect square in the first period on the left, and place its root on the right, after the manner of*

---

131. Give the rule for the extraction of the square root of numbers? What is the first step? What the second? What the third? What the fourth? What the fifth?

a quotient in division; then subtract the square of this root from the first period, and bring down the second period for a remainder:

III. *Double the root already found, and place the result on the left for a divisor. Seek how many times the divisor is contained in the remainder, exclusive of the right-hand figure, and place the figure in the root and also at the right of the divisor:*

IV. *Multiply the divisor, thus augmented, by the last figure of the root, and subtract the product from the remainder, and bring down the next period for a new remainder. But if any of the products should be greater than the remainder, diminish the last figure of the root by one:*

V. *Double the whole root already found, for a new divisor, and continue the operation as before, until all the periods are brought down.*

132. Note.—1. If, after all the periods are brought down, there is no remainder, the given number is a perfect square.

2. The number of places of figures in the root will always be equal to the number of periods into which the given number is divided.

3. If the given number has not an exact root, there will be a remainder after all the periods are brought down, in which case ciphers may be annexed, forming new periods, for each of which there will be one decimal place in the root.

---

132. What takes place when the given number is a perfect square? How many places of figures will there be in the root? If the given number is not a perfect square, what may be done after all the periods are brought down?

EXAMPLES.

1. What is the square root of 36729?

In this example there are two periods of decimals, and, hence, two places of decimals in the root.

$$\begin{array}{r|l|l}
 & \dot{3}\ 6\dot{7}\ 2\dot{9} & 191.64+ \\
 & 1 & \\ \hline
2\ 9 & 267 & \\
 & 261 & \\ \hline
38\ 1 & 629 & \\
 & 381 & \\ \hline
382\ 6 & 24800 & \\
 & 22956 & \\ \hline
3832\ 4 & 184400 & \\
 & 153296 & \\ \hline
 & 31104 \text{ Rem.} &
\end{array}$$

2. To find the square root of 7225. *Ans.* 85.
3. To find the square root of 17689. *Ans.* 133.
4. To find the square root of 994009. *Ans.* 997.
5. To find the square root of 85673536. *Ans.* 9256.
6. To find the square root of 67798756. *Ans.* 8234.
7. To find the square root of 978121. *Ans.* 989.
8. To find the square root of 956484. *Ans.* 978.
9. What is the square root of 36372961? *Ans.* 6031.
10. What is the square root of 22071204? *Ans.* 4698.
11. What is the square root of 106929? *Ans.* 327.
12. What of 12088868379025? *Ans.* 3476905.
13. What of 2268741? *Ans.* 1506.23 +
14. What of 7596796? *Ans.* 2756.22 +
15. What is the square root of 96? *Ans.* 9.79795 +.
16. What is the square root of 153? *Ans.* 12.36931 +.
17. What is the square root of 101. *Ans.* 10.04987 +.

18. What of 285970396644? *Ans.* 534762.

19. What of 41605800625? *Ans.* 203975.

20. What of 48303584206084? *Ans.* 6950078.

EXTRACTION OF THE SQUARE ROOT OF FRACTIONS.

133. Since the square or second power of a fraction is obtained by squaring the numerator and denominator separately, it follows that

The square root of a fraction will be equal to the square root of the numerator divided by the square root of the denominator.

For example, the square root of $\frac{a^2}{b^2}$ is equal to $\frac{a}{b}$: for,

$$\frac{a}{b} \times \frac{a}{b} = \frac{a^2}{b^2}.$$

1. What is the square root of $\frac{1}{4}$? *Ans.* $\frac{1}{2}$.

2. What is the square root of $\frac{9}{16}$? *Ans.* $\frac{3}{4}$.

3. What is the square root of $\frac{64}{81}$? *Ans.* $\frac{8}{9}$.

4. What is the square root of $\frac{256}{361}$? *Ans.* $\frac{16}{19}$.

5. What is the square root of $\frac{16}{64}$? *Ans.* $\frac{1}{2}$.

6. What is the square root of $\frac{4096}{61009}$? *Ans.* $\frac{64}{247}$.

7. What is the square root of $\frac{582169}{956484}$? *Ans.* $\frac{763}{978}$.

---

133. To what is the square root of a fraction equal?

134. If the numerator and denominator are not perfect squares, the root of the fraction cannot be exactly found. We can, however, easily find the approximate root.

RULE.

Multiply both terms of the fraction by the denominator: Then extract the square root of the numerator, and divide this root by the root of the denominator; the quotient will be the approximate root.

1. Find the square root of $\frac{3}{5}$.

Multiplying the numerator and denominator by 5

$$\sqrt{\frac{3}{5}} = \sqrt{\frac{15}{25}} = \frac{\sqrt{15}}{5} = (3.8729 +) \div 5;$$

hence, $(3.8729 +) \div 5 = .7745 +$ = *Ans.*

2. What is the square root of $\frac{7}{4}$?	*Ans.* 1.32287 +.
3. What is the square root of $\frac{14}{9}$?	*Ans.* 1.24721 +.
4. What is the square root of $11\frac{11}{16}$?	*Ans.* 3.41869 +.
5. What is the square root of $7\frac{13}{36}$?	*Ans.* 2.71313 +.
6. What is the square root of $8\frac{15}{49}$?	*Ans.* 2.88203 +.
7. What is the square root of $\frac{5}{12}$?	*Ans.* 0.64549 +.
8. What is the square root of $10\frac{3}{10}$?	*Ans.* 3.20936 +.

---

134. What is the rule when the numerator and denominator are not perfect squares?

135. Finally, instead of the last method, we may, if we please,

Change the common fraction into a decimal, and continue the division until the number of decimal places is double the number of places required in the root. Then extract the root of the decimal by the last rule.

EXAMPLES.

1. Extract the square of $\frac{11}{14}$ to within .001. This number, reduced to decimals, is 0.785714 to within 0.000001; but the root of 0.785714 to the nearest unit, is .886; hence, 0.886 is the root of $\frac{11}{14}$ to within .001.

2. Find the $\sqrt{2\frac{13}{15}}$ to within 0.0001. *Ans.* 1.6931 +.

3. What is the square root of $\frac{1}{17}$? *Ans.* 0.24253 +.

4. What is the square root of $\frac{7}{8}$? *Ans.* 0.93541 +.

5. What is the square root of $\frac{5}{3}$? *Ans.* 1.29099 +.

EXTRACTION OF THE SQUARE ROOT OF MONOMIALS.

136. In order to discover the process for extracting the square root of a monomial, we must see how its square is formed.

By the rule for the multiplication of monomials (Art. 42), we have,

$$(5a^2b^3c)^2 = 5a^2b^3c \times 5a^2b^3c = 25a^4b^6c^2;$$

---

135. What is a second method of finding the approximate root?
136. Give the rule for extracting the square root of monomials?

that is, in order to square a monomial, it is necessary to *square its coefficient and double the exponent of each of the letters.* Hence, to find the square root of a monomial, we have the following

RULE.

I. *Extract the square root of the coefficient for a new coefficient:*

II. *Divide the exponent of each letter by* 2, *and then annex all the letters with their new exponents.*

Since like signs in two factors give a plus sign in the product, the square of $-a$, as well as that of $+a$, will be $+a^2$; hence, the square root of a^2 is either $+a$, or $-a$. Also, the square root of $25a^2b^4$, is either $+5ab^2$, or $-5ab^2$. Whence we conclude, that if a monomial is positive, its square root may be affected either with the sign $+$ or $-$; thus, $\sqrt{9a^4} = \pm 3a^2$; for, $+3a^2$ or $-3a^2$, squared, gives $+9a^4$. The double sign $\pm$, with which the root is affected, is read *plus and minus.*

EXAMPLES.

1. What is the square root of $64a^6b^4$?

$$\sqrt{64a^6b^4} = +8a^3b^2; \text{ for } +8a^3b^2 \times +8a^3b^2 = +64a^6b^4$$

and, $\sqrt{64a^6b^4} = -8a^3b^2$; for $-8a^3b^2 \times -8a^3b^2 = +64a^6b^4$

Hence, $\sqrt{64a^6b^4} = \pm 8a^3b^4$.

2. Find the square root of $625a^2b^8c^6$. $\pm 25ab^4c^3$.

3. Find the square root of $576a^4b^6c^8$. $\pm 24a^2b^3c^4$.

4. Find the square root of $196x^6y^2z^4$. $\pm 14x^3yz^2$.

5. Find the square root of $441a^8b^6c^{10}d^{16}$. $\pm 21a^4b^3c^5d^8$.

6. Find the square root of $784a^{12}b^{14}c^{16}d^2$. $\pm 28a^6b^7c^8d$.

7. Find the square root of $81a^8b^4c^6$. $\pm 9a^4b^2c^3$.

NOTES.—**137.** 1. From the preceding rule it follows, that when a monomial is a perfect square, *its numerical coefficient is a perfect square, and all its exponents even numbers.* Thus, $25a^4b^2$ is a perfect square.

2. If the proposed monomial were *negative*, it would be impossible to extract its square root, since it has just been shown (Art. 136) that the square of every quantity, whether positive or negative, is essentially positive. Therefore,

$$\sqrt{-9}, \quad \sqrt{-4a^2}, \quad \sqrt{-8a^2b},$$

are algebraic symbols which indicate operations that cannot be performed. They are called *imaginary quantities*, or rather, *imaginary expressions*, and are frequently met with in the resolution of equations of the second degree.

IMPERFECT SQUARES.

138. When the coefficient is not a *perfect square*, or when the exponent of any letter is *uneven*, the monomial is an *imperfect square:* thus, $98ab^4$ is an *imperfect square.* Its root is then indicated by means of the radical sign; thus,

$$\sqrt{98ab^4}.$$

Such quantities are called, *radical quantities*, or *radicals of the second degree:* hence,

A RADICAL QUANTITY, is the indicated root of an imperfect power.

---

137. When is a monomial a perfect square? What monomials are these whose square roots cannot be extracted? What are such expressions called?

138. When is a monomial an imperfect square? What are such quantities called? What is a radical quantity?

TRANSFORMATION OF RADICALS.

139. Let a and b denote any two numbers, and p the product of their square roots: then,

$$\sqrt{a} \times \sqrt{b} = p \quad . \quad . \quad . \quad . \quad . \quad (1.)$$

Squaring both members, we have,

$$a \times b = p^2 \quad . \quad . \quad . \quad . \quad (2.)$$

Then, extracting the square root of both members of (2),

$$\sqrt{ab} = p \quad . \quad . \quad . \quad . \quad . \quad (3.)$$

And since the second members are the same in Equations (1) and (3), the first members are equal: that is,

The square root of the product of two quantities is equal to the product of their square roots.

140. Let a and b denote any two numbers, and q the quotient of their square roots; then,

$$\frac{\sqrt{a}}{\sqrt{b}} = q \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad (1.)$$

Squaring both members, we have,

$$\frac{a}{b} = q^2 \quad . \quad . \quad . \quad . \quad . \quad . \quad (2.)$$

then extracting the square root of both members of (2),

$$\sqrt{\frac{a}{b}} = q \quad . \quad . \quad . \quad . \quad . \quad . \quad (3)$$

and since the second members are the same in Equations (1) and (3), the first members are equal; that is,

---

139. To what is the square root of the product of two quantities equal?

140. To what is the square root of the quotient of two quantities equal?

The square root of the quotient of two quantities is equal to the quotient of their square roots.

These principles enable us to *transform* radical expressions, or to *reduce* them to simpler forms; thus, the expression,

$$98ab^4 = 49b^4 \times 2a;$$

hence, $\sqrt{98ab^4} = \sqrt{49b^4 \times 2a};$

and by the principle of (Art. 139),

$$\sqrt{49b^4 \times 2a} = \sqrt{49b^4} \times \sqrt{2a} = 7b^2\sqrt{2a}.$$

In like manner,

$$\sqrt{45a^2b^3c^2d} = \sqrt{9a^2b^2c^2 \times 5bd} = 3abc\sqrt{5bd}.$$

$$\sqrt{864a^2b^5c^{11}} = \sqrt{144a^2b^4c^{10} \times 6bc} = 12ab^2c^5\sqrt{6bc}.$$

The COEFFICIENT of a radical is the quantity without the sign; thus, in the expressions,

$$7b^2\sqrt{2a}, \quad 3abc\sqrt{5bd}, \quad 12ab^2c^5\sqrt{6bc},$$

the quantities $7b^2$, $3abc$, $12ab^2c^5$, are *coefficients of the radicals.*

141. Hence, to simplify a radical of the second degree, we have the following

RULE.

I. *Divide the expression under the radical sign into two factors, one of which shall be a perfect square:*

II. *Extract the square root of the perfect square, and then multiply this root by the indicated square root of the remaining factor.*

---

141. Give the rule for simplifying radicals of the second degree. How do you determine whether a given number has a factor which is a perfect square?

NOTE.—To determine if a given number has any factor which is a perfect square, we examine and see if it is divisible by either of the perfect squares,

4, 9, 16, 25, 36, 49, 64, 81, &c.;

if it is not, we conclude that it does not contain a factor which is a perfect square.

EXAMPLES.

Reduce the following radicals to their simplest form:

1. $\sqrt{75a^3bc}$. *Ans.* $5a\sqrt{3abc}$.
2. $\sqrt{128b^5a^6d^2}$. *Ans.* $8b^2a^3d\sqrt{2b}$.
3. $\sqrt{32a^9b^8c}$. *Ans.* $4a^4b^4\sqrt{2ac}$.
4. $\sqrt{256a^2b^4c^8}$. *Ans.* $16ab^2c^4$.
5. $\sqrt{1024a^9b^7c^5}$. *Ans.* $32a^4b^3c^2\sqrt{abc}$.
6. $\sqrt{729a^7b^5c^6d}$. *Ans.* $27a^3b^2c^3\sqrt{abd}$.
7. $\sqrt{675a^7b^5c^2d}$. *Ans.* $15a^3b^2c\sqrt{3abd}$.
8. $\sqrt{1445a^3c^8d^4}$. *Ans.* $17ac^4d^2\sqrt{5a}$.
9. $\sqrt{1008a^9d^7m^8}$. *Ans.* $12a^4d^3m^4\sqrt{7ad}$.
10. $\sqrt{2156a^{10}b^8c^6}$. *Ans.* $14a^5b^4c^3\sqrt{11}$.
11. $\sqrt{405a^7b^6d^8}$. *Ans.* $9a^3b^3d^4\sqrt{5a}$.

142. NOTES.—1. A *coefficient*, or a *factor of a coefficient*, may be carried under the radical sign, *by squaring it.* Thus,

1. $3a^2\sqrt{bc} = \sqrt{(3a^2)^2 \times bc} = \sqrt{9a^4bc}$.
2. $2ab\sqrt{d} = 2\sqrt{a^2b^2d} = \sqrt{4a^2b^2d}$.

---

142. How may a coefficient or factor be carried under the radical sign To what is the square root of a negative quantity equal?

3. $4(a+b)\sqrt{a-b}=4\sqrt{(a+b)^2(a-b)}=4\sqrt{(a^2-b^2)(a+b)}$

4. $5bc\sqrt{a^2-c^2} = 5\sqrt{b^2c^2(a^2-c^2)}$.

2. The square root of a negative quantity may also be simplified; thus,

$$\sqrt{-9} = \sqrt{9 \times -1} = \sqrt{9} \times \sqrt{-1} = 3\sqrt{-1},$$

and, $\sqrt{-4a^2} = \sqrt{4a^2} \times \sqrt{-1} = 2a\sqrt{-1}$; also

$$\sqrt{-8a^2b} = \sqrt{4a^2 \times -2b} = 2a\sqrt{-2b} = 2a\sqrt{2b} \times \sqrt{-1};$$

that is, *the square root of a negative quantity is equal to the square root of the same quantity with a positive sign, multiplied into the square root of* -1.

Reduce the following:

1. $\sqrt{-64a^2b^2}$. *Ans.* $8ab\sqrt{-1}$.
2. $\sqrt{-128a^4b^5}$. *Ans.* $8a^2b^2\sqrt{2b}\sqrt{-1}$.
3. $\sqrt{-72a^5b^7c^6}$. *Ans.* $6a^2b^3c^3\sqrt{2ab}\sqrt{-1}$.
4. $\sqrt{-48a^3bc^5}$. *Ans.* $4ac^2\sqrt{3abc}\sqrt{-1}$.

ADDITION OF RADICALS.

143. SIMILAR RADICALS, of the second degree, are those in which the quantities under the sign are the same. Thus, the radicals $3\sqrt{b}$, and $5c\sqrt{b}$ are similar, and so also are $9\sqrt{2}$, and $7\sqrt{2}$.

144. Radicals are added like other algebraic quantities hence, the following

---

143. What are similar radicals of the second degree?

144. Give the rule for the addition of radicals of the second degree?

RULE.

I. *If the radicals are similar, add their coefficients, and to the sum annex the common radical:*

II. *If the radicals are not similar, connect them together with their proper signs.*

Thus, $3a\sqrt{b} + 5c\sqrt{b} = (3a + 5c)\sqrt{b}$.

In like manner,

$$7\sqrt{2a} + 3\sqrt{2a} = (7 + 3)\sqrt{2a} = 10\sqrt{2a}.$$

NOTES.—1. Two radicals, which do not appear to be similar at first sight, may become so by transformation (Art. 141.)

For example,

$$\sqrt{48ab^2} + b\sqrt{75a} = 4b\sqrt{3a} + 5b\sqrt{3a} = 9b\sqrt{3a};$$

$$2\sqrt{45} + 3\sqrt{5} = 6\sqrt{5} + 3\sqrt{5} = 9\sqrt{5}.$$

2. When the radicals are not similar, the addition or subtraction can only be indicated. Thus, in order to add $3\sqrt{b}$ to $5\sqrt{a}$, we write,

$$5\sqrt{a} + 3\sqrt{b}.$$

Add together the following:

1. $\sqrt{27a^2}$ and $\sqrt{48a^2}$. *Ans.* $7a\sqrt{3}$.

2. $\sqrt{50a^4b^2}$ and $\sqrt{72a^4b^2}$. *Ans.* $11a^2b\sqrt{2}$.

3. $\sqrt{\frac{3a^2}{5}}$ and $\sqrt{\frac{a^2}{15}}$. *Ans.* $4a\sqrt{\frac{1}{15}}$.

4. $\sqrt{125}$ and $\sqrt{500a^2}$. *Ans.* $(5 + 10a)\sqrt{5}$.

5. $\sqrt{\frac{50}{147}}$ and $\sqrt{\frac{100}{294}}$ *Ans.* $\frac{10}{21}\sqrt{6}$.

6. $\sqrt{98a^2x}$ and $\sqrt{36x^2 - 36a^2}$.
Ans. $7a\sqrt{2x} + 6\sqrt{x^2 - a^2}$.

7. $\sqrt{98a^2x}$ and $\sqrt{288a^4x^5}$. *Ans.* $(7a + 12a^2x^2)\sqrt{2x}$.

8. $\sqrt{72}$ and $\sqrt{128}$. *Ans.* $14\sqrt{2}$.

9. $\sqrt{27}$ and $\sqrt{147}$. *Ans.* $10\sqrt{3}$.

10. $\sqrt{\frac{2}{3}}$ and $\sqrt{\frac{27}{50}}$. *Ans.* $\frac{19}{30}\sqrt{6}$.

11. $2\sqrt{a^2b}$ and $3\sqrt{64bx^4}$. *Ans.* $(2a + 24x^2)\sqrt{b}$.

12. $\sqrt{243}$ and $10\sqrt{363}$. *Ans.* $119\sqrt{3}$.

13. $\sqrt{320a^2b^2}$ and $\sqrt{245a^8b^6}$. *Ans.* $(8ab + 7a^4b^3)\sqrt{5}$.

14. $\sqrt{75a^6b^7}$ and $\sqrt{300a^6b^5}$. *Ans.* $(5a^3b^3 + 10a^3b^2)\sqrt{3b}$.

SUBTRACTION OF RADICALS.

145. Radicals are subtracted like other algebraic quantities; hence, the following

RULE.

I. *If the radicals are similar, subtract the coefficient of the subtrahend from that of the minuend, and to the difference annex the common radical:*

II. *If the radicals are not similar, indicate the operation by the minus sign.*

EXAMPLES.

1. What is the difference between $3a\sqrt{b}$ and $a\sqrt{b}$?

Here, $3a\sqrt{b} - a\sqrt{b} = 2a\sqrt{b}$. *Ans.*

---

145. Give the rule for the subtraction of radicals.

2. From $9a\sqrt{27b^2}$ subtract $6a\sqrt{27b^2}$.

First, $9a\sqrt{27b^2} = 27ab\sqrt{3}$, and $6a\sqrt{27b^2} = 13ab\sqrt{3}$;

and, $27ab\sqrt{3} - 18ab\sqrt{3} = 9ab\sqrt{3}$. *Ans.*

Find the differences between the following:

3. $\sqrt{75}$ and $\sqrt{48}$. *Ans.* $\sqrt{3}$.

4. $\sqrt{24a^2b^2}$ and $\sqrt{54b^4}$. *Ans.* $(2ab - 3b^2)\sqrt{6}$.

5. $\sqrt{\frac{3}{5}}$ and $\sqrt{\frac{5}{27}}$. *Ans.* $\frac{4}{45}\sqrt{15}$.

6. $\sqrt{128a^3b^2}$ and $\sqrt{32a^9}$. *Ans.* $(8ab - 4a^4)\sqrt{2a}$.

7. $\sqrt{48a^3b^3}$ and $\sqrt{9ab}$. *Ans.* $4ab\sqrt{3ab} - 3\sqrt{ab}$.

8. $\sqrt{242a^5b^5}$ and $\sqrt{2a^3b^3}$. *Ans.* $(11a^2b^2 - ab)\sqrt{2ab}$.

9. $\sqrt{\frac{3}{4}}$ and $\sqrt{\frac{3}{9}}$. *Ans.* $\frac{1}{6}\sqrt{3}$.

10. $\sqrt{320a^2}$ and $\sqrt{80a^2}$. *Ans.* $4a\sqrt{5}$.

11. $\sqrt{720a^3b^3}$ and $\sqrt{245abc^2d^2}$.
Ans. $(12ab - 7cd)\sqrt{5ab}$.

12. $\sqrt{968a^2b^2}$ and $\sqrt{200a^2b^2}$. *Ans.* $12ab\sqrt{2}$.

13. $\sqrt{112a^8b^6}$ and $\sqrt{28a^8b^6}$. *Ans.* $2a^4b^3\sqrt{7}$.

MULTIPLICATION OF RADICALS.

146. Radicals are multiplied like other algebraic quantities; hence, we have the following

RULE.

I. *Multiply the coefficients together for a new coefficient:*

---

146. Give the rule for the multiplication of radicals.

II. *Multiply together the quantities under the radical signs:*

III. *Then reduce the result to its simplest form.*

1. Multiply $3a\sqrt{bc}$ by $2\sqrt{ab}$.

$$3a\sqrt{bc} \times 2\sqrt{ab} = 3a \times 2 \times \sqrt{bc} \times \sqrt{ab}.$$

which, by Art. 139, $= 6a\sqrt{b^2ac} = 6ab\sqrt{ac}$.

Multiply the following:

2. $3\sqrt{5ab}$ and $4\sqrt{20a}$. *Ans.* $120a\sqrt{b}$,

3. $2a\sqrt{bc}$ and $3a\sqrt{bc}$. *Ans.* $6a^2bc$.

4. $2a\sqrt{a^2+b^2}$ and $-3a\sqrt{a^2+b^2}$. *A.* $-6a^2(a^2+b^2.)$

5. $2ab\sqrt{a+b}$ and $ac\sqrt{a-b}$. *Ans.* $2a^2bc\sqrt{a^2-b^2}$.

6. $3\sqrt{2}$ and $2\sqrt{8}$. *Ans.* 24.

7. $\frac{5}{3}\sqrt{\frac{3}{8}a^2b}$ and $\frac{2}{10}\sqrt{\frac{2}{5}c^2b}$. *Ans.* $\frac{1}{30}abc\sqrt{15}$.

8. $2x+\sqrt{b}$ and $2x-\sqrt{b}$. *Ans.* $4x^2-b$

9. $\sqrt{a+2\sqrt{b}}$ and $\sqrt{a-2\sqrt{b}}$. *Ans.* $\sqrt{a^2-4b}$.

10. $3a\sqrt{27a^3}$ by $\sqrt{2a}$. *Ans.* $9a^3\sqrt{6}$

DIVISION OF RADICALS.

147. Radical quantities are divided like other algebraic quantities; hence, we have the following

RULE.

I. *Divide the coefficient of the dividend by the coefficient of the divisor, for a new coefficient:*

---

147. Give the rule for the division of radicals.

II. *Divide the quantities under the radicals, in the same manner:*

III. *Then reduce the result to its simplest form.*

EXAMPLES.

1. Divide $8a\sqrt{b^3c}$ by $4a\sqrt{bc^3}$.

$$\frac{8a}{4a} = 2, \quad \text{new coefficient.}$$

Art. 140, $\dfrac{\sqrt{b^3c}}{\sqrt{bc^3}} = \sqrt{\dfrac{b^3c}{bc^3}} = \sqrt{\dfrac{b^2}{c^2}} = \dfrac{b}{c}$;

hence, the quotient is $2 \times \dfrac{b}{c} = \dfrac{2b}{c}$.

2. Divide $5a\sqrt{b}$ by $2b\sqrt{c}$. *Ans.* $\dfrac{5a}{2b}\sqrt{\dfrac{b}{c}}$.

3. Divide $12ac\sqrt{6bc}$ by $4c\sqrt{2b}$. *Ans.* $3a\sqrt{3c}$.

4. Divide $6a\sqrt{96b^4}$ by $3\sqrt{8b^2}$. *Ans.* $4ab\sqrt{3}$.

5. Divide $4a^2\sqrt{50b^5}$ by $2a^2\sqrt{5b}$. *Ans.* $2b^2\sqrt{10}$.

6. Divide $26a^3b\sqrt{81a^2b^2}$ by $13a\sqrt{9ab}$. *A.* $6a^2b\sqrt{ab}$.

7. Divide $84a^3b^4\sqrt{27ac}$ by $42ab\sqrt{3a}$. *A.* $6a^2b^3\sqrt{c}$.

8. Divide $\sqrt{\frac{1}{8}a^2}$ by $\sqrt{2}$. *Ans.* $\frac{1}{4}a$.

9. Divide $6a^2b^2\sqrt{20a^3}$ by $12\sqrt{5a}$. *Ans.* a^3b^2.

10. Divide $6a\sqrt{10b^2}$ by $3\sqrt{5}$. *Ans.* $2ab\sqrt{2}$.

11. Divide $48b^4\sqrt{15}$ by $2b^2\sqrt{\frac{1}{15}}$ *Ans.* $360b^2$.

12. Divide $8a^2b^4c^3\sqrt{7d^3}$ by $2a\sqrt{28d}$. *Ans.* $2ab^4c^3d$.

13. Divide $96a^4c^3\sqrt{98b^5}$ by $48abc\sqrt{2b}$. *A.* $14a^3bc^2$.

14. Divide $27a^5b^6\sqrt{21a^3}$ by $\sqrt{7a}$. *Ans.* $27a^6b^6\sqrt{3}$.

15. Divide $18a^8b^6\sqrt{8a^4}$ by $6ab\sqrt{a^2}$. *Ans.* $6a^8b^5\sqrt{2}$.

SQUARE ROOT OF POLYNOMIALS.

148. Before explaining the rule for the extraction of the square root of a polynomial, let us first examine the squares of several polynomials: we have,

$$(a + b)^2 = a^2 + 2ab + b^2,$$
$$(a + b + c)^2 = a^2 + 2ab + b^2 + 2(a + b)c + c^2,$$
$$(a + b + c + d)^2 = a^2 + 2ab + b^2 + 2(a + b)c + c^2 + 2(a + b + c)d + d^2.$$

The *law* by which these squares are formed can be enunciated thus:

The square of any polynomial is equal to the square of the first term, plus twice the product of the first term by the second, plus the square of the second; plus twice the first two terms multiplied by the third, plus the square of the third; plus twice the first three terms multiplied by the fourth, plus the square of the fourth; and so on.

149. Hence, to extract the square root of a polynomial, we have the following

RULE.

I. *Arrange the polynomial with reference to one of its letters, and extract the square root of the first term: this will give the first term of the root:*

---

148. What is the square of a binomial equal to? What is the square of a trinomial equal to? To what is the square of any polynomial equal?

149. Give the rule for extracting the square root of a polynomial? What is the first step? What the second? What the third? What the fourth?

II. *Divide the second term of the polynomial by double the first term of the root, and the quotient will be the second term of the root:*

III. *Then form the square of the algebraic sum of the two terms of the root found, and subtract it from the first polynomial, and then divide the first term of the remainder by double the first term of the root, and the quotient will be the third term:*

IV. *Form the double product of the sum of the first and second terms by the third, and add the square of the third; then subtract this result from the last remainder, and divide the first term of the result so obtained, by double the first term of the root, and the quotient will be the fourth term. Then proceed in a similar manner to find the other terms.*

EXAMPLES.

1. Extract the square root of the polynomial,

$$49a^2b^2 - 24ab^3 + 25a^4 - 30a^3b + 16b^4.$$

First arrange it with reference to the letter a.

$$\begin{array}{r|l}
25a^4 - 30a^3b + 49a^2b^2 - 24ab^3 + 16b^4 & \underline{5a^2 - 3ab + 4b^2} \\
\underline{25a^4 - 30a^3b + 9a^2b^2 \qquad\qquad\qquad\quad} & 10a^2 \\
40a^2b^2 - 24ab^3 + 16b^4 & \quad . \quad . \quad \textit{1st Rem.} \\
\underline{40a^2b^2 - 24ab^3 + 16b^4} & \\
0 \; . \; . \; . \; . \; . & \quad . \quad . \quad \textit{2d Rem.}
\end{array}$$

After having arranged the polynomial with reference to a, extract the square root of $25a^4$; this gives $5a^2$, which is placed at the right of the polynomial: then divide the second term, $-30a^3b$, by the double of $5a^2$, or $10a^2$; the quotient is $-3ab$, which is placed at the right of $5a^2$. Hence, the first two terms of the root are $5a^2 - 3ab$. Squaring this binomial, it becomes $25a^4 - 30a^3b + 9a^2b^2$, which, subtracted from the proposed polynomial, gives a remainder, of which the first term is $40a^2b^2$. Dividing this first term by $10a^2$, (the double of $5a^2$), the quotient is $+ 4b^2$; this is the third term of the root, and is written on the right of the first two terms. By forming the double product of $5a^2 - 3ab$ by $4b^2$, squaring $4b^2$, and taking the sum, we find the polynomial $40a^2b^2 - 24ab^3 + 16b^4$, which, subtracted from the first remainder, gives 0. Therefore, $5a^2 - 3ab + 4b^2$ is the required root.

2. Find the square root of $a^4 + 4a^3x + 6a^2x^2 + 4ax^3 + x^4$.
Ans. $a^2 + 2ax + x^2$.

3. Find the square root of $a^4 - 4a^3x + 6a^2x^2 - 4ax^3 + x^4$.
Ans. $a^2 - 2ax + x^2$.

4. Find the square root of
$4x^6 + 12x^5 + 5x^4 - 2x^3 + 7x^2 - 2x + 1$.
Ans. $2x^3 + 3x^2 - x + 1$

5. Find the square root of
$9a^4 - 12a^3b + 28a^2b^2 - 16ab^3 + 16b^4$.
Ans. $3a^2 - 2ab + 4b^2$

6. What is the square root of
$x^4 - 4ax^3 + 4a^2x^2 - 4x^2 + 8ax + 4$?
Ans. $x^2 - 2ax - 2$.

7. What is the square root of
$9x^2 - 12x + 6xy + y^2 - 4y + 4$?
Ans. $3x + y - 2$.

8. What is the square root of $y^4 - 2y^2x^2 + 2x^2 - 2y^2 + 1 + x^4$?
Ans. $y^2 - x^2 - 1$.

9. What is the square root of $9a^4b^4 - 30a^3b^3 + 25a^2b^2$?
Ans. $3a^2b^2 - 5ab$.

10. Find the square root of
$25a^4b^2 - 40a^3b^2c + 76a^2b^2c^2 - 48ab^2c^3 + 36b^2c^4 - 30a^4bc$
$+ 24a^3bc^2 - 36a^2bc^3 + 9a^4c^2$.
Ans. $5a^2b - 3a^2c - 4abc + 6bc^2$.

150. We will conclude this subject with the following remarks:

1st. A binomial can never be a perfect square, since we know that the square of the most simple polynomial, viz., a binomial, contains three distinct parts, which cannot experience any reduction amongst themselves. Thus, the expression $a^2 + b^2$, is not a perfect square; it wants the term $\pm 2ab$, in order that it should be the square of $a \pm b$.

2d. In order that a trinomial, when arranged, may be a perfect square, its two extreme terms must be squares, and the middle term must be the double product of the square roots of the two others. Therefore, to obtain the square root of a trinomial when it is a perfect square: *Extract the roots of the two extreme terms, and give these roots the same or contrary signs, according as the middle term is positive or negative. To verify it, see if the double product of the two roots is the same as the middle term of the trinomial.* Thus,

$$9a^6 - 48a^4b^2 + 64a^2b^4, \text{ is a perfect square,}$$

since, $\sqrt{9a^6} = 3a^3$, and $\sqrt{64a^2b^4} = -8ab^2$;

and also,

$$2 \times 3a^3 \times -8ab^2 = -48a^4b^2 = \text{the middle term.}$$

But, $4a^2 + 14ab + 9b^2$ is not a perfect square: for, although $4a^2$ and $+9b^2$ are the squares of $2a$ and $3b$, yet $2 \times 2a \times 3b$ is not equal to $14ab$.

3d. In the series of operations required by the general rule, when the first term of one of the remainders is not exactly divisible by twice the first term of the root, we may

---

150. Can a binomial ever be a perfect power? Why not? When is a trinomial a perfect square? When, in extracting the square root, we find that the first term of the remainder is not divisible by twice the root, is the polynomial a perfect power or not?

conclude that the proposed polynomial is not a perfect square. This is an evident consequence of the course of reasoning by which we have arrived at the general rule for extracting the square root.

4th. When the polynomial is *not a perfect square*, it may sometimes be simplified (See Art. 139).

Take, for example, the expression, $\sqrt{a^3b + 4a^2b^2 + 4ab^3}$.

The quantity under the radical is not a perfect square; but it can be put under the form $ab(a^2 + 4ab + 4b^2.)$ Now, the factor within the parenthesis is evidently the square of $a + 2b$, whence, we may conclude that,

$$\sqrt{a^3b + 4a^2b^2 + 4ab^3} = (a + 2b)\sqrt{ab}.$$

2. Reduce $\sqrt{2a^2b - 4ab^2 + 2b^3}$ to its simplest form.

Ans. $(a - b)\sqrt{2b}$.

CHAPTER VIII.

EQUATIONS OF THE SECOND DEGREE.

EQUATIONS CONTAINING ONE UNKNOWN QUANTITY.

151. An Equation of the second degree containing but one unknown quantity, is one in which the greatest exponent is equal to 2. Thus,

$$x^2 = a, \quad ax^2 + bx = c,$$

are equations of the second degree.

152. Let us see to what form every equation of the second degree may be reduced.

Take any equation of the second degree, as,

$$(1 + x)^2 - \frac{3}{4}x - 10 = 5 - \frac{x}{4} + \frac{x^2}{2}.$$

Clearing of fractions, and performing indicated operations, we have,

$$4 + 8x + 4x^2 - 3x - 40 = 20 - x + 2x^2.$$

Transposing the unknown terms to the first member, the known terms to the second, and arranging with reference to the powers of x, we have,

$$4x^2 - 2x^2 + 8x - 3x + x = 20 + 40 - 4;$$

---

151. What is an equation of the second degree? Give an example.

152. To what form may every equation of the second degree be reduced?

and, by reducing,

$$2x^2 + 6x = 56;$$

dividing by the coefficient of x^2, we have,

$$x^2 + 3x = 28$$

If we denote the coefficient of x by $2p$, and the second member by q, we have,

$$x^2 + 2px = q.$$

This is called the *reduced equation.*

153. When the reduced equation is of this form, it contains three terms, and is called a *complete equation.* The terms are,

FIRST TERM.—The second power of the unknown quantity, with a plus sign.

SECOND TERM.—The first power of the unknown quantity, with a coefficient.

THIRD TERM.—A known term, in the second member.

Every equation of the second degree may be reduced to this form, by the following

RULE.

I. *Clear the equation of fractions, and perform all the indicated operations:*

II. *Transpose all the unknown terms to the first member, and all the known terms to the second member:*

---

153. How many terms are there in a complete equation? What is the first term? What is the second term? What is the third term? How many operations are there in reducing an equation of the second degree to the required form? What is the first? What the second? What the third? What the fourth?

III. *Reduce all the terms containing the square of the unknown quantity to a single term, one factor of which is the square of the unknown quantity; reduce, also, all the terms containing the first power of the unknown quantity, to a single term:*

IV. *Divide both members of the resulting equation by the coefficient of the square of the unknown quantity.*

154. A Root of an equation is such a value of the unknown quantity as, being substituted for it, will satisfy the equation; that is, make the two members equal.

The Solution of an equation is the operation of finding its roots.

INCOMPLETE EQUATIONS.

155. It may happen, that $2p$, the coefficient of the first power of x, in the equation $x + 2px = q$, is equal to 0. In this case, the first power of x will disappear, and the equation will take the form,

$$x^2 = q \quad . \; . \; . \; . \; . \; . \; . \quad (1.)$$

This is called an incomplete equation; hence,

An incomplete equation, when reduced, contains but two terms; the square of the unknown quantity, and a known term.

156. Extracting the square root of both members of Equation (1), we have,

$$x = \pm\sqrt{q}.$$

---

154. What is the root of an equation? What is the solution of an equation?

155. What form will the reduced equation take when the coefficient ot x is 0? What is the equation then called? How many terms are there in an incomplete equation? What are they?

156. What is the rule for the solution of an incomplete equation? How many roots are there in every incomplete equation? How do the roots compare with each other?

Hence, for the solution of incomplete equations:

RULE.

I. *Reduce the equation to the form* $x^2 = q$*:*

II. *Then extract the square root of both members.*

NOTE.—There will be two roots, numerically equal, but having contrary signs. Denoting the first by x', and the second by x'', we have,

$$x' = +\sqrt{q}, \text{ and } x'' = -\sqrt{q}.$$

VERIFICATION.

Substituting $+\sqrt{q}$, or $-\sqrt{q}$, for x, in Equation (1), we have,

$$(+\sqrt{q})^2 = q; \text{ and, } (-\sqrt{q})^2 = q;$$

hence, both satisfy the equation; they are, therefore, roots. (Art. 154.)

EXAMPLES.

1. What are the values of x in the equation,

$$3x^2 + 8 = 5x^2 - 10?$$

By transposing, $3x^2 - 5x^2 = -10 - 8$.

Reducing, $-2x^2 = -18$.

Dividing by -2, $x^2 = 9$.

Extracting square root, $x = \pm\sqrt{9} = +3$ and -3.

Hence, $x' = +3$, and $x'' = -3$.

2. What are the roots of the equation,

$$3x^2 + 6 = 4x^2 - 10?$$

Ans. $x' = +4$, $x'' = -4$.

3. What are the roots of the equation,

$$\frac{1}{3}x^2 - 8 = \frac{x^2}{9} + 10\,?$$

Ans. $x' = +9, \quad x'' = -9.$

4. What are the roots of the equation,

$$4x^2 + 13 - 2x^2 = 45\,?$$

Ans. $x' = +4, \quad x'' = -4.$

5. What are the roots of the equation,

$$6x^2 - 7 = 3x^2 + 5\,?$$

Ans. $x' = +2, \quad x'' = -2.$

6. What are the roots of the equation,

$$8 + 5x^2 = \frac{x^2}{5} + 4x^2 + 28\,?$$

Ans. $x' = +5, \quad x'' = -5.$

7. What are the roots of the equation,

$$\frac{3x^2 + 5}{8} - \frac{x^2 + 29}{3} = 117 - 5x^2\,?$$

Ans. $x' = +5, \quad x'' = -5.$

8. What are the roots of the equation,

$$x^2 + ab = 5x^2\,?$$

Ans. $x' = +\frac{1}{2}\sqrt{ab}, \quad x'' = -\frac{1}{2}\sqrt{ab}.$

9 What are the roots of the equation,

$$x\sqrt{a + x^2} = b + x^2\,?$$

Ans. $x' = \dfrac{b}{\sqrt{a - 2b}}, \quad x'' = -\dfrac{b}{\sqrt{a - 2b}}$

PROBLEMS.

1. What number is that which being multiplied by itself the product will be 144?

Let $x =$ the number: then,

$$x \times x = x^2 = 144.$$

It is plain that the value of x will be found by extracting the square root of both members of the equation: that is,

$$\sqrt{x^2} = \sqrt{144}: \text{ that is, } x = 12.$$

2. A person being asked how much money he had, said, if the number of dollars be squared and 6 be added, the sum will be 42: how much had he?

Let $x =$ the number of dollars.

Then, by the conditions,

$$x^2 + 6 = 42;$$

hence, $x^2 = 42 - 6 = 36,$

and, $x = 6.$ *Ans.* \$6.

3. A grocer being asked how much sugar he had sold to a person, answered, if the square of the number of pounds be multiplied by 7, the product will be 1575. How many pounds had he sold?

Denote the number of pounds by x. Then, by the conditions of the question,

$$7x^2 = 1575;$$

hence, $x^2 = 225,$

and, $x = 15.$ *Ans.* 15.

4. A person being asked his age, said, if from the square

of my age in years, you take 192 years, the remainder will be the square of half my age: what was his age?

Denote the number of years in his age by x.

Then, by the conditions of the question,

$$x^2 - 192 = \left(\frac{1}{2}x\right)^2 = \frac{x^2}{4},$$

and by clearing the fractions,

$$4x^2 - 768 = x^2;$$

hence, $$4x^2 - x^2 = 768,$$

and, $$3x^2 = 768,$$

$$x^2 = 256$$

$$x = 16 \qquad \textit{Ans.}\ 16 \text{ years.}$$

5. What number is that whose eighth part multiplied by its fifth part and the product divided by 4, will give a quotient equal to 40?

Let $x =$ the number.

By the conditions of the question,

$$\left(\frac{1}{8}x \times \frac{1}{5}x\right) \div 4 = 40;$$

hence, $$\frac{x^2}{160} = 40;$$

by clearing of fractions,

$$x^2 = 6400,$$

$$x = 80. \qquad \textit{Ans.}\ 80.$$

6. Find a number such that one-third of it multiplied by one fourth shall be equal to 108. *Ans.* 36.

7. What number is that whose sixth part multiplied by its fifth part and the product divided by ten, will give a quotient equal to 3? *Ans.* 30.

8. What number is that whose square, plus 18, will be equal to half the square, plus $30\frac{1}{2}$? *Ans.* 5.

9. What numbers are those which are to each other as 1 to 2, and the difference of whose squares is equal to 75?

Let $x =$ the less number.

Then, $2x =$ the greater.

Then, by the conditions of the question,

$$4x^2 - x^2 = 75;$$

hence, $$3x^2 = 75,$$

and by dividing by 3, $x^2 = 25$, and $x = 5$,

and, $$2x = 10.$$

Ans. 5 and 10

10. What two numbers are those which are to each other as 5 to 6, and the difference of whose squares is 44?

Let $x =$ the greater number.

Then, $\frac{5}{6}x =$ the less.

By the conditions of the problem,

$$x^2 - \frac{25}{36}x^2 = 44;$$

by clearing of fractions,

$$36x^2 - 25x^2 = 1584;$$

hence, $$11x^2 = 1584,$$

and, $$x^2 = 144;$$

hence, $$x = 12,$$

and, $$\frac{5}{6}x = 10.$$

Ans. 10 and 12.

11. What two numbers are those which are to each other as 3 to 4, and the difference of whose squares is 28?

Ans. 6 and 8.

12. What two numbers are those which are to each other as 5 to 11, and the sum of whose squares is 584?

Ans. 10 and 22.

13. *A* says to *B*, my son's age is one quarter of yours, and the difference between the squares of the numbers representing their ages is 240: what were their ages?

Ans. { Eldest, 16,
Younger, 4.

Two unknown quantities.

157. When there are two or more unknown quantities:

I. *Eliminate one of the unknown quantities by* Art. 113*:*

II. *Then extract the square root of both members of the equation.*

PROBLEMS.

1. There is a room of such dimensions, that the difference of the sides multiplied by the less, is equal to 36, and the product of the sides is equal to 360: what are the sides?

Let $x =$ the length of the less side;

$y =$ the length of the greater.

Then, by the first condition,

$$(y - x)x = 36;$$

and by the 2d,

$$xy = 360.$$

---

157. How do you proceed when there are two or more unknown quantities?

From the first equation, we have,

$$xy - x^2 = 36;$$

and by subtraction, $x^2 = 324.$

Hence, $x = \sqrt{324} = 18;$

$$y = \frac{360}{18} = 20.$$

Ans. $x = 18$, $y = 20$.

2. A merchant sells two pieces of muslin, which together measure 12 yards. He received for each piece just so many dollars per yard as the piece contained yards. Now, he gets four times as much for one piece as for the other: how many yards in each piece?

Let x = the number of yards in the larger piece;
y = the number of yards in the shorter piece.

Then, by the conditions of the question,

$$x + y = 12.$$

$x \times x = x^2$ = what he got for the larger piece;
$y \times y = y^2$ = what he got for the shorter;
and, $x^2 = 4y^2$, by the 2d condition,
$x = 2y$, by extracting the square root.

Substituting this value of x in the first equation, we have,

$$y + 2y = 12;$$

and, consequently, $y = 4,$
and, $x = 8.$

Ans. 8 and 4.

3. What two numbers are those whose product is 30, and the quotient of the greater by the less, $3\frac{1}{3}$? *Ans.* 10 and 3.

4. The product of two numbers is a, and their quotient b: what are the numbers?

Ans. $\sqrt{ab}$, and $\sqrt{\frac{a}{b}}$.

5. The sum of the squares of two numbers is 117, and the difference of their squares 45 : what are the numbers?

Ans. 9 and 6.

6. The sum of the squares of two numbers is a, and the difference of their squares is b: what are the numbers?

Ans. $x = \sqrt{\frac{a+b}{2}}, \quad y = \sqrt{\frac{a-b}{2}}.$

7. What two numbers are those which are to each other as 3 to 4, and the sum of whose squares is 225?

Ans. 9 and 12

8. What two numbers are those which are to each other as m to n, and the sum of whose squares is equal to a^2?

Ans. $\frac{ma}{\sqrt{m^2+n^2}}, \quad \frac{na}{\sqrt{m^2+n^2}}.$

9. What two numbers are those which are to each other as 1 to 2, and the difference of whose squares is 75?

Ans. 5 and 10.

10. What two numbers are those which are to each other as m to n, and the difference of whose squares is equal to b^2?

Ans. $\frac{mb}{\sqrt{m^2-n^2}}, \quad \frac{nb}{\sqrt{m^2-n^2}}.$

11. A certain sum of money is placed at interest for six months, at 8 per cent. per annum. Now, if the sum put at interest be multiplied by the number expressing the interest, the product will be $562500: what is the principal at interest?

Ans. $3750.

12. A person distributes a sum of money between a number of women and boys. The number of women is to the number of boys as 3 to 4. Now, the boys receive one-half as many dollars as there are persons, and the women, twice as many dollars as there are boys, and together they receive

138 dollars: how many women were there, and how many boys?

Ans. { 36 women.
48 boys.

COMPLETE EQUATIONS.

158 The reduced form of the complete equation (Art. 153) is,

$$x^2 + 2px = q.$$

Comparing the first member of this equation with the square of a binomial (Art. 54), we see that it needs but the square of half the coefficient of x, to make it a perfect square. Adding p^2 to both members (Ax. 1, Art. 102), we have,

$$x^2 + 2px + p^2 = q + p^2.$$

Then, extracting the square root of both members (Ax. 5), we have,

$$x + p = \pm\sqrt{q + p^2}.$$

Transposing p to the second member, we have,

$$x = -p \pm \sqrt{q + p^2}.$$

Hence, there are two roots, one corresponding to the *plus* sign of the radical, and the other to the *minus* sign. Denoting these roots by x' and x'', we have,

$$x' = -p + \sqrt{q + p^2}, \text{ and } x'' = -p - \sqrt{q + p^2}.$$

The root denoted by x' is called the *first* root; that denoted by x'' is called the *second root.*

---

158. What is the form of the reduced equation of the second degree? What is the square of the binomial $x + p$? How many of those terms are found in the first term of the reduced equation? What must be added to make the first member a perfect square? How many roots are there in every equation of the first degree? What is the first root equal to? What is the second equal to?

159. The operation of squaring half the coefficient of x and adding the result to both members of the equation, is called *Completing the Square.* For the solution of every complete equation of the second degree, we have the following

RULE.

I. *Reduce the equation to the form,* $x^2 + 2px = q$ *:*

II. *Take half the coefficient of the second term, square it, and add the result to both members of the equation:*

III. *Then extract the square root of both members; after which, transpose the known term to the second member.*

NOTE.—Although, in the beginning, the student should complete the square and then extract the square root, yet he should be able, in all cases, to write the roots immediately, by the following (See Art. 158)

RULE.

I. *The first root is equal to half the coefficient of the second term of the reduced equation, taken with a contrary sign, plus the square root of the second member increased by the square of half the coefficient of the second term:*

II. *The second root is equal to half the coefficient of the second term of the reduced equation, taken with a contrary sign, minus the square root of the second member increased by the square of half the coefficient of the second term.*

160. We will now show that the complete equation of

---

159. What is the operation of completing the square? How many operations are there in the solution of every equation of the second degree? What is the first? What the second? What the third? Give the rule for writing the roots without completing the square?

160. How many forms will the complete equation of the second degree assume? On what will these forms depend? What are the signs of $2p$

the second degree will take four forms, dependent on the signs of $2p$ and q.

1st. Let us suppose $2p$ to be positive, and q positive; we shall then have,

$$x^2 + 2px = q. \quad . \quad . \quad . \quad . \quad (1.)$$

2d. Let us suppose $2p$ to be negative, and q positive we shall then have,

$$x^2 - 2px = q. \quad . \quad . \quad . \quad . \quad (2.)$$

3d. Let us suppose $2p$ to be positive, and q negative; we shall then have,

$$x^2 + 2px = -q. \quad . \quad . \quad . \quad (3.)$$

4th. Let us suppose $2p$ to be negative, and q negative; we shall then have,

$$x^2 - 2px = -q. \quad . \quad . \quad . \quad (4.)$$

As these are all the combinations of signs that can take place between $2p$ and q, we conclude that every complete equation of the second degree will be reduced to one or the other of these four forms:

$$x^2 + 2px = +q, \quad . \quad . \quad \text{1st form.}$$
$$x^2 - 2px = +q, \quad . \quad . \quad \text{2d form.}$$
$$x^2 + 2px = -q, \quad . \quad . \quad \text{3d form.}$$
$$x^2 - 2px = -q, \quad . \quad . \quad \text{4th form.}$$

EXAMPLES OF THE FIRST FORM.

1. What are the values of x in the equation,

$$2x^2 + 8x = 64?$$

If we first divide by the coefficient 2, we obtain

$$x^2 + 4x = 32.$$

---

and q in the first form? What in the second? What in the third? What in the fourth?

Then, completing the square,

$$x^2 + 4x + 4 = 32 + 4 = 36.$$

Extracting the root,

$$x + 2 = \pm\sqrt{36} = +6, \text{ and } -6.$$

Hence, $x' = -2 + 6 = +4;$

and, $x'' = -2 - 6 = -8.$

Hence, in this form, the smaller root, numerically, is positive and the larger negative.

VERIFICATION.

If we take the positive value, viz.: $x' = +4$,

the equation, $x^2 + 4x = 32$,

gives $4^2 + 4 \times 4 = 32;$

and if we take the negative value of x, viz.: $x'' = -8$,

the equation, $x^2 + 4x = 32$,

gives $(-8)^2 + 4(-8) = 64 - 32 = 32;$

from which we see that either of the values of x, viz.: $x' = +4$, or $x'' = -8$, will satisfy the equation.

2. What are the values of x in the equation,

$$3x^2 + 12x - 19 = -x^2 - 12x + 89?$$

By transposing the terms, we have,

$$3x^2 + x^2 + 12x + 12x = 89 + 19;$$

and by reducing,

$$4x^2 + 24x = 108;$$

and dividing by the coefficient of x^2,

$$x^2 + 6x = 27.$$

Now, by completing the square,

$$x^2 + 6x + 9 = 36;$$

extracting the square root,

$$x + 3 = \pm\sqrt{36} = +6, \text{ and } -6;$$

hence, $$x' = +6 - 3 = +3;$$

and, $$x'' = -6 - 3 = -9.$$

VERIFICATION.

If we take the plus root, the equation,

$$x^2 + 6x = 27,$$

gives $$(3)^2 + 6(3) = 27;$$

and for the negative root,

$$x^2 + 6x = 27,$$

gives $$(-9)^2 + 6(-9) = 81 - 54 = 27.$$

3. What are the values of x in the equation,

$$x^2 - 10x + 15 = \frac{x^2}{5} - 34x + 155?$$

By clearing of fractions, we have,

$$5x^2 - 50x + 75 = x^2 - 170x + 775;$$

by transposing and reducing, we obtain,

$$4x^2 + 120x = 700;$$

then, dividing by the coefficient of x^2, we have,

$$x^2 + 30x = 175;$$

and by completing the square,

$$x^2 + 30x + 225 = 400;$$

and by extracting the square root,

$$x + 15 = \pm\sqrt{400} = +20, \text{ and } -20.$$

Hence, $x' = +5$, and $x'' = -35$.

VERIFICATION.

For the plus value of x, the equation,

$$x^2 + 30x = 175,$$

gives, $(5)^2 + 30 \times 5 = 25 + 150 = 175.$

And for the negative value of x, we have,

$$(-35)^2 + 30(-35) = 1225 - 1050 = 175.$$

4. What are the values of x in the equation,

$$\frac{5}{6}x^2 - \frac{1}{2}x + \frac{3}{4} = 8 - \frac{2}{3}x - x^2 + \frac{273}{12}?$$

Clearing of fractions, we have,

$$10x^2 - 6x + 9 = 96 - 8x - 12x^2 + 273;$$

transposing and reducing,

$$22x^2 + 2x = 360;$$

dividing both members by 22,

$$x^2 + \frac{2}{22}x = \frac{360}{22}.$$

Add $\left(\frac{1}{22}\right)^2$ to both members, and the equation becomes,

$$x^2 + \frac{2}{22}x + \left(\frac{1}{22}\right)^2 = \frac{360}{22} + \left(\frac{1}{22}\right)^2;$$

whence, by extracting the square root,

$$x + \frac{1}{22} = \pm\sqrt{\frac{360}{22} + \left(\frac{1}{22}\right)^2},$$

therefore,

$$x' = -\frac{1}{22} + \sqrt{\frac{360}{22} + \left(\frac{1}{22}\right)^2},$$

and,

$$x'' = -\frac{1}{22} - \sqrt{\frac{360}{22} + \left(\frac{1}{22}\right)^2}.$$

It remains to perform the numerical operations. In the first place,

$$\frac{360}{22} + \left(\frac{1}{22}\right)^2,$$

must be reduced to a single number, having $(22)^2$ for its denominator. Now,

$$\frac{360}{22} + \left(\frac{1}{22}\right)^2 = \frac{360 \times 22 + 1}{(22)^2} = \frac{7921}{(22)^2};$$

extracting the square root of 7921, we find it to be 89; therefore,

$$\pm\sqrt{\frac{360}{22} + \left(\frac{1}{22}\right)^2} = \pm\frac{89}{22}.$$

Consequently, the plus value of x is,

$$x' = -\frac{1}{22} + \frac{89}{22} = \frac{88}{22} = 4,$$

and the negative value is,

$$x'' = -\frac{1}{22} - \frac{89}{22} = -\frac{45}{11};$$

that is, one of the two values of x which will satisfy the proposed equation is a positive whole number, and the other a negative fraction.

NOTE.—Let the pupil be exercised in writing the roots, in the last five, and in the following examples, *without completing the square.*

5. What are the values of x in the equation,

$$3x^2 + 2x - 9 = 76\,?$$

$$Ans. \begin{cases} x' = 5. \\ x'' = -\,5\tfrac{2}{3}. \end{cases}$$

6. What are the values of x in the equation,

$$2x^2 + 8x + 7 = \frac{5x}{4} - \frac{x^2}{8} + 197\,?$$

$$Ans. \begin{cases} x' = 8. \\ x'' = -\,11\tfrac{3}{17}. \end{cases}$$

7. What are the values of x in the equation,

$$\frac{x^2}{4} - \frac{x}{3} + 15 = \frac{x^2}{9} - 8x + 95\tfrac{1}{4}\,?$$

$$Ans. \begin{cases} x' = 9. \\ x'' = -\,64\tfrac{1}{5}. \end{cases}$$

8. What are the values of x in the equation,

$$\frac{x^2}{1} - \frac{5x}{4} - 8 = \frac{x}{2} - 7x + 6\tfrac{1}{2}\,?$$

$$Ans. \begin{cases} x' = 2. \\ x'' = -\,7\tfrac{1}{4}. \end{cases}$$

9. What are the values of x in the equation,

$$\frac{x^2}{2} + \frac{x}{4} = \frac{x^2}{5} - \frac{x}{10} + \frac{13}{20}$$

$$Ans. \begin{cases} x' = 1. \\ x'' = -\,2\tfrac{1}{6}. \end{cases}$$

EXAMPLES OF THE SECOND FORM.

1. What are the values of x in the equation,

$$x^2 - 8x + 10 = 19\,?$$

By transposing,

$$x^2 - 8x = 19 - 10 = 9;$$

then, by completing the square,

$$x^2 - 8x + 16 = 9 + 16 = 25;$$

and by extracting the root,

$$x - 4 = \pm\sqrt{25} = +5, \text{ or } -5.$$

Hence,

$$x' = 4 + 5 = 9, \text{ and } x'' = 4 - 5 = -1.$$

That is, in this form, the larger root, numerically, is positive, and the lesser negative.

VERIFICATION.

If we take the positive value of x, the equation,

$$x^2 - 8x = 9, \text{ gives } (9)^2 - 8 \times 9 = 81 - 72 = 9;$$

and if we take the negative value, the equation,

$$x^2 - 8x = 9, \text{ gives } (-1)^2 - 8(-1) = 1 + 8 = 9;$$

from which we see that both roots alike satisfy the equation.

2. What are the values of x in the equation,

$$\frac{x^2}{2} + \frac{x}{3} - 15 = \frac{x^2}{4} + x - 14\tfrac{3}{4}?$$

By clearing of fractions, we have,

$$6x^2 + 4x - 180 = 3x^2 + 12x - 177,$$

and by transposing and reducing,

$$3x^2 - 8x = 3;$$

and dividing by the coefficient of x^2, we obtain,

$$x^2 - \frac{8}{3}x = 1.$$

Then, by completing the square, we have,

$$x^2 - \frac{8}{3}x + \frac{16}{9} = 1 + \frac{16}{9} = \frac{25}{9};$$

and by extracting the square root,

$$x - \frac{4}{3} = \pm\sqrt{\frac{25}{9}} = +\frac{5}{3}, \text{ and } -\frac{5}{3}$$

Hence,

$$x' = \frac{4}{3} + \frac{5}{3} = +3, \text{ and } x'' = \frac{4}{3} - \frac{5}{3} = -\frac{1}{3}.$$

VERIFICATION.

For the positive root of x, the equation,

$$x^2 - \frac{8}{3}x = 1,$$

gives $$3^2 - \frac{8}{3} \times 3 = 9 - 8 = 1;$$

and for the negative root, the equation,

$$x^2 - \frac{8}{3}x = 1,$$

gives $$\left(-\frac{1}{3}\right)^2 - \frac{8}{3} \times -\frac{1}{3} = \frac{1}{9} + \frac{8}{9} = 1.$$

3. What are the values of x in the equation,

$$\frac{x^2}{2} - \frac{x}{3} + 7\tfrac{3}{8} = 8?$$

Clearing of fractions, and dividing by the coefficient of x^2, we have,

$$x^2 - \frac{2}{3}x = 1\tfrac{1}{4}.$$

Completing the square, we have,

$$x^2 - \frac{2}{3}x + \frac{1}{9} = 1\tfrac{1}{4} + \frac{1}{9} = \frac{49}{36};$$

then, by extracting the square root, we have,

$$x - \frac{1}{3} = \pm\sqrt{\frac{49}{36}} = +\frac{7}{6}, \text{ and } -\frac{7}{6};$$

hence,

$$x' = \frac{1}{3} + \frac{7}{6} = \frac{9}{6} = 1\tfrac{1}{2}, \text{ and } x'' = \frac{1}{3} - \frac{7}{6} = -\frac{5}{6}.$$

VERIFICATION.

If we take the positive root of x, the equation,

$$x^2 - \frac{2}{3}x = 1\tfrac{1}{4},$$

gives $\quad (1\tfrac{1}{2})^2 - \frac{2}{3} \times 1\tfrac{1}{2} = 2\tfrac{1}{4} - 1 = 1\tfrac{1}{4};$

and for the negative root, the equation,

$$x^2 - \frac{2}{3}x = 1\tfrac{1}{4},$$

gives $\left(-\frac{5}{6}\right)^2 - \frac{2}{3} \times -\frac{5}{6} = \frac{25}{36} + \frac{10}{18} = \frac{45}{36} = 1\tfrac{1}{4}$

4. What are the values of x in the equation,

$$4a^2 - 2x^2 + 2ax = 18ab - 18b^2?$$

By transposing, changing the signs, and dividing by 2, the equation becomes,

$$x^2 - ax = 2a^2 - 9ab + 9b^2;$$

whence, completing the square,

$$x^2 - ax + \frac{a^2}{4} = \frac{9a^2}{4} - 9ab + 9b^2;$$

extracting the square root,

$$x = \frac{a}{2} \pm \sqrt{\frac{9a^2}{4} - 9ab + 9b^2}.$$

Now, the square root of $\frac{9a^2}{4} - 9ab + 9b^2$, is evidently $\frac{3a}{2} - 3b$. Therefore,

$$x = \frac{a}{2} \pm \left(\frac{3a}{2} - 3b\right), \text{ and } \begin{cases} x' = 2a - 3b. \\ x'' = - a + 3b. \end{cases}$$

What will be the numerical values of x, if we suppose $a = 6$, and $b = 1$?

5. What are the values of x in the equation,

$$\frac{1}{3}x - 4 - x^2 + 2x - \frac{4}{5}x^2 = 45 - 3x^2 + 4x?$$

$$Ans. \begin{cases} x' = 7.12 \\ x'' = -5.73 \end{cases} \text{to within } 0.01.$$

6. What are the values of x in the equation,

$$8x^2 - 14x + 10 = 2x + 34?$$

$$Ans. \begin{cases} x' = 3. \\ x'' = -1. \end{cases}$$

7. What are the values of x in the equation,

$$\frac{x^2}{4} - 30 + x = 2x - 22?$$

$$Ans. \begin{cases} x' = 8. \\ x'' = -4. \end{cases}$$

8. What are the values of x in the equation,

$$x^2 - 3x + \frac{x^2}{2} = 9x + 13\tfrac{1}{2}?$$

$$Ans. \begin{cases} x' = 9. \\ x'' = -1. \end{cases}$$

9. What are the values of x in the equation,

$$2ax - x^2 = -2ab - b^2?$$

$$Ans. \begin{cases} x' = 2a + b \\ x'' = -b. \end{cases}$$

10. What are the values of x in the equation,

$$a^2 + b^2 - 2bx + x^2 = \frac{m^2x^2}{n^2}?$$

$$Ans. \begin{cases} x' = \dfrac{n}{n^2 - m^2}\left(bn + \sqrt{a^2m^2 + b^2m^2 - a^2n^2}\right) \\ x'' = \dfrac{n}{n^2 - m^2}\left(bn - \sqrt{a^2m^2 + b^2m^2 - a^2n^2}\right) \end{cases}$$

EXAMPLES OF THE THIRD FORM.

1. What are the values of x in the equation,

$$x^2 + 4x = -3?$$

First, by completing the square, we have,

$$x^2 + 4x + 4 = -3 + 4 = 1;$$

and by extracting the square root,

$$x + 2 = \pm\sqrt{1} = +1, \text{ and } -1;$$

hence, $x' = -2 + 1 = -1$; and $x'' = -2 - 1 = -3$

That is, in this form both the roots are negative.

VERIFICATION.

If we take the first negative value, the equation,

$$x^2 + 4x = -3,$$

gives $(-1)^2 + 4(-1) = 1 - 4 = -3;$

and by taking the second value, the equation,

$$x^2 + 4x = -3,$$

gives $(-3)^2 + 4(-3) = 9 - 12 = -3;$

hence, both values of x satisfy the given equation.

2. What are the values of x in the equation,

$$-\frac{x^2}{2} - 5x - 16 = 12 + \frac{1}{2}x^2 + 6x\,?$$

By transposing and reducing, we have,

$$-x^2 - 11x = 28;$$

then, dividing by -1, the coefficient of x^2, we have,

$$x^2 + 11x = -28;$$

then, by completing the square,

$$x^2 + 11x + 30.25 = 2.25;$$

hence, $x + 5.5 = \pm\sqrt{2.25} = +1.5$, and -1.5;

consequently, $x' = -4$, and $x'' = -7$.

3. What are the values of x in the equation,

$$-\frac{x^2}{8} - 2x - 5 = \frac{7}{8}x^2 + 5x + 5\,?$$

$$Ans. \begin{cases} x' = -2. \\ x'' = -5. \end{cases}$$

4. What are the values of x in the equation,

$$2x^2 + 8x = -2\tfrac{2}{3} - \frac{2}{3}x\,?$$

$$Ans. \begin{cases} x' = -\tfrac{1}{3}. \\ x'' = -4. \end{cases}$$

5. What are the values of x in the equation,

$$4x^2 + \frac{3}{5}x + 3x = -14x - 3\tfrac{1}{5} - 4x^2\,?$$

$$Ans. \begin{cases} x' = -\tfrac{1}{5}. \\ x'' = -2. \end{cases}$$

6. What are the values of x in the equation,

$$-x^2 - 4 - \frac{3}{4}x = \frac{4x^2}{2} + 24x + 2?$$

$$Ans. \begin{cases} x' = -\frac{1}{4}. \\ x'' = -8. \end{cases}$$

7. What are the values of x in the equation,

$$\frac{1}{9}x^2 + 7x + 20 = -\frac{8}{9}x^2 - 11x - 60?$$

$$Ans. \begin{cases} x' = -8. \\ x'' = -10. \end{cases}$$

8. What are the values of x in the equation,

$$\frac{5}{6}x^2 - x + \frac{1}{2} = -9\tfrac{1}{8}x - \frac{1}{6}x^2 - \frac{1}{2}?$$

$$Ans. \begin{cases} x' = -\frac{1}{8}. \\ x'' = -8. \end{cases}$$

9. What are the values of x in the equation,

$$\frac{4}{5}x^2 + 5x + \frac{1}{4} = -\frac{1}{5}x^2 - 5\tfrac{1}{10}x - \frac{3}{4}?$$

$$Ans. \begin{cases} x' = -\frac{1}{10}. \\ x'' = -10. \end{cases}$$

10. What are the values of x in the equation,

$$x - x^2 - 3 = 6x + 1?$$

$$Ans. \begin{cases} x' = -1. \\ x'' = -4. \end{cases}$$

11. What are the values of x in the equation,

$$x^2 + 4x - 90 = -93?$$

$$Ans. \begin{cases} x' = -1. \\ x'' = -3. \end{cases}$$

EXAMPLES OF THE FOURTH FORM.

1. What are the values of x in the equation,

$$x^2 - 8x = -7?$$

By completing the square, we have,

$$x^2 - 8x + 16 = -7 + 16 = 9$$

then, by extracting the square root,

$$x - 4 = \pm\sqrt{9} = +3, \text{ and } -3;$$

hence, $x' = +7$, and $x'' = +1$.

That is, in this form, both the roots are positive.

VERIFICATION.

If we take the greater root, the equation,

$x^2 - 8x = -7$, gives, $7^2 - 8 \times 7 = 49 - 56 = -7$;

and for the lesser, the equation,

$x^2 - 8x = -7$, gives, $1^2 - 8 \times 1 = 1 - 8 = -7$

hence, both of the roots will satisfy the equation.

2. What are the values of x in the equation,

$$-1\tfrac{1}{2}x^2 + 3x - 10 = 1\tfrac{1}{2}x^2 - 18x + \frac{40}{2}?$$

By clearing of fractions, we have,

$$-3x^2 + 6x - 20 = 3x^2 - 36x + 40;$$

then, by collecting the similar terms,

$$-6x^2 + 42x = 60;$$

then, by dividing by the coefficient of x^2, which is -6, we have,

$$x^2 - 7x = -10.$$

By completing the square, we have,

$$x^2 - 7x + 12.25 = 2.25,$$

and by extracting the square root of both members,

$$x - 3.5 = \pm\sqrt{2.25} = +1.5, \text{ and } -1.5;$$

hence,

$$x' = 3.5 + 1.5 = 5, \text{ and } x'' = 3.5 - 1.5 = 2.$$

VERIFICATION.

If we take the greater root, the equation.

$$x^2 - 7x = -10, \text{ gives, } 5^2 - 7 \times 5 = 25 - 35 = -10;$$

and if we take the lesser root, the equation,

$$x^2 - 7x = -10, \text{ gives, } 2^2 - 7 \times 2 = 4 - 14 = -10.$$

3. What are the values of x in the equation,

$$-3x + 2x^2 + 1 = 17\tfrac{4}{5}x - 2x^2 - 3?$$

By transposing and collecting the terms, we have,

$$4x^2 - 20\tfrac{4}{5}x = -4;$$

then dividing by the coefficient of x^2, we have,

$$x^2 - 5\tfrac{1}{5}x = -1.$$

By completing the square, we obtain,

$$x^2 - 5\tfrac{1}{5}x + \frac{169}{25} = -1 + \frac{169}{25} = \frac{144}{25};$$

and by extracting the root,

$$x^2 - 2\tfrac{3}{5} = \pm\sqrt{\frac{144}{25}} = +\frac{12}{5}, \text{ and } -\frac{12}{5};$$

hence,

$$x' = 2\tfrac{3}{5} + \frac{12}{5} = 5, \text{ and, } x'' = 2\tfrac{3}{5} - \frac{12}{5} = \frac{1}{5}.$$

VERIFICATION.

If we take the greater root, the equation,

$$x^2 - 5\tfrac{1}{5}x = -1, \text{ gives, } 5^2 - 5\tfrac{1}{5} \times 5 = 25 - 26 = -1;$$

and if we take the lesser root, the equation,

$$x^2 - 5\tfrac{1}{5}x = -1, \text{ gives, } \left(\frac{1}{5}\right)^2 - 5\tfrac{1}{5} \times \frac{1}{5} = \frac{1}{25} - \frac{26}{25} = -1.$$

4. What are the values of x in the equation,

$$\frac{1}{7}x^2 - 3x + \frac{1}{2} = -\frac{6}{7}x^2 + \frac{1}{4}x - \frac{1}{4}?$$

$$Ans. \begin{cases} x' = 3 \\ x'' = \tfrac{1}{4}. \end{cases}$$

5. What are the values of x in the equation,

$$-4x^2 - \frac{1}{7}x + 1\tfrac{1}{7} = -5x^2 + 8x?$$

$$Ans. \begin{cases} x' = 8. \\ x'' = \tfrac{1}{7}. \end{cases}$$

6. What are the values of x in the equation,

$$-4x^2 + \frac{8}{20}x - \frac{1}{40} = -3x^2 - \frac{1}{20}x + \frac{1}{40}?$$

$$Ans. \begin{cases} x' = \tfrac{1}{4}. \\ x'' = \tfrac{1}{5}. \end{cases}$$

7. What are the values of x in the equation,

$$x^2 - 10\tfrac{1}{10}x = -1?$$

$$Ans. \begin{cases} x' = 10. \\ x'' = \tfrac{1}{10}. \end{cases}$$

8. What are the values of x in the equation,

$$-27x + \frac{17x^2}{5} + 100 = \frac{2x^2}{5} + 12x - 26?$$

$$Ans. \begin{cases} x' = 7. \\ x'' = 6. \end{cases}$$

9. What are the values of x in the equation,

$$\frac{8x^2}{3} - 22x + 15 = -\frac{7x^2}{3} + 28x - 30?$$

$$Ans. \begin{cases} x' = 9. \\ x'' = 1. \end{cases}$$

10. What are the values of x in the equation,

$$2x^2 - 30x + 3 = -x^2 + 3\tfrac{3}{10}x - \frac{3}{10}?$$

$$Ans. \begin{cases} x' = 11. \\ x'' = \tfrac{1}{10}. \end{cases}$$

PROPERTIES OF EQUATIONS OF THE SECOND DEGREE.

FIRST PROPERTY.

161. We have seen (Art. 153), that every complete equation of the second degree may be reduced to the form,

$$x^2 + 2px = q. \quad . \quad . \quad . \quad . \quad (1.)$$

Completing the square, we have,

$$x^2 + 2px + p^2 = q + p^2;$$

transposing $q + p^2$ to the first member,

$$x^2 + 2px + p^2 - (q + p^2) = 0. \quad . \quad (2.)$$

Now, since $x^2 + 2px + p^2$ is the square of $x + p$, and $q + p^2$ the square of $\sqrt{q + p^2}$, we may regard the first member as the difference between two squares. Factoring, (Art. 56), we have,

$$(x + p + \sqrt{q + p^2})\,(x + p - \sqrt{q + p^2}) = 0. \quad . \quad (3.)$$

This equation can be satisfied only in two ways:

1st. By attributing such a value to x as shall render the first factor equal to 0; or,

---

161. To what form may every equation of the second degree be reduced? What form will this equation take after completing the square and transposing to the first member? After factoring? In how many ways may Equation (3) be satisfied? What are they? How many roots has every equation of the second degree?

2d. By attributing such a value to x as shall render the second factor equal to 0.

Placing the second factor equal to 0, we have,

$$x + p - \sqrt{q+p^2} = 0; \text{ and } x' = -p + \sqrt{q+p^2}. \qquad (4.)$$

Placing the first factor equal to 0, we have,

$$+p + \sqrt{q+p^2} = 0; \text{ and } x'' = -p - \sqrt{q+p^2}. \qquad (5.)$$

Since every supposition that will satisfy Equation (3), will also satisfy Equation (1), from which it was derived, it follows, that x' and x'' are roots of Equation (1); also, that

Every equation of the second degree has two roots, and only two.

NOTE.—The two roots denoted by x' and x'', are the same as found in Art. 158.

SECOND PROPERTY.

162. We have seen (Art. 161), that every equation of the second degree may be placed under the form,

$$(x + p + \sqrt{q + p^2})\,(x + p - \sqrt{q + p^2}) = 0.$$

By examining this equation, we see that the first factor may be obtained by subtracting the *second* root from the unknown quantity x; and the second factor by subtracting the *first* root from the unknown quantity x; hence,

Every equation of the second degree may be resolved into two binomial factors of the first degree, the first terms, in both factors, being the unknown quantity, and the second terms, the roots of the equation, taken with contrary signs.

---

162. Into how many binomial factors of the first degree may every equation of the second degree be resolved? What are the first terms of these factors? What the second?

THIRD PROPERTY.

163. If we add Equations (4) and (5), Art. 161, we have,

$$\begin{aligned} x' &= -p + \sqrt{q+p^2} \\ x'' &= -p - \sqrt{q+p^2} \\ \hline x' + x'' &= -2p; \end{aligned} \qquad \text{that is,}$$

In every reduced equation of the second degree, the sum of the two roots is equal to the coefficient of the second term taken with a contrary sign.

FOURTH PROPERTY.

164. If we multiply Equations (4) and (5), Art. 161, member by member, we have,

$$\begin{aligned} x' \times x'' &= (-p + \sqrt{q+p^2})(-p - \sqrt{q+p^2}) \\ &= p^2 - (q+p^2) = -q; \text{ that is,} \end{aligned}$$

In every equation of the second degree, the product of the two roots is equal to the known term in the second member, taken with a contrary sign.

FORMATION OF EQUATIONS OF THE SECOND DEGREE.

165. By taking the converse of the second property, (Art. 162), we can form equations which shall have given roots; that is, if they are known, we can find the corresponding equations by the following

RULE.

I. *Subtract each root from the unknown quantity:*

---

163. What is the algebraic sum of the roots equal to in every equation of the second degree?

164. What is the product of the roots equal to?

165. How will you find the equation when the roots are known?

II. *Multiply the results together, and place their product equal to* 0.

EXAMPLES.

Note.—Let the pupil prove, *in every case*, that the roots will satisfy the third and fourth properties.

1. If the roots of an equation are 4 and -5, what is the equation? *Ans.* $x^2 + x = 20$.

2. What is the equation when the roots are 1 and -3? *Ans.* $x^2 + 2x = 3$.

3. What is the equation when the roots are 9 and -10? *Ans.* $x^2 + x = 90$.

4. What is the equation whose roots are 6 and -10? *Ans.* $x^2 + 4x = 60$.

5. What is the equation whose roots are 4 and -3? *Ans.* $x^2 - x = 12$.

6. What is the equation whose roots are 10 and $-\frac{1}{10}$? *Ans.* $x^2 - 9\frac{9}{10}x = 1$.

7. What is the equation whose roots are 8 and -2? *Ans.* $x^2 - 6x = 16$.

8. What is the equation whose roots are 16 and -5? *Ans.* $x^2 - 11x = 80$

9. What is the equation whose roots are -4 and -5? *Ans.* $x^2 + 9x = -20$.

10. What is the equation whose roots are -6 and -7? *Ans.* $x^2 + 13x = -42$.

11. What is the equation whose roots are $-\frac{3}{4}$ and -2? *Ans.* $x^2 + 2\frac{3}{4}x = -\frac{3}{2}$.

12. What is the equation whose roots are -2 and -3? *Ans.* $x^2 + 5x = -6$.

13. What is the equation whose roots are 4 and 3?

Ans. $x^2 - 7x = -12$.

14. What is the equation whose roots are 12 and 2?

Ans. $x^2 - 14x = -24$.

15. What is the equation whose roots are 18 and 2?

Ans. $x^2 - 20x = -36$.

16. What is the equation whose roots are 14 and 3?

Ans. $x^2 - 17x = -42$.

17. What is the equation whose roots are $\frac{4}{9}$ and $-\frac{9}{4}$?

Ans. $x^2 + \frac{65}{36}x = 1$.

18. What is the equation whose roots are 5 and $-\frac{2}{3}$?

Ans. $x^2 - \frac{13}{3}x = \frac{10}{3}$.

19. What is the equation whose roots are a and b?

Ans. $x^2 - (a + b)x = -ab$.

20. What is the equation whose roots are c and $-d$?

Ans. $x^2 - (c - d)x = cd$.

TRINOMIAL EQUATIONS OF THE SECOND DEGREE.

165.* A trinomial equation of the second degree contains three kinds of terms:

1st. A term involving the unknown quantity to the second degree.

2d. A term involving the unknown quantity to the first degree; and

3d. A known term. Thus,

$$x^2 - 4x - 12 = 0,$$

is a trinomial equation of the second degree.

FACTORING.

165.** What are the factors of the trinomial equation,

$$x^2 - 4x - 12 = 0?$$

A trinomial equation of the second degree may always be reduced to one of the four forms (Art. 160), by simply transposing the known term to the second member, and then solving the equation. Thus, from the above equation, we have,

$$x^2 - 4x = 12.$$

Resolving the equation, we find the two roots to be $+6$ and -2; therefore~~, the~~ factors are, $x - 6$, and $x + 2$ (Art. 162).

Since the sum of the two roots is equal to the coefficient of the second term, taken with a contrary sign (Art. 163); and the product of the two roots is equal to the known term in the second member, taken with a contrary sign, or to the third term of the trinomial, taken with the *same sign:* hence it follows, that any trinomial may be factored by inspection, when two numbers can be discovered *whose algebraic sum is equal to the coefficient of the second term, and whose product is equal to the third term.*

EXAMPLES

1. What are the factors of the trinomial, $x^2 - 9x - 36$?

It is seen, by inspection, that -12 and $+3$ will fulfil the conditions of roots. For, $12 - 3 = 9$; that is, the coefficient of the second term with a contrary sign; and $12 \times -3 = -36$, the third term of the trinomial; hence, the factors are, $x - 12$, and $x + 3$.

2. What are the factors of $x^2 - 7x - 30 = 0$?

Ans. $x - 10$, and $x + 3$

3. What are the factors of $x^2 + 15x + 36 = 0$?

Ans. $x + 12$, and $x + 3$.

4. What are the factors of $x^2 - 12x - 28 = 0$?

Ans. $x - 14$, and $x + 2$.

5. What are the factors of $x^2 - 7x - 8 = 0$?

Ans. $x - 8$, and $x + 1$.

TRINOMIAL EQUATIONS OF THE FORM

$$x^{2n} + 2px^n = q.$$

In the above equation, the exponent of x, in the first term, is double the exponent of x in the second term.

$$x^6 - 4x^3 = 32, \quad \text{and} \quad x^4 + 4x^2 = 117,$$

are both equations of this form, and may be solved by the rules already given for the solution of equations of the second degree.

In the equation,

$$x^{2n} + 2px^n = q,$$

we see that the first member will become a perfect square, by adding to it the square of half the coefficient of x^n; thus,

$$x^{2n} + 2px^n + p^2 = q + p^2,$$

in which the first member is a perfect square. Then, extracting the square root of both members, we have,

$$x^n + p = \pm \sqrt{q + p^2};$$

hence,

$$x^n = -p \pm \sqrt{q + p^2};$$

then, by taking the nth root of both members,

$$x' = \sqrt[n]{-p + \sqrt{q + p^2}},$$

and

$$x'' = \sqrt[n]{-p - \sqrt{-p + p^2}}.$$

EXAMPLES.

1. What are the values of x in the equation.

$$x^6 + 6x^3 = 112?$$

Completing the square,

$$x^6 + 6x^3 + 9 = 112 + 9 = 121;$$

then, extracting the square root of both members,

$$x^3 + 3 = \pm \sqrt{121} = \pm 11; \text{ hence,}$$

$$x' = \sqrt[3]{-3 + 11}, \quad \text{and} \quad x'' = \sqrt[3]{-3 - 11}; \text{ hence,}$$

$$x' = \sqrt[3]{8} = 2, \quad \text{and} \quad x'' = \sqrt[3]{-14} = -\sqrt[3]{14}.$$

2. What are the values of x in the equation,

$$x^4 - 8x^2 = 9?$$

Completing the square, we have,

$$x^4 - 8x^2 + 16 = 9 + 16 = 25.$$

Extracting the square root of both members,

$$x^2 - 4 = \pm \sqrt{25} = \pm 5; \text{ hence,}$$

$$x' = \pm \sqrt{4 + 5}, \quad \text{and} \quad x'' = \pm \sqrt{4 - 5}; \text{ hence,}$$

$$x' = +3 \text{ and } -3; \text{ and } x'' = +\sqrt{-1} \text{ and } -\sqrt{-1}.$$

3. What are the values of x in the equation,

$$x^6 + 20x^3 = 69?$$

Completing the square,

$$x^6 + 20x^3 + 100 = 69 + 100 = 169.$$

Extracting the square root of both members,

$$x^3 + 10 = \pm \sqrt{169} = \pm 13; \text{ hence,}$$

$$x' = \sqrt[3]{-10 + 13}, \quad \text{and} \quad x'' = \sqrt[3]{-10 - 13}.$$

$$x' = \sqrt[3]{3}, \quad \text{and} \quad x'' = \sqrt[3]{-23}.$$

4. What are the values of x in the equation,

$$x^4 - 2x^2 = 3?$$

Ans. $x' = \pm \sqrt{3}$, and $x'' = \pm \sqrt{-1}$.

5. What are the values of x in the equation,

$$x^6 + 8x^3 = 9?$$

Ans. $x' = 1$, and $x'' = \sqrt[3]{-9}$.

6. Given $x \pm \sqrt{9x + 4} = 12$, to find x.

Transposing x to the second member, and then squaring,

$$9x + 4 = x^2 - 24x + 144;$$

$$\therefore \quad x^2 - 33x = -140;$$

and, $\quad x' = 28$, and $x'' = 5$.

7. $4x \pm 4\sqrt{x + 2} = 7$. *Ans.* $x' = 4\frac{1}{4}$, $x'' = \frac{1}{4}$.

8. $x \pm \sqrt{5x + 10} = 8$. *Ans.* $x' = 18$, $x'' = 3$.

NUMERICAL VALUES OF THE ROOTS.

166. We have seen (Art. 160), that by attributing all possible signs to $2p$ and q, we have the four following forms:

$$x^2 + 2px = q. \quad \ldots\ldots \quad (1.)$$

$$x^2 - 2px = q. \quad \ldots\ldots \quad (2.)$$

$$x^2 + 2px = -q. \quad \ldots\ldots \quad (3.)$$

$$x^2 - 2px = -q. \quad \ldots\ldots \quad (4.)$$

---

166. To how many forms may every equation of the second degree be reduced? What are they?

First Form.

167. Since q is positive, we know, from Property Fourth, that the product of the roots must be negative· hence, *the roots have contrary signs.* Since the coefficient $2p$ is positive, we know, from Property Third, that the algebraic sum of the roots is negative; hence, *the negative root is numerically the greater.*

Second Form.

168. Since q is positive, the product of the roots must be negative; hence, *the roots have contrary signs.* Since $2p$ is negative, the algebraic sum of the roots must be positive; hence, *the positive root is numerically the greater.*

Third Form.

169. Since q is negative, the product of the roots is positive (Property Fourth); hence, *the roots have the same sign.* Since $2p$ is positive, the sum of the roots must be negative; hence, *both are negative.*

Fourth Form.

170. Since q is negative, the product of the roots is positive; hence, *the roots have the same sign.* Since $2p$ is negative, the sum of the roots is positive; hence, *the roots are both positive.*

---

167. What sign has the product of the roots in the first form? How are their signs? Which root is numerically the greater? Why?

168. What sign has the product of the roots in the second form? How are the signs of the roots? Which root is numerically the greater?

169. What sign has the product of the roots in the third form? How are their signs?

170. What sign has the product of the roots in the fourth form? How are the signs of the roots?

First and Second Forms.

171. If we make $q = 0$, the first form becomes,

$$x^2 + 2px = 0, \quad \text{or} \quad x(x + 2p) = 0;$$

which shows that one root is equal to 0, and the other to $-2p$.

Under the same supposition, the second form becomes,

$$x^2 - 2px = 0, \quad \text{or} \quad x(x - 2p) = 0;$$

which shows that one root is equal to 0, and the other to $2p$. Both of these results are as they should be; since, when q, the product of the roots, becomes 0, one of the factors must be 0; and hence, one root must be 0.

Third and Fourth Forms.

172. If, in the Third and Fourth Forms, $q > p^2$, the quantity under the radical sign will become *negative;* hence, *its square root cannot be extracted* (Art. 137). Under this supposition, the values of x are *imaginary.* How are these results to be interpreted?

If a given number be divided into two parts, their product will be the greatest possible, when the parts are equal.

Denote the number by $2p$, and the difference of the parts by d; then,

$$p + \frac{d}{2} = \text{the greater part,} \quad \text{(Page 120.)}$$

and, $$p - \frac{d}{2} = \text{the less part,}$$

and, $$p^2 - \frac{d^2}{4} = P, \text{ their product.}$$

---

171. If we make $q = 0$, to what does the first form reduce? What, then, are its roots? Under the same supposition, to what does the second form reduce? What are, then, its roots?

172. If $q > p^2$, in the third and fourth forms, what takes place?

If a number be divided into two parts, when will the product be the greatest possible?

It is plain. that the product P will *increase*, as d *diminishes*, and that it will be the *greatest possible* when $d = 0$; for then there will be no negative quantity to be subtracted from p^2, in the first member of the equation. But when $d = 0$, the parts are equal; hence, *the product of the two parts is the greatest when they are equal.*

In the equations,

$$x^2 + 2px = -q, \quad x^2 - 2px = -q,$$

$2p$ is the sum of the roots, and $-q$ their product; and hence, by the principle just established, the product q, can never be greater than p^2. This condition fixes a limit to the value of q. If, then, we make $q > p^2$, we pass this limit, and express, by the equation, a condition which cannot be fulfilled; and this incompatibility of the conditions is made apparent by the values of x becoming imaginary. Hence, we conclude that,

When the values of the unknown quantity are imaginary, the conditions of the proposition are incompatible with each other.

EXAMPLES.

1. Find two numbers, whose sum shall be 12 and product 46.

Let x and y be the numbers.

By the 1st condition, $x + y = 12$;

and by the 2d, $xy = 46$.

The first equation gives,

$$x = 12 - y.$$

Substituting this value for x in the second, we have,

$$12y - y^2 = 46;$$

and changing the signs of the terms, we have,

$$y^2 - 12y = -46$$

Then, by completing the square,

$$y^2 - 12y + 36 = -46 + 36 = -10;$$

which gives, $y' = 6 + \sqrt{-10},$

and. $y'' = 6 - \sqrt{-10};$

both of which values are imaginary, as indeed they should be, since the conditions are incompatible.

2. The sum of two numbers is 8, and their product 20: what are the numbers?

Denote the numbers by x and y.

By the first condition,

$$x + y = 8;$$

and by the second, $xy = 20.$

The first equation gives,

$$x = 8 - y.$$

Substituting this value of x in the second, we have,

$$8y - y^2 = 20;$$

changing the signs, and completing the square, we have,

$$y^2 - 8y + 16 = -4;$$

and by extracting the root,

$$y' = 4 + \sqrt{-4}, \text{ and } y'' = 4 - \sqrt{-4}.$$

These values of y may be put under the forms (Art. 142),

$$y = 4 + 2\sqrt{-1}, \text{ and } y = 4 - 2\sqrt{-1}.$$

3 What are the values of x in the equation,

$$x^2 + 2x = -10?$$

$$Ans. \begin{cases} x' = -1 + 3\sqrt{-1}. \\ x'' = -1 - 3\sqrt{-1}. \end{cases}$$

PROBLEMS.

1. Find a number such, that twice its square, added to three times the number, shall give 65.

Let x denote the unknown number. Then, the equation of the problem is,

$$2x^2 + 3x = 65;$$

whence,

$$x = -\frac{3}{4} \pm \sqrt{\frac{65}{2} + \frac{9}{16}} = -\frac{3}{4} \pm \frac{23}{4}.$$

Therefore,

$$x' = -\frac{3}{4} + \frac{23}{4} = 5, \quad \text{and} \quad x'' = -\frac{3}{4} - \frac{23}{4} = -\frac{13}{2}.$$

Both these values satisfy the equation of the problem. For,

$$2 \times (5)^2 + 3 \times 5 = 2 \times 25 + 15 = 65;$$

and, $$2\left(-\frac{13}{2}\right)^2 + 3 \times -\frac{13}{2} = \frac{169}{2} - \frac{39}{2} = \frac{130}{2} = 65.$$

NOTES.—1. If we restrict the enunciation of the problem to its arithmetical sense, in which "added" means *numerical increase*, the first value of x only will satisfy the conditions of the problem.

2. If we give to "added," its algebraical signification (when it may mean subtraction as well as addition), the problem may be thus stated:

To find a number such, that twice its square diminished by three times the number, shall give 65.

The second value of x will satisfy this enunciation; for,

$$2\left(\frac{13}{2}\right)^2 - 3 \times \frac{13}{2} = \frac{169}{2} - \frac{39}{2} = 65.$$

3. The root which results from giving the plus sign to the radical, is, generally, an answer to the question in its arithmetical sense. The second root generally satisfies the problem under a modified statement.

Thus, in the example, it was required to find a number, of which twice the square, *added* to three times the number, shall give 65. Now, in the arithmetical sense, *added* means increased; but in the algebraic sense, it implies diminution when the quantity added is negative. In this sense, the second root satisfies the enunciation.

2. A certain person purchased a number of yards of cloth for 240 cents. If he had purchased 3 yards *less* of the same cloth for the same sum, it would have cost him 4 cents more per yard: how many yards did he buy?

Let x denote the number of yards purchased.

Then, $\frac{240}{x}$ will denote the price per yard.

If, for 240 cents, he had purchased three yards less, that is, $x - 3$ yards, the price per yard, under this hypothesis, would have been denoted by $\frac{240}{x-3}\cdot$ But, by the conditions, this last cost must exceed the first by 4 cents. Therefore, we have the equation,

$$\frac{240}{x-3} - \frac{240}{x} = 4;$$

whence, by reducing, $x^2 - 3x = 180,$

and, $x = \frac{3}{2} \pm \sqrt{\frac{9}{4} + 180} = \frac{3 \pm 27}{2};$

therefore, $x' = 15$, and $x'' = -12.$

NOTES.—1. The value, $x' = 15$, satisfies the enunciation in its arithmetical sense. For, if 15 yards cost 240 cents,

$240 \div 15 = 16$ cents, the price of 1 yard; and $240 \div 12 = 20$ cents, the price of 1 yard under the second supposition.

2. The second value of x is an answer to the following Problem:

A certain person purchased a number of yards of cloth for 240 cents. If he had paid the same for three yards *more*, it would have cost him 4 cents *less* per yard: how many yards did he buy?

This would give the equation of condition,

$$\frac{240}{x} - \frac{240}{x+3} = 4; \text{ or,}$$

$$x^2 - 3x = 180;$$

the same equation as found before; hence,

A single equation will often state two or more arithmetical problems.

This arises from the fact that the language of Algebra is more comprehensive than that of Arithmetic.

3. A man having bought a horse, sold it for $24. At the sale he lost as much per cent. on the price of the horse, as the horse cost him dollars: what did he pay for the horse?

Let x denote the number of dollars that he paid for the horse. Then, $x - 24$ will denote the loss he sustained. But as he lost x per cent. by the sale, he must have lost $\frac{x}{100}$ upon each dollar, and upon x dollars he lost a sum denoted by $\frac{x^2}{100}$; we have, then, the equation,

$$\frac{x^2}{100} = x - 24; \text{ whence, } x^2 - 100x = -2400,$$

and, $x = 50 \pm \sqrt{2500 - 2400} = 50 \pm 10.$

Therefore, $x' = 60,$ and $x'' = 40.$

Both of these roots will satisfy the problem.

For, if the man gave \$60 for the horse, and sold him for \$24, he lost \$36. From the enunciation, he should have lost 60 per cent. of \$60; that is,

$$\frac{60}{100} \text{ of } 60 = \frac{60 \times 60}{100} = 36;$$

therefore, \$60 satisfies the enunciation.

Had he paid \$40 for the horse, he would have lost by the sale, \$16. From the enunciation, he should have lost 40 per cent. of \$40; that is,

$$\frac{40}{100} \text{ of } 40 = \frac{40 \times 40}{100} = 16;$$

therefore, \$40 satisfies the enunciation.

4. The sum of two numbers is 11, and the sum of their squares is 61: what are the numbers? *Ans.* 5 and 6.

5. The difference of two numbers is 3, and the sum of their squares is 89: what are the numbers? *Ans.* 5 and 8.

6. A grazier bought as many sheep as cost him £60, and after reserving fifteen out of the number, he sold the remainder for £54, and gained 2*s.* a head on those he sold: how many did he buy? *Ans.* 75.

7. A merchant bought cloth, for which he paid £33 15*s.*, which he sold again at £2 8*s.* per piece, and gained by the bargain as much as one piece cost him: how many pieces did he buy? *Ans.* 15.

8. The difference of two numbers is 9, and their sum, multiplied by the greater, is equal to 266: what are the numbers? *Ans.* 14 and 5

9. To find a number, such that if you subtract it from 10, and multiply the remainder by the number itself, the product will be 21. *Ans.* 7 or 3.

10. A person traveled 105 miles. If he had traveled 2 miles an hour slower, he would have been 6 hours longer in completing the same distance: how many miles did he travel per hour? *Ans.* 7 miles.

11. A person purchased a number of sheep, for which he paid $224. Had he paid for each twice as much, plus 2 dollars, the number bought would have been denoted by twice what was paid for each: how many sheep were purchased? *Ans.* 32.

12. The difference of two numbers is 7, and their sum multiplied by the greater, is equal to 130: what are the numbers? *Ans.* 10 and 3.

13. Divide 100 into two such parts, that the sum of their squares shall be 5392. *Ans.* 64 and 36.

14. Two square courts are paved with stones a foot square; the larger court is 12 feet larger than the smaller one, and the number of stones in both pavements is 2120: how long is the smaller pavement? *Ans.* 26 feet.

15. Two hundred and forty dollars are equally distributed among a certain number of persons. The same sum is again distributed amongst a number greater by 4. In the latter case each receives 10 dollars less than in the former: how many persons were there in each case. *Ans.* 8 and 12.

16. Two partners, *A* and *B*, gained 360 dollars. *A's* money was in trade 12 months, and he received, for principal and profit, 520 dollars. *B's* money was 600 dollars, and was in trade 16 months: how much capital had *A*? *Ans.* 400 dollars.

EQUATIONS INVOLVING MORE THAN ONE UNKNOWN QUANTITY.

173. Two simultaneous equations, each of the second degree, and containing two unknown quantities, will, when combined, generally give rise to an equation of the fourth degree. Hence, only particular cases of such equations can be solved by the methods already given.

FIRST.

Two simultaneous equations, involving two unknown quantities, can always be solved when one is of the first and the other of the second degree.

EXAMPLES.

1. Given $\left\{\begin{array}{l} x + y = 14 \\ x^2 + y^2 = 100 \end{array}\right\}$ to find x and y.

By transposing y in the first equation, we have,

$$x = 14 - y;$$

and by squaring both members,

$$x^2 = 196 - 28y + y^2.$$

Substituting this value for x^2 in the second equation, we have,

$$196 - 28y + y^2 + y^2 = 100;$$

from which we have,

$$y^2 - 14y = -48.$$

By completing the square,

$$y^2 - 14y + 49 = 1;$$

---

173. When may two simultaneous equations of the second degree be solved?

and by extracting the square root,

$$y - 7 = \pm\sqrt{1} = +1, \text{ and } -1;$$

hence, $y' = 7 + 1 = 8$, and $y'' = 7 - 1 = 6$.

If we take the greater value, we find $x = 6$; and if we take the lesser, we find $x = 8$.

$$Ans. \begin{cases} x' = 8, & x'' = 6. \\ y' = 6, & y'' = 8 \end{cases}$$

VERIFICATION.

For the greater value, $y = 8$, the equation,

$$x + y = 14, \text{ gives } 6 + 8 = 14;$$

and, $x^2 + y^2 = 100$, gives $36 + 64 = 100$.

For the value $y = 6$, the equation,

$$x + y = 14, \text{ gives } 8 + 6 = 14;$$

and, $x^2 + y^2 = 100$, gives $64 + 36 = 100$.

Hence, both sets of values satisfy the given equation.

2. Given $\left\{ \begin{array}{l} x - y = 3 \\ x^2 - y^2 = 45 \end{array} \right\}$ to find x and y.

Transposing y in the first equation, we have,

$$x = 3 + y;$$

then, squaring both members,

$$x^2 = 9 + 6y + y^2.$$

Substituting this value for x^2, in the second equation, we have,

$$9 + 6y + y^2 - y^2 = 45;$$

whence, we have,

$$6y = 36, \text{ and } y = 6.$$

Substituting this value of y, in the first equation, we have,

$$x - 6 = 3,$$

and, consequently, $x' = 3 + 6 = 9.$

VERIFICATION.

$$x - y = 3, \text{ gives } 9 - 6 = 3;$$

and, $$x^2 - y^2 = 45, \text{ gives } 81 - 36 = 45.$$

Solve the following simultaneous equations:

3. $\left\{\begin{array}{l} x + y = 12 \\ x^2 - y^2 = 24 \end{array}\right\}$ *Ans.* $\left\{\begin{array}{l} x' = 7. \\ y' = 5. \end{array}\right.$

4. $\left\{\begin{array}{l} x - y = 3 \\ x^2 + y^2 = 117 \end{array}\right\}$ *Ans.* $\left\{\begin{array}{ll} x' = 9, & x'' = -6. \\ y' = 6, & y'' = -9. \end{array}\right.$

5. $\left\{\begin{array}{l} x + y = 9 \\ x^2 - 2xy + y^2 = 1 \end{array}\right\}$ *Ans.* $\left\{\begin{array}{ll} x' = 5, & x'' = 5. \\ y' = 4, & y'' = 4. \end{array}\right.$

6. $\left\{\begin{array}{l} x - y = 5 \\ x^2 + 2xy + y^2 = 225 \end{array}\right\}$

Ans. $\left\{\begin{array}{ll} x' = 10, & x'' = -5. \\ y' = 5, & y'' = -10. \end{array}\right.$

SECOND.

174. *Two simultaneous equations of the second degree, which are homogeneous with respect to the unknown quantity, can always be solved.*

EXAMPLES.

1. Given $\left\{\begin{array}{lr} x^2 + 3xy = 22 & \ldots\ldots\ (1.) \\ x^2 + 3xy + 2y^2 = 40 & \ldots\ldots\ (2.) \end{array}\right.$

to find x and y.

---

174. When may two simultaneous equations of the second degree be solved?

Assume $x = ty$, t being any auxiliary unknown quantity.

Substituting this value of x in Equations (1) and (2), we have,

$$t^2y^2 + 3ty^2 = 22, \quad \therefore \quad y^2 = \frac{22}{t^2 + 3t}; \qquad (3.)$$

$$t^2y^2 + 3ty^2 + 2y^2 = 40, \quad \therefore \quad y^2 = \frac{40}{t^2 + 3t + 2}; \qquad (4)$$

hence,
$$\frac{22}{t^2 + 3t} = \frac{40}{t^2 + 3t + 2};$$

hence,
$$22t^2 + 66t + 44 = 40t^2 + 120t;$$

reducing,
$$t^2 + 3t = \frac{22}{9};$$

whence,
$$t' = \frac{2}{3}, \text{ and } t'' = -\frac{11}{3}.$$

Substituting either of these values in Equations (3) or (4), we find,

$$y' = +3, \text{ and } y'' = -3$$

Substituting the plus value of y, in Equation (1), we have,

$$x^2 + 9x = 22;$$

from which we find,

$$x' = +2, \text{ and } x'' = -11.$$

If we take the negative value, $y'' = -3$, we have, from Equation (1),

$$x^2 - 9x = 22;$$

from which we find,

$$x' = +11, \text{ and } x'' = -2.$$

VERIFICATION.

For the values $y' = +3$, and $x' = +2$, the given equation,

$$x^2 + 3xy = 22,$$

gives, $2^2 + 3 \times 2 \times 3 = 4 + 18 = 22;$

and for the second value, $x'' = -11$, the same equation,

$$x^2 + 3xy = 22,$$

gives, $(-11)^2 + 3 \times -11 \times 3 = 121 - 99 = 22.$

If, now, we take the second value of y, that is, $y'' = -3$, and the corresponding values of x, viz., $x' = +11$, and $x'' = -2$; for $x' = +11$, the given equation,

$$x^2 + 3xy = 22,$$

gives, $11^2 + 3 \times 11 \times -3 = 121 - 99 = 22;$

and for $x'' = -2$, the same equation,

$$x^2 + 3xy = 22,$$

gives, $(-2)^2 + 3 \times -2 \times -3 = 4 + 18 = 22.$

The verifications could be made in the same way by employing Equation (2).

NOTE.—In equations similar to the above, we generally find but a single pair of values, corresponding to the values in this equation, of $y' = +3$, and $x' = +2$.

The complete solution would give four pairs of values.

2. $\begin{cases} x^2 - y^2 = -9 \\ y^2 - xy = 5 \end{cases}$ *Ans.* $\begin{cases} x = 4. \\ y = 5. \end{cases}$

3. $\begin{cases} xy - y^2 = -7 \\ y^2 + x^2 = 85 \end{cases}$ *Ans.* $\begin{cases} x = 6. \\ y = 7. \end{cases}$

4. $\begin{cases} 2x^2 + 3xy = 470 \\ y^2 - xy = -9 \end{cases}$ *Ans.* $\begin{cases} x = 10. \\ y = 9 \end{cases}$

5. $\begin{cases} 5xy - 3y^2 = 32 \\ x^2 + y^2 + 3xy = 71 \end{cases}$ *Ans.* $\begin{cases} x = 7. \\ y = 1. \end{cases}$

THIRD.—PARTICULAR CASES.

175 Many other equations of the second degree may be so transformed, as to be brought under the rules of solution already given. The seven following formulas will aid in such transformation.

(**1.**)

When the sum and difference are known:

$$x + y = s$$
$$x - y = d.$$

Then, page 132, Example 8,

$$x = \frac{s+d}{2} = \frac{1}{2}s + \frac{1}{2}d, \quad \text{and} \quad y = \frac{s-d}{2} = \frac{1}{2}s - \frac{1}{2}d.$$

(**2.**)

When the sum and product are known:

$$x + y = s \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad (1.)$$
$$xy = p \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad (2.)$$
$$x^2 + 2xy + y^2 = s^2, \text{ by squaring } (1);$$
$$4xy = 4p, \text{ by mult. } (2) \text{ by } 4.$$
$$x^2 - 2xy + y^2 = s^2 - 4p, \text{ by subtraction.}$$
$$x - y = \pm\sqrt{s^2 - 4p}, \text{ by ext. root.}$$

But, $x + y = s;$

hence $$x = \frac{s}{2} \pm \frac{1}{2}\sqrt{s^2 - 4p}.$$

and, $$y = \frac{s}{2} \mp \frac{1}{2}\sqrt{s^2 - 4p}.$$

---

175. What is the first formula of this article? What the second? Third? Fourth? Fifth? Sixth? Seventh?

(3.)

When the difference and product are known:

$$\begin{aligned}
x - y &= d \;.\;.\;.\;.\;.\;.\;.\;.\;. \quad (1.)\\
xy &= p \;.\;.\;.\;.\;.\;.\;.\;.\;. \quad (2.)\\
x^2 - 2xy + y^2 &= d^2, \text{ by squaring } (1)\cdot\\
4xy &= 4p, \text{ mult. } (2) \text{ by } 4.\\
\hline
x^2 + 2xy + y^2 &= d^2 + 4p, \text{ by adding.}\\
x + y &= \pm\sqrt{d^2 + 4p}\\
x - y &= d\\
\hline
x &= \tfrac{1}{2}d \pm \tfrac{1}{2}\sqrt{d^2 + 4p}.\\
y &= -\tfrac{1}{2}d \pm \tfrac{1}{2}\sqrt{d^2 + 4p}.
\end{aligned}$$

4.)

When the sum of the squares and product are known.

$x^2 + y^2 = s\;.\,.\;(1.)\quad xy = p\;.\,.\;(2.)\quad \therefore 2xy = 2p\;.\,.\;(3.)$

Adding (1) and (3), $x^2 + 2xy + y^2 = s + 2p$;

hence, $$x + y = \pm\sqrt{s + 2p} \quad (4.)$$

Subtracting (3) from (1), $x^2 - 2xy + y^2 = s - 2p$;

hence, $$x - y = \pm\sqrt{s - 2p} \quad (5.)$$

Combining (4) and (5), $x = \tfrac{1}{2}\sqrt{s + 2p} + \tfrac{1}{2}\sqrt{s - 2p}$,

and, $$y = \tfrac{1}{2}\sqrt{s + 2p} - \tfrac{1}{2}\sqrt{s - 2p}.$$

(5.)

When the sum and sum of the squares are known:

$$\begin{aligned}
x + y &= s \;.\;.\;.\;.\;.\;. \quad (1.)\\
x^2 + y^2 &= s' \;.\;.\;.\;.\;.\;. \quad (2.)\\
x^2 + 2xy + y^2 &= s^2 \text{ by squaring } (1)\\
\hline
2xy &= s^2 - s'\\
xy &= \frac{s^2 - s'}{2} = p. \quad (3.)
\end{aligned}$$

By putting $xy = p$, and combining Equations (1) and (3), by Formula (2), we find the values of x and y.

(6.)

When the sum and sum of the cubes are known:

$$\begin{array}{rcll}
x + y & = & 8 & \quad .\ .\ .\ .\ \ (1.) \\
x^3 + y^3 & = & 152 & \quad .\ .\ .\ .\ \ (2.) \\
\underline{x^3 + 3x^2y + 3xy^2 + y^3} & = & \underline{512} & \quad \text{by cubing (1).} \\
3x^2y + 3xy^2 & = & 360 & \quad \text{by subtraction.} \\
3xy(x + y) & = & 360 & \quad \text{by factoring.} \\
3xy(8) & = & 360 & \quad \text{from Equa. (1)} \\
24xy & = & 360 & \\
\text{hence,} \qquad xy & = & 15 & \quad .\ .\ .\ .\ \ (3.)
\end{array}$$

Combining (1) and (3), we find $x = 5$ and $y = 3$

(7.)

When we have an equation of the form,

$$(x + y)^2 + (x + y) = q.$$

Let us assume $x + y = z$.

Then the given equation becomes,

$$z^2 + z = q; \quad \text{and} \quad z = -\frac{1}{2} \pm \sqrt{q + \frac{1}{4}}.$$

$$x + y = -\frac{1}{2} \pm \sqrt{q + \frac{1}{4}}.$$

EXAMPLES.

1. Given $\left\{\begin{array}{rcll} xz & = & y^2 & (1) \\ x + y + z & = & 7 & (2) \\ x^2 + y^2 + z^2 & = & 21 & (3) \end{array}\right\}$ to find x, y, and z.

Transposing y in Equation (2), we have,

$$x + z = 7 - y; \quad . \; . \; . \quad (4.)$$

then, squaring the members, we have,

$$x^2 + 2xz + z^2 = 49 - 14y + y^2.$$

If now we substitute for $2xz$, its value taken from Equation (1), we have,

$$x^2 + 2y^2 + z^2 = 49 - 14y + y^2;$$

and cancelling y^2, in each member, there results,

$$x^2 + y^2 + z^2 = 49 - 14y.$$

But, from Equation (3), we see that each member of the last equation is equal to 21; hence,

$$49 - 14y = 21,$$

and, $\quad 14y = 49 - 21 = 28,$

hence, $\quad y = \frac{28}{14} = 2.$

Substituting this value of y in Equation (1), gives,

$$xz = 4;$$

and substituting it in Equation (4), gives,

$$x + z = 5, \quad \text{or} \quad x = 5 - z.$$

Substituting this value of x, in the previous equation, we obtain

$$5z - z^2 = 4, \quad \text{or} \quad z^2 - 5z = -4;$$

and by completing the square, we have,

$$z^2 - 5z + 6.25 = 2.25.$$

and, $\quad z - 2.5 = \pm\sqrt{2.25} = +1.5, \quad \text{or} \quad -1.5$

hence, $z = 2.5 + 1.5 = 4$, and $z = +2.5 - 1.5 = 1$

2. $$\left.\begin{array}{rl}\text{Given} & x + \sqrt{xy} + y = 19 \\ \text{and} & x^2 + xy + y^2 = 133\end{array}\right\}\text{ to find } x \text{ and } y.$$

Dividing the second equation by the first, we have,

$$\begin{array}{lr}
 & x - \sqrt{xy} + y = 7 \\
\text{but,} & x + \sqrt{xy} + y = 19 \\ \hline
\text{hence, by addition,} & 2x + 2y = 26 \\
\text{or,} & x + y = 13 \\
\text{and substituting in 1st Equa.,} & \sqrt{xy} + 13 = 19 \\
\text{or, by transposing,} & \sqrt{xy} = 6 \\
\text{and by squaring,} & xy = 36.
\end{array}$$

$$\begin{array}{lr}
\text{Equation 2d, is} & x^2 + xy + y^2 = 133 \\
\text{and from the last, we have,} & 3xy = 108 \\ \hline
\text{Subtracting,} & x^2 - 2xy + y^2 = 25 \\
\text{hence,} & x - y = \pm 5 \\
\text{but,} & x + y = 13
\end{array}$$

hence, $x = 9$ and 4; and $y = 4$ and 9.

PROBLEMS.

1. Find two numbers, such that their sum shall be 15 and the sum of their squares 113.

Let x and y denote the numbers; then,

$$x + y = 15, \quad (1.) \qquad \text{and} \qquad x^2 + y^2 = 113. \quad (2.)$$

From Equation (1), we have,

$$x^2 = 225 - 30y + y^2$$

Substituting this value in Equation (2),

$$225 - 30y + y^2 + y^2 = 113;$$

hence, $$2y^2 - 30y = -112,$$

$$y^2 - 15y = -\ 56,$$

hence, $$y' = 8, \quad \text{and} \quad y'' = 7.$$

The first value of y being substituted in Equation (1), gives $x' = 7$; and the second, $x'' = 8$. Hence, the numbers are 7 and 8.

2. To find two numbers, such that their product added to their sum shall be 17, and their sum taken from the sum of their squares shall leave 22.

Let x and y denote the numbers; then, from the conditions,

$$(x + y) + xy = 17. \quad . \ . \ . \quad (1.)$$

$$x^2 + y^2 - (x + y) = 22. \quad . \ . \ . \quad (2.)$$

Multiplying Equation (1) by 2, we have,

$$2xy + 2(x + y) = 34. \quad . \ . \ . \quad (3.)$$

Adding (2) and (3), we have,

$$x^2 + 2xy + y^2 + (x + y) = 56;$$

hence, $$(x + y)^2 + (x + y) = 56. \quad . \ . \quad (4.)$$

Regarding $(x + y)$ as a single unknown quantity (page 248),

$$x + y = -\frac{1}{2} \pm \sqrt{56 + \frac{1}{4}} = 7.$$

Substituting this value in Equation (1), we have,

$$7 + xy = 17, \quad \text{and} \quad y = 5.$$

Hence, the numbers are 2 and 5.

3. What two numbers are those whose sum is 8, and sum of their squares 34? *Ans.* 5 and 3.

4. It is required to find two such numbers, that the first shall be to the second as the second is to 16, and the sum of whose squares shall be 225? *Ans.* 9 and 12.

5. What two numbers are those which are to each other as 3 to 5, and whose squares added together make 1666? *Ans.* 21 and 35.

6. There are two numbers whose difference is 7, and half their product plus 30 is equal to the square of the less number: what are the numbers? *Ans.* 12 and 19.

7. What two numbers are those whose sum is 5, and the sum of their cubes 35? *Ans.* 2 and 3.

8. What two numbers are those whose sum is to the greater as 11 to 7, and the difference of whose squares is 132? *Ans.* 14 and 8.

9. Divide the number 100 into two such parts, that the product may be to the sum of their squares as 6 to 13. *Ans.* 40 and 60

10. Two persons, *A* and *B*, departed from different places at the same time, and traveled towards each other. On meeting, it appeared that *A* had traveled 18 miles more than *B*; and that *A* could have gone *B's* journey in $15\frac{3}{4}$ days, but *B* would have been 28 days in performing *A's* journey: how far did each travel? *Ans.* { *A*, 72 miles. *B*, 54 miles.

11. There are two numbers whose difference is 15, and half their product is equal to the cube of the lesser number: what are those numbers? *Ans.* 3 and 18.

12. What two numbers are those whose sum, multiplied by the greater, is equal to 77; and whose difference, multiplied by the less, is equal to 12?

Ans. 4 and 7, or $\frac{3}{2}\sqrt{2}$ and $\frac{11}{2}\sqrt{2}$.

13. Divide 100 into two such parts, that the sum of their square roots may be 14. *Ans.* 64 and 36.

14. It is required to divide the number 24 into two such parts, that their product may be equal to 35 times their difference. *Ans.* 10 and 14.

15. The sum of two numbers is 8, and the sum of their cubes is 152: what are the numbers? *Ans.* 3 and 5.

16. Two merchants each sold the same kind of stuff; the second sold 3 yards more of it than the first, and together they receive 35 dollars. The first said to the second, "I would have received 24 dollars for your stuff;" the other replied, "And I should have received $12\frac{1}{2}$ dollars for yours:" how many yards did each of them sell?

$$\text{Ans.}\begin{cases}\text{1st merchant } x' = 15, & \text{or, } x'' = 5.\\ \text{2d} \quad \text{“} \quad y' = 18, & \quad\;\; y'' = 8.\end{cases}$$

17. A widow possessed 13,000 dollars, which she divided into two parts, and placed them at interest in such a manner that the incomes from them were equal. If she had put out the first portion at the same rate as the second, she would have drawn for this part 360 dollars interest; and if she had placed the second out at the same rate as the first, she would have drawn for it 490 dollars interest: what were the two rates of interest? *Ans.* 7 and 6 per cent.

18. Find three numbers, such that the difference between the third and second shall exceed the difference between the second and first by 6; that the sum of the numbers shall be 33, and the sum of their squares 467.

Ans. 5, 9, and 19.

19. What number is that which, being divided by the product of its two digits, the quotient will be 3; and if 18 be added to it, the resulting number will be expressed by the digits inverted? *Ans.* 24.

20. What two numbers are those which are to each other as m to n, and the sum of whose squares is b?

Ans. $\dfrac{m\sqrt{b}}{\sqrt{m^2+n^2}}, \quad \dfrac{n\sqrt{b}}{\sqrt{m^2+n^2}}.$

21. What two numbers are those which are to each other as m to n, and the difference of whose squares is b?

Ans. $\dfrac{m\sqrt{b}}{\sqrt{m^2-n^2}}, \quad \dfrac{n\sqrt{b}}{\sqrt{m^2-n^2}}.$

22. Required to find three numbers, such that the product of the first and second shall be equal to 2; the product of the first and third equal to 4, and the sum of the squares of the second and third equal to 20. *Ans.* 1, 2, and 4.

23. It is required to find three numbers, whose sum shall be 38, the sum of their squares 634, and the difference between the second and first greater by 7 than the difference between the third and second. *Ans.* 3, 15, and 20.

24. Required to find three numbers, such that the product of the first and second shall be equal to a; the product of the first and third equal to b; and the sum of the squares of the second and third equal to c.

$$\text{Ans.} \begin{cases} x = \sqrt{\dfrac{a^2+b^2}{c}}. \\ y = a\sqrt{\dfrac{c}{a^2+b^2}}. \\ z = b\sqrt{\dfrac{c}{a^2+b^2}}. \end{cases}$$

25. What two numbers are those, whose sum, multiplied by the greater, gives 144; and whose difference, multiplied by the less, gives 14? *Ans.* 9 and 7.

CHAPTER IX.

OF PROPORTIONS AND PROGRESSIONS.

176. Two quantities of the same kind may be compared, the one with the other, in two ways:

1st. By considering *how much* one is greater or less than the other, which is shown by their difference; and,

2d. By considering *how many times* one is greater or less than the other, which is shown by their quotient.

Thus, in comparing the numbers 3 and 12 together, with respect to their difference, we find that 12 *exceeds* 3, by 9; and in comparing them together with respect to their quotient, we find that 12 *contains* 3, four times, or that 12 is 4 times as great as 3.

The first of these methods of comparison is called *Arithmetical Proportion*, and the second, *Geometrical Proportion.*

Hence, *Arithmetical Proportion considers the relation of quantities with respect to their difference, and Geometrical Proportion the relation of quantities with respect to their quotient.*

---

176. In how many ways may two quantities be compared the one with the other? What does the first method consider? What the second? What is the first of these methods called? What is the second called? How then do you define the two proportions?

OF ARITHMETICAL PROPORTION AND PROGRESSION.

177. If we have four numbers, 2, 4, 8, and 10, of which the difference between the first and second is equal to the difference between the third and fourth, these numbers are said to be in arithmetical proportion. The first term 2 is called an *antecedent*, and the second term 4, with which it is compared, a *consequent*. The number 8 is also called an antecedent, and the number 10, with which it is compared, a consequent.

When the difference between the first and second is equal to the difference between the third and fourth, the four numbers are said to be in proportion. Thus, the numbers,

$$2, \quad 4, \quad 8, \quad 10,$$

are in arithmetical proportion.

178. When the difference between the first antecedent and consequent is the same as between any two consecutive terms of the proportion, the proportion is called an *arithmetical progression*. Hence, a *progression by differences*, or an *arithmetical progression*, is a series in which the successive terms are continually increased or decreased by a constant number, which is called the *common difference* of the progression.

Thus, in the two series,

$$1, \quad 4, \quad 7, \quad 10, \quad 13, \quad 16, \quad 19, \quad 22, \quad 25, \ldots$$
$$60, \quad 56, \quad 52, \quad 48, \quad 44, \quad 40, \quad 36, \quad 32, \quad 28, \ldots$$

---

177. When are four numbers in arithmetical proportion? What is the first called? What is the second called? What is the third called? What is the fourth called?

178. What is an arithmetical progression? What is the number called by which the terms are increased or diminished? What is an increasing progression? What is a decreasing progression? Which term is only an antecedent? Which only a consequent?

the first is called an *increasing progression*, of which the common difference is 3, and the second, a *decreasing progression*, of which the common difference is 4.

In general, let a, b, c, d, e, f, ... denote the terms of a progression by differences; it has been agreed to write them thus:

$$a \,.\, b \,.\, c \,.\, d \,.\, e \,.\, f \,.\, g \,.\, h \,.\, i \,.\, k \ldots$$

This series is read, a is to b, as b is to c, as c is to d, as d is to e, &c. This is a series of *continued equi-differences*, in which each term is at the same time an antecedent and a consequent, with the exception of the first term, which is only an *antecedent*, and the last, which is only a *consequent*.

179. Let d denote the common difference of the progresion,

$$a \,.\, b \,.\, c \,.\, e \,.\, f \,.\, g \,.\, h. \text{ \&c.},$$

which we will consider increasing.

From the definition of the progression, it evidently follows that,

$$b = a + d, \quad c = b + d = a + 2d, \quad e = c + d = a + 3d;$$

and, in general, any term of the series is equal to *the first term, plus as many times the common difference as there are preceding terms.*

Thus, let l be any term, and n the number which marks the place of it; the expression for this *general term* is,

$$l = a + (n - 1)d.$$

Hence, for finding the last term, we have the following

---

179. Give the rule for finding the last term of a series when the progression is increasing.

RULE.

I. *Multiply the common difference by the number of terms less one:*

II. *To the product add the first term; the sum will be the last term.*

EXAMPLES.

The formula,

$$l = a + (n - 1)d,$$

serves to find any term whatever, without determining all those which precede it.

1. If we make $n = 1$, we have, $l = a$; that is, the series will have but one term.

2. If we make $n = 2$, we have, $l = a + d$; that is, the series will have two terms, and the second term is equal to the first, plus the common difference.

3. If $a = 3$, and $d = 2$, what is the 3d term? *Ans.* 7.

4. If $a = 5$, and $d = 4$, what is the 6th term? *Ans.* 25.

5. If $a = 7$, and $d = 5$, what is the 9th term? *Ans.* 47.

6. If $a = 8$, and $d = 5$, what is the 10th term? *Ans.* 53

7. If $a = 20$, and $d = 4$, what is the 12th term? *Ans.* 64.

8. If $a = 40$, and $d = 20$, what is the 50th term? *Ans.* 1020.

9. If $a = 45$, and $d = 30$, what is the 40th term? *Ans.* 1215.

10. If $a = 30$, and $d = 20$, what is the 60th term?
Ans. 1210.

11. If $a = 50$, and $d = 10$, what is the 100th term?
Ans. 1040.

12. To find the 50th term of the progression,

$$1 . 4 . 7 . 10 . 13 . 16 . 19 \ldots$$

we have, $l = 1 + 49 \times 3 = 148$.

13. To find the 60th term of the progression,

$$1 . 5 . 9 . 13 . 17 . 21 . 25 \ldots$$

we have, $l = 1 + 59 \times 4 = 237$.

180. If the progression were a decreasing one, we should have,

$$l = a - (n - 1)d.$$

Hence, to find the last term of a decreasing progression, we have the following

RULE.

I. *Multiply the common difference by the number of terms less one:*

II. *Subtract the product from the first term; the remainder will be the last term.*

EXAMPLES.

1. The first term of a decreasing progression is 60, the number of terms 20, and the common difference 3: what is the last term?

$l = a - (n - 1)d$, gives $l = 60 - (20 - 1)3 = 60 - 57 = 3$.

---

180. Give the rule for finding the last term of a series, when the progression is decreasing.

2. The first term is 90, the common difference 4, and the number of terms 15: what is the last term? *Ans.* 34.

3. The first term is 100, the number of terms 40, and the common difference 2: what is the last term? *Ans.* 22.

4. The first term is 80, the number of terms 10, and the common difference 4: what is the last term? *Ans.* 44.

5. The first term is 600, the number of terms 100, and the common difference 5: what is the last term?
Ans. 105.

6. The first term is 800, the number of terms 200, and the common difference 2: what is the last term?
Ans. 402.

181. A progression by differences being given, it is proposed to prove that, *the sum of any two terms, taken at equal distances from the two extremes, is equal to the sum of the two extremes.*

That is, if we have the progression,

$$2 \,.\, 4 \,.\, 6 \,.\, 8 \,.\, 10 \,.\, 12,$$

we wish to prove generally, that,

$$4 + 10, \quad \text{or} \quad 6 + 8,$$

is equal to the sum of the two extremes, 2 and 12.

Let $a \,.\, b \,.\, c \,.\, e \,.\, f \,\ldots\, i \,.\, k \,.\, l$, be the proposed progression, and n the number of terms.

We will first observe that, if x denotes a term which has p terms before it, and y a term which has p terms after it, we have, from what has been said,

---

181. In every progression by differences, what is the sum of the two extremes equal to? What is the rule for finding the sum of an arithmetical series?

$$x = a + p \times d,$$

and, $$y = l - p \times d;$$

whence, by addition, $x + y = a + l$,

which proves the proposition.

Referring to the previous example, if we suppose, in the first place, x to denote the second term 4, then y will denote the term 10, next to the last. If x denotes the third term 6, then y will denote 8, the third term from the last.

To apply this principle in finding the sum of the terms of a progression, write the terms, as below, and then again, in an inverse order, viz.:

$$a \,.\, b \,.\, c \,.\, d \,.\, e \,.\, f \,.\,.\,.\, i \,.\, k \,.\, l.$$

$$l \,.\, k \,.\, i \,.\,.\,.\,.\,.\,.\,.\,.\,.\,.\, c \,.\, b \,.\, a.$$

Calling S the sum of the terms of the first progression, $2S$ will be the sum of the terms of both progressions, and we shall have,

$$2S = (a+l) + (b+k) + (c+i) \ldots + (i+c) + (k+b) + (l+a).$$

Now, since all the parts, $a + l$, $b + k$, $c + i \ldots$ are equal to each other, and their number equal to n,

$$2S = (a + l) \times n, \quad \text{or} \quad S = \left(\frac{a + l}{2}\right) \times n.$$

Hence, for finding the sum of an arithmetical series, we have the following

RULE.

I. *Add the two extremes together, and take half their sum:*

II. *Multiply this half-sum by the number of terms; the product will be the sum of the series.*

EXAMPLES.

1. The extremes are 2 and 16, and the number of terms 8: what is the sum of the series?

$$S = \left(\frac{a+l}{2}\right) \times n, \quad \text{gives} \quad S = \frac{2+16}{2} \times 8 = 72.$$

2. The extremes are 3 and 27, and the number of terms 12: what is the sum of the series? *Ans.* 180.

3. The extremes are 4 and 20, and the number of terms 10: what is the sum of the series? *Ans.* 120.

4. The extremes are 100 and 200, and the number of terms 80: what is the sum of the series? *Ans.* 12000.

5. The extremes are 500 and 60, and the number of terms 20: what is the sum of the series? *Ans.* 5600

6. The extremes are 800 and 1200, and the number of terms 50: what is the sum of the series? *Ans.* 50000.

182. In arithmetical proportion there are five members to be considered:

1st. The first term, a.
2d. The common difference, d.
3d. The number of terms, n.
4th. The last term, l.
5th. The sum, S.

The formulas,

$$l = a + (n-1)d, \quad \text{and} \quad S = \left(\frac{a+l}{2}\right) \times n,$$

contain five quantities, a, d, n, l, and S, and consequently give rise to the following general problem, viz.: *Any three*

---

182. How many numbers are considered in arithmetical proportion? What are they? In every arithmetical progression, what is the common difference equal to?

of these five quantities being given, to determine the other two.

We already know the value of S in terms of a, n, and l. From the formula,

$$l = a + (n - 1)d,$$

we find, $$a = l - (n - 1)d.$$

That is: *The first term of an increasing arithmetical progression is equal to the last term, minus the product of the common difference by the number of terms less one.*

From the same formula, we also find,

$$d = \frac{l - a}{n - 1}.$$

That is: *In any arithmetical progression, the common difference is equal to the last term, minus the first term, divided by the number of terms less one.*

The last term is 16, the first term 4, and the number of terms 5: what is the common difference?

The formula, $$d = \frac{l - a}{n - 1}$$

gives, $$d = \frac{16 - 4}{4} = 3.$$

2. The last term is 22, the first term 4, and the number of terms 10: what is the common difference? *Ans.* 2.

183. The last principle affords a solution to the following question:

To find a number m *of arithmetical means between two given numbers* a *and* b.

---

183. How do you find any number of arithmetical means between two given numbers?

To resolve this question, it is first necessary to find the common difference. Now, we may regard a as the first term of an arithmetical progression, b as the last term, and the required means as intermediate terms. The number of terms of this progression will be expressed by $m + 2$.

Now, by substituting in the above formula, b for l, and $m + 2$ for n, it becomes,

$$d = \frac{b - a}{m + 2 - 1} = \frac{b - a}{m + 1};$$

that is: *The common difference of the required progression is obtained by dividing the difference between the given numbers*, a *and* b, *by the required number of means plus one.*

Having obtained the common difference, d, form the second term of the progression, or the *first arithmetical mean*, by adding d to the first term a. The *second mean* is obtained by augmenting the first mean by d, &c.

1. Find three arithmetical means between the extremes 2 and 18.

The formula, $$d = \frac{b - a}{m + 1},$$

gives, $$d = \frac{18 - 2}{4} = 4;$$

hence, the progression is,

$$2 . 6 . 10 . 14 . 18.$$

2. Find twelve arithmetical means between 12 and 77.

The formula, $$d = \frac{b - a}{m + 1},$$

gives, $$d = \frac{77 - 12}{13} = 5;$$

hence, the progression is,

$$12 . 17 . 22 . 27 \quad . . . 77.$$

184. REMARK.—If the same number of arithmetical means are inserted between all the terms, taken two and two, these terms, and the arithmetical means united, will form one and the same progression.

For, let $a \,.\, b \,.\, c \,.\, e \,.\, f \ldots$ be the proposed progression, and m the number of means to be inserted between a and b, b and c, c and $e \ldots$ &c.

From what has just been said, the common difference of each partial progression will be expressed by

$$\frac{b-a}{m+1}, \quad \frac{c-b}{m+1}, \quad \frac{e-c}{m+1} \ldots$$

expressions which are equal to each other, since a, b, $c \ldots$ are in progression; therefore, the common difference is the same in each of the partial progressions; and, since the *last term* of the first forms the *first term* of the second, &c., we may conclude, that all of these partial progressions form a single progression.

EXAMPLES.

1. Find the sum of the first fifty terms of the progression 2 . 9 . 16 . 23 . . .

For the 50th term, we have,

$$l = 2 + 49 \times 7 = 345.$$

Hence, $S = (2 + 345) \times \dfrac{50}{2} = 347 \times 25 = 8675.$

2. Find the 100th term of the series 2 . 9 . 16 . 23 . . .
Ans. 695

3. Find the sum of 100 terms of the series 1 . 3 . 5 . 7 . 9 . . . .
Ans 10000.

4. The greatest term is 70, the common difference 3, and the number of terms 21: what is the least term and the sum of the series?

Ans. Least term, 10; sum of series, 840.

5. The first term is 4, the common difference 8, and the number of terms 8: what is the last term, and the sum of the series? *Ans.* $\begin{cases}\text{Last term, } 60\\ \text{Sum } = 256.\end{cases}$

6. The first term is 2, the last term 20, and the number of terms 10: what is the common difference? *Ans.* 2.

7. Insert four means between the two numbers 4 and 19: what is the series? *Ans.* 4 . 7 . 10 . 13 . 16 . 19.

8. The first term of a decreasing arithmetical progression is 10, the common difference one-third, and the number of terms 21: required the sum of the series. *Ans.* 140.

9. In a progression by differences, having given the common difference 6, the last term 185, and the sum of the terms 2945: find the first term, and the number of terms.

Ans. First term = 5; number of terms, 31.

10. Find nine arithmetical means between each antecedent and consequent of the progression 2 . 5 . 8 . 11 . 14 . . .

Ans. Common diff., or $d = 0.3$.

11. Find the number of men contained in a triangular battalion, the first rank containing one man, the second 2, the third 3, and so on to the n^{th}, which contains n. In other words, find the expression for the sum of the natural numbers 1, 2, 3 . . ., from 1 to n inclusively.

Ans. $S = \frac{n(n+1)}{2}$.

12. Find the sum of the n first terms of the progression of uneven numbers, 1 . 3 . 5 . 7 . 9, . . . *Ans.* $S = n^2$.

13. One hundred stones being placed on the ground in a straight line, at the distance of 2 yards apart, how far will a person travel who shall bring them one by one to a basket, placed at a distance of 2 yards from the first stone?

Ans. 11 miles, 840 yards.

GEOMETRICAL PROPORTION AND PROGRESSION.

185. *Ratio* is the quotient arising from dividing one quantity by another quantity of the same kind, regarded as a standard. Thus, if the numbers 3 and 6 have the same unit, the ratio of 3 to 6 will be expressed by

$$\frac{6}{3} = 2.$$

And in general, if A and B represent quantities of the same kind, the ratio of A to B will be expressed by

$$\frac{B}{A}.$$

186. The character $\propto$ indicates that one quantity is proportional to another. Thus,

$$A \propto B,$$

is read, A proportional to B.

If there be four numbers,

$$2, \quad 4, \quad 8, \quad 16,$$

having such values that the second divided by the first is equal to the fourth divided by the third, the numbers are

---

185. What is ratio? What is the ratio of 3 to 6? Of 4 to 12?

186. What is proportion? How do you express that four numbers are in proportion? What are the numbers called? What are the first and fourth terms called? What the second and third?

said to form a proportion. And in general, if there be four quantities, A, B, C, and D, having such values that,

$$\frac{B}{A} = \frac{D}{C},$$

then, A is said to have the same ratio to B that C has to D; or, the ratio of A to B is equal to the ratio of C to D. When four quantities have this relation to each other, compared together two and two, they are said to form a geometrical proportion.

To express that the ratio of A to B is equal to the ratio of C to D, we write the quantities thus,

$$A : B :: C : D;$$

and read, A is to B as C to D.

The quantities which are compared, the one with the other, are called *terms* of the proportion. The first and last terms are called the *two extremes*, and the second and third terms, the *two means*. Thus, A and D are the extremes, and B and C the means.

187. Of four terms of a proportion, the first and third are called the *antecedents*, and the second and fourth the *consequents*; and the last is said to be a fourth proportional to the other three, taken in order. Thus, in the last proportion A and C are the antecedents, and B and D the consequents.

188. Three quantities are in proportion, when the first has the same ratio to the second that the second has to the

---

187. In four proportional quantities, what are the first and third called? What the second and fourth?

188. When are three quantities proportional? What is the middle one called?

third; and then the middle term is said to be a mean proportional between the other two. For example,

3 : 6 : : 6 : 12;

and 6 is a mean proportional between 3 and 12.

189. Four quantities are said to be in proportion by *inversion*, or *inversely*, when the consequents are made the antecedents, and the antecedents the consequents.

Thus, if we have the proportion,

3 : 6 : : 8 · 16,

the inverse proportion would be,

6 : 3 : : 16 . 8.

190. Quantities are said to be in proportion by *alternation*, or *alternately*, when antecedent is compared with antecedent, and consequent with consequent.

Thus, if we have the proportion,

3 : 6 : : 8 : 16,

the alternate proportion would be,

3 : 8 : : 6 : 16.

191. Quantities are said to be in proportion by *composition*, when the sum of the antecedent and consequent is compared either with antecedent or consequent

Thus, if we have the proportion,

2 : 4 : : 8 : 16,

---

189. When are quantities said to be in proportion by inversion, or inversely?

190. When are quantities in proportion by alternation?

191. When are quantities in proportion by composition?

the proportion by composition would be,

$$2+4 : 2 :: 8+16 : 8;$$

and, $$2+4 : 4 :: 8+16 : 16.$$

192. Quantities are said to be in proportion by *division*, when the difference of the antecedent and consequent is compared either with antecedent or consequent.

Thus, if we have the proportion,

$$3 : 9 :: 12 : 36,$$

the proportion by division will be,

$$9-3 : 3 :: 36-12 : 12;$$

and, $$9-3 : 9 :: 36-12 : 36;$$

193. Equi-multiples of two or more quantities are the products which arise from multiplying the quantities by the same number.

Thus, if we have any two numbers, as 6 and 5, and multiply them both by any number, as 9, the equi-multiples will be 54 and 45; for,

$$6 \times 9 = 54, \text{ and } 5 \times 9 = 45.$$

Also, $m \times A$, and $m \times B$, are equi-multiples of A and B, the common multiplier being m.

194. Two quantities A and B, which may change their values, *are reciprocally or inversely proportional, when one is proportional to unity divided by the other, and then their product remains constant.*

---

192. When are quantities in proportion by division?

193. What are equi-multiples of two or more quantities?

194. When are two quantities said to be reciprocally proportional?

We express this reciprocal or inverse relation thus,

$$A \propto \frac{1}{B}.$$

in which A is said to be inversely proportional to B.

195. If we have the proportion,

$$A : B :: C : D,$$

we have, $$\frac{B}{A} = \frac{D}{C}, \text{ (Art. 186)};$$

and by clearing the equation of fractions, we have,

$$BC = AD.$$

That is: *Of four proportional quantities, the product of the two extremes is equal to the product of the two means.*

This general principle is apparent in the proportion between the numbers,

$$2 : 10 :: 12 : 60,$$

which gives, $$2 \times 60 = 10 \times 12 = 120.$$

196. If four quantities, A, B, C, D, are so related to each other, that

$$A \times D = B \times C,$$

we shall also have, $$\frac{B}{A} = \frac{D}{C};$$

and hence, $$A : B :: C : D.$$

That is: *If the product of two quantities is equal to the product of two other quantities, two of them may be made the extremes, and the other two the means of a proportion.*

---

195. If four quantities are proportional, what is the product of the two means equal to?

196. If the product of two quantities is equal to the product of two other quantities, may the four be placed in a proportion? How?

Thus, if we have,

$$2 \times 8 = 4 \times 4,$$

we also have,

$$2 : 4 :: 4 : 8.$$

197. If we have three proportional quantities,

$$A : B :: B : C,$$

we have,

$$\frac{B}{A} = \frac{C}{B};$$

hence,

$$B^2 = AC.$$

That is: *If three quantities are proportional, the square of the middle term is equal to the product of the two extremes*

Thus, if we have the proportion,

$$3 : 6 :: 6 : 12,$$

we shall also have,

$$6 \times 6 = 6^2 = 3 \times 12 = 36.$$

198. If we have,

$$A : B :: C : D, \text{ and consequently, } \frac{B}{A} = \frac{D}{C},$$

multiply both members of the last equation by $\frac{C}{B}$, and we then obtain,

$$\frac{C}{A} = \frac{D}{B};$$

and, hence,

$$A : C :: B : D.$$

That is: *If four quantities are proportional, they will be in proportion by alternation.*

---

197. If three quantities are proportional, what is the product of the extremes equal to?

198. If four quantities are proportional, will they be in proportion by alternation?

Let us take, as an example,

$$10 : 15 :: 20 \quad 30.$$

We shall have, by alternating the terms,

$$10 : 20 :: 15 : 30.$$

199. If we have,

$$A : B :: C : D, \text{ and } A : B :: E : F,$$

we shall also have,

$$\frac{B}{A} = \frac{D}{C}, \text{ and } \frac{B}{A} = \frac{F}{E};$$

hence, $\frac{D}{C} = \frac{F}{E}$, and $C : D :: E : F$.

That is: *If there are two sets of proportions having an antecedent and consequent in the one, equal to an antecedent and consequent of the other, the remaining terms will be proportional.*

If we have the two proportions,

$$2 : 6 :: 8 : 24, \text{ and } 2 : 6 :: 10 : 30,$$

we shall also have,

$$8 : 24 :: 10 : 30.$$

200. If we have,

$$A : B :: C : D, \text{ and consequently, } \frac{B}{A} = \frac{D}{C},$$

we have, by dividing 1 by each member of the equation,

$$\frac{A}{B} = \frac{C}{D}, \text{ and consequently, } B : A :: D : C.$$

---

199. If you have two sets of proportions having an antecedent and consequent in each, equal; what will follow?

200. If four quantities are in proportion, will they be in proportion when taken inversely?

That is: *Four proportional quantities will be in proportion, when taken inversely.*

To give an example in numbers, take the proportion,

$$7 : 14 :: 8 : 16;$$

then, the inverse proportion will be,

$$14 : 7 :: 16 : 8,$$

in which the ratio is one-half.

201. The proportion,

$$A : B :: C : D, \text{ gives, } A \times D = B \times C.$$

To each member of the last equation add $B \times D$. We shall then have,

$$(A + B) \times D = (C + D) \times B;$$

and by separating the factors, we obtain,

$$A + B : B :: C + D : D.$$

If, instead of adding, we subtract $B \times D$ from both members, we have,

$$(A - B) \times D = (C - D) \times B;$$

which gives,

$$A - B : B :: C - D : D.$$

That is: *If four quantities are proportional, they will be in proportion by composition or division.*

Thus, if we have the proportion,

$$9 : 27 :: 16 : 48,$$

---

201. If four quantities are in proportion, will they be in proportion by composition? Will they be in proportion by division? What is the difference between composition and division?

we shall have, by composition,

$$9 + 27 \cdot 27 :: 16 + 48 : 48;$$

that is, $$36 : 27 :: 64 : 48,$$

in which the ratio is three-fourths.

The same proportion gives us, by division,

$$27 - 9 : 27 :: 48 - 16 : 48;$$

that is, $$18 : 27 :: 32 : 48,$$

in which the ratio is one and one-half.

202. If we have,

$$\frac{B}{A} = \frac{D}{C},$$

and multiply the numerator and denominator of the first member by any number m, we obtain,

$$\frac{mB}{mA} = \frac{D}{C}, \text{ and } mA : mB :: C : D.$$

That is: *Equal multiples of two quantities have the same ratio as the quantities themselves.*

For example, if we have the proportion,

$$5 : 10 :: 12 : 24,$$

and multiply the first antecedent and consequent by 6, we have,

$$30 : 60 :: 12 : 24,$$

in which the ratio is still 2.

203. The proportions,

$$A : B :: C : D, \text{ and } A : B :: E : F,$$

---

202. Have equal multiples of two quantities the same ratio as the quantities?

203. Suppose the antecedent and consequent be augmented or diminished by quantities having the same ratio?

give, $A \times D = B \times C$, and $A \times F = B \times E$;

adding and subtracting these equations, we obtain,

$A(D \pm F) = B(C \pm E)$, or $A : B :: C \pm E : D \pm F.$

That is: *If C and D, the antecedent and consequent, be augmented or diminished by quantities E and F, which have the same ratio as C to D, the resulting quantities will also have the same ratio.*

Let us take, as an example, the proportion,

$$9 : 18 :: 20 : 40,$$

in which the ratio is 2.

If we augment the antecedent and consequent by the numbers 15 and 30, which have the same ratio, we shall have,

$$9 + 15 : 18 + 30 :: 20 : 40;$$

that is, $24 : 48 :: 20 : 40,$

in which the ratio is still 2.

If we diminish the second antecedent and consequent by these numbers respectively, we have,

$$9 : 18 :: 20 - 15 : 40 - 30;$$

that is, $9 : 18 :: 5 : 10,$

in which the ratio is till 2.

204. If we have several proportions,

$A : B :: C : D$, which gives $A \times D = B \times C$,
$A : B :: E : F$, which gives $A \times F = B \times E$,
$A : B :: G : H$, which gives $A \times H = B \times G$,
&c., &c.,

---

204. In any number of proportions having the same ratio, how will any one antecedent be to its consequent?

we shall have, by addition,

$$A(D + F + H) = B(C + E + G);$$

and by separating the factors,

$$A : B :: C + E + G : D + F + H.$$

That is: *In any number of proportions having the same ratio, any antecedent will be to its consequent as the sum of the antecedents to the sum of the consequents.*

Let us take, for example,

$$2 : 4 :: 6 : 12, \quad \text{and} \quad 1 : 2 :: 3 : 6, \quad \text{\&c.}$$

Then $2 : 4 :: 6 + 3 : 12 + 6$;

that is, $2 : 4 :: 9 : 18$,

in which the ratio is still 2.

205. If we have four proportional quantities,

$$A : B :: C : D, \quad \text{we have,} \quad \frac{B}{A} = \frac{D}{C};$$

and raising both members to any power whose exponent is n, or extracting any root whose index is n, we have,

$$\frac{B^n}{A^n} = \frac{D^n}{C^n}, \quad \text{and consequently,}$$

$$A^n : B^n :: C^n : D^n.$$

That is: *If four quantities are proportional, their like powers or roots will be proportional.*

If we have, for example,

$$2 : 4 :: 3 : 6,$$

we shall have $2^2 : 4^2 :: 3^2 : 6^2$;

---

205. In four proportional quantities, how are like powers or roots?

that is, $4 : 16 :: 9 : 36,$

in which the terms are proportional, the ratio being 4.

206. Let there be two sets of proportions,

$$A : B :: C : D, \text{ which gives } \frac{B}{A} = \frac{D}{C};$$

$$E : F :: G : H, \text{ which gives } \frac{F}{E} = \frac{H}{G}.$$

Multiply them together, member by member, we have,

$$\frac{B \times F}{A \times E} = \frac{D \times H}{C \times G},$$

$$A \times E : B \times F :: C \times G : D \times H.$$

That is: *In two sets of proportional quantities, the products of the corresponding terms are proportional.*

Thus, if we have the two proportions,

$8 : 16 :: 10 : 20,$

and, $3 : 4 :: 6 : 8,$

we shall have, $24 : 64 :: 60 : 160.$

GEOMETRICAL PROGRESSION.

207. We have thus far only considered the case in which the ratio of the first term to the second is the same as that of the third to the fourth.

---

206. In two sets of proportions, how are the products of the corresponding terms?

207. What is a geometrical progression? What is the ratio of the progression? If any term of a progression be multiplied by the ratio, what will the product be? If any term be divided by the ratio, what

If we have the farther condition, that the ratio of the second term to the third shall also be the same as that of the first to the second, or of the third to the fourth, we shall have a series of numbers, each one of which, divided by the preceding one, will give the same ratio. Hence, if any term be multiplied by this quotient, the product will be the succeeding term. A series of numbers so formed, is called a *geometrical progression*. Hence,

A *Geometrical Progression*, or *progression by quotients*, is a series of terms, each of which is equal to the preceding term multiplied by a *constant number*, which number is called the *ratio* of the progression. Thus,

$$1 : 3 : 9 : 27 : 81 : 243, \text{ \&c.},$$

is a geometrical progression, in which the ratio is 3. It is written by merely placing two dots between the terms.

Also, $$64 : 32 : 16 : 8 : 4 : 2 : 1,$$

is a geometrical progression in which the ratio is *one-half*.

In the first progression each term is contained three times in the one that follows, and hence the ratio is 3. In the second, each term is contained one-half times in the one which follows, and hence the ratio is one-half.

The first is called an *increasing* progression, and the second a *decreasing* progression.

Let a, b, c, d, e, f, . . . be numbers, in a progression by quotients · they are written thus:

$$a : b : c : d : e : f : g \ldots$$

and it is enunciated in the same manner as a progression by differences. It is necessary, however, to make the distinc-

---

will the quotient be? How is a progression by quotients written? Which of the terms is only an antecedent? Which only a consequent? How may each of the others be considered?

tion, that one is a series formed by equal differences, and the other a series formed by equal quotients or ratios. It should be remarked that each term is at the same time an antecedent and a consequent, except the first, which is only an antecedent, and the last, which is only a consequent.

208. Let r denote the ratio of the progression,

$$a : b : c : d \ldots$$

r being > 1 when the progression is *increasing*, and $r < 1$, when it is *decreasing*. Then, since,

$$\frac{b}{a} = r, \quad \frac{c}{b} = r, \quad \frac{d}{c} = r, \quad \frac{e}{d} = r, \quad \&c.,$$

we have,

$$b = ar, \quad c = br = ar^2, \quad d = cr = ar^3, \quad e = dr = ar^4,$$
$$f = er = ar^5 \ldots$$

that is, the second term is equal to ar, the third to ar^2, the fourth to ar^3, the fifth to ar^4, &c.; and in general, the nth term, that is, one which has $n - 1$ terms before it, is expressed by ar^{n-1}.

Let l be this term: we then have the formula,

$$l = ar^{n-1},$$

by means of which we can obtain any term without being obliged to find all the terms which precede it. Hence, to find the last term of a progression, we have the following

RULE.

I. *Raise the ratio to a power whose exponent is one less than the number of terms.*

II. *Multiply the power thus found by the first term: the product will be the required term.*

---

208. By what letter do we denote the ratio of a progression? In an increasing progression is r greater or less than 1? In a decreasing pro-

EXAMPLES.

1. Find the 5th term of the progression,

$$2 : 4 : 8 : 16 \ldots$$

in which the first term is 2, and the common ratio 2.

$$5\text{th term} = 2 \times 2^4 = 2 \times 16 = 32. \textit{ Ans.}$$

2. Find the 8th term of the progression,

$$2 : 6 : 18 : 54 \ldots$$

$$8\text{th term} = 2 \times 3^7 = 2 \times 2187 = 4374. \textit{ Ans.}$$

3. Find the 6th term of the progression,

$$2 : 8 : 32 : 128 \ldots$$

$$6\text{th term} = 2 \times 4^5 = 2 \times 1024 = 2048. \textit{ Ans}$$

4. Find the 7th term of the progression,

$$3 : 9 : 27 : 81 \ldots$$

$$7\text{th term} = 3 \times 3^6 = 3 \times 729 = 2187. \textit{ Ans.}$$

5. Find the 6th term of the progression,

$$4 : 12 : 36 : 108 \ldots$$

$$6\text{th term} = 4 \times 3^5 = 4 \times 243 = 972. \textit{ Ans.}$$

6. A person agreed to pay his servant 1 cent for the first day, two for the second, and four for the third, doubling every day for ten days: how much did he receive on the tenth day? *Ans.* $5.12.

---

gression is r greater or less than 1? If a is the first term and r the ratio, what is the second term equal to? What the third? What the fourth? What is the last term equal to? Give the rule for finding the last term.

7. What is the 8th term of the progression,

$$9 : 36 : 144 : 576 \ldots$$

8th term $= 9 \times 4^7 = 9 \times 16384 = 147456$. *Ans.*

8. Find the 12th term of the progression,

$$64 : 16 : 4 : 1 : \frac{1}{4} \ldots$$

12th term $= 64\left(\frac{1}{4}\right)^{11} = \frac{4^3}{4^{11}} = \frac{1}{4^8} = \frac{1}{65536}$. *Ans.*

209. We will now proceed to determine the sum of n terms of a progression,

$$a : b : c : d : e : f : \ldots : i : k : l;$$

l denoting the nth term.

We have the equations (Art. 208),

$$b = ar, \quad c = br, \quad d = cr, \quad e = dr, \ldots k = ir, \quad l = kr,$$

and by adding them all together, member to member, we deduce,

$$\underbrace{b+c+d+e+\ldots+k+l}_{\text{Sum of 1st members.}} = \underbrace{(a+b+c+d+\ldots+i+k)r}_{\text{Sum of 2d members.}};$$

in which we see that the first member contains all the terms but a, and the polynomial, within the parenthesis in the second member, contains all the terms but l. Hence, if we call the sum of the terms S, we have,

$$S - a = (S - l)r = Sr - lr, \quad \therefore Sr - S = lr - a$$

whence, $$S = \frac{lr - a}{r - 1}.$$

---

209. Give the rule for finding the sum of the series. What is the first step? What the second? What the third?

Therefore, to obtain the sum of all the terms, or sum of the series of a geometrical progression, we have the

RULE.

I. *Multiply the last term by the ratio:*

II. *Subtract the first term from the product:*

III. *Divide the remainder by the ratio diminished by* 1 *and the quotient will be the sum of the series.*

1. Find the sum of eight terms of the progression,

$$2 : 6 : 18 : 54 : 162 \ldots 2 \times 3^7 = 4374.$$

$$S = \frac{lr - a}{r - 1} = \frac{13122 - 2}{2} = 6560.$$

2. Find the sum of the progression,

$$2 : 4 : 8 : 16 : 32.$$

$$S = \frac{lr - a}{r - 1} = \frac{64 - 2}{1} = 62.$$

3. Find the sum of ten terms of the progression,

$$2 : 6 : 18 : 54 : 162 \ldots 2 \times 3^9 = 39366.$$

Ans. 59048.

4. What debt may be discharged in a year, or twelve months, by paying $1 the first month, $2 the second month, $4 the third month, and so on, each succeeding payment being double the last; and what will be the last payment?

Ans. { Debt, . . . $4095
Last payment, $2048

5. A daughter was married on New-Year's day. Her father gave her 1*s.*, with an agreement to double it on the first of the next month, and at the beginning of each succeeding month to double what she had previously received. How much did she receive? *Ans.* £204 15*s.*

6. A man bought ten bushels of wheat, on the condition that he should pay 1 cent for the first bushel, 3 for the second, 9 for the third, and so on to the last: what did he pay for the last bushel, and for the ten bushels?

Ans. { Last bushel, $196 83.
Total cost, $295 24.

7. A man plants 4 bushels of barley, which, at the first harvest, produced 32 bushels; these he also plants, which, in like manner, produce 8 fold; he again plants all his crop, and again gets 8 fold, and so on for 16 years: what is his last crop, and what the sum of the series?

Ans. { Last, 140737488355328 bush.
Sum, 160842843834660.

210. When the progression is decreasing, we have, $r < 1$, and $l < a$; the above formula,

$$S = \frac{lr - a}{r - 1},$$

for the sum, is then written under the form,

$$S = \frac{a - lr}{1 - r},$$

in order that the two terms of the fraction may be positive.

1. Find the sum of the terms of the progression,

$$32 : 16 : 8 : 4 : 2$$

$$S = \frac{a - lr}{1 - r} = \frac{32 - 2 \times \frac{1}{2}}{\frac{1}{2}} = \frac{31}{\frac{1}{2}} = 62.$$

---

210. What is the formula for the sum of the series of a decreasing progression?

2. Find the sum of the first twelve terms of the progression,

$$64 : 16 : 4 : 1 : \frac{1}{4} : \ldots : 64\left(\frac{1}{4}\right)^{11}, \text{ or } \frac{1}{65536}.$$

$$S = \frac{a - lr}{1 - r} = \frac{64 - \frac{1}{65536} \times \frac{1}{4}}{\frac{3}{4}} = \frac{256 - \frac{1}{65536}}{3} = 85 + \frac{65535}{196608}$$

211. Remark.—We perceive that the principal difficulty consists in obtaining the numerical value of the last term, a tedious operation, even when the number of terms is not very great.

3. Find the sum of six terms of the progression,

512 : 128 : 32 . . .

Ans. $682\frac{1}{2}$

4. Find the sum of seven terms of the progression,

2187 : 729 : 243 . . .

Ans. 3279

5. Find the sum of six terms of the progression,

972 : 324 : 108 . . .

Ans. 1456.

6. Find the sum of eight terms of the progression,

147456 : 36864 : 9216 . . .

Ans. 196605.

OF PROGRESSIONS HAVING AN INFINITE NUMBER OF TERMS

212. Let there be the decreasing progression,

$$a : b : c : d : e : f : \ldots$$

---

212. When the progression is decreasing, and the number of terms infinite, what is the expression for the value of the sum of the series?

containing an indefinite number of terms. In the formula,

$$S = \frac{a - lr}{1 - r},$$

substitute for l its value, ar^{n-1}, (Art. 208), and we have,

$$S = \frac{a - ar^n}{1 - r},$$

which expresses the sum of n terms of the progression. This may be put under the form,

$$S = \frac{a}{1 - r} - \frac{ar^n}{1 - r}.$$

Now, since the progression is decreasing, r is a proper fraction; and r^n is also a fraction, which diminishes as n increases. Therefore, the greater the number of terms we take, the more will $\frac{a}{1 - r} \times r^n$ diminish, and consequently, the more will the entire sum of all the terms approximate to an equality with the first part of S, that is, to $\frac{a}{1 - r}$. Finally, when n is taken greater than any given number, or $n =$ infinity, then $\frac{a}{1 - r} \times r^n$ will be less than any given number, or will become equal to 0; and the expression, $\frac{a}{1 - r}$, will then represent the true value of the sum of all the terms of the series. Whence we may conclude, that the expression for *the sum of the terms of a decreasing progression, in which the number of terms is infinite*, is,

$$S = \frac{a}{1 - r};$$

that is, *equal to the first term, divided by* 1 *minus the ratio.*

This is, properly speaking, the *limit* to which the *partial sums* approach, as we take a greater number of terms in the progression. The difference between these sums and $\frac{a}{1-r}$, may be made as small as we please, but will only become *nothing* when the number of terms is infinite.

EXAMPLES.

1. Find the sum of

$$1 : \frac{1}{3} : \frac{1}{9} : \frac{1}{27} : \frac{1}{81}, \text{ to infinity.}$$

We have, for the expression of the sum of the terms,

$$S = \frac{a}{1-r} = \frac{1}{1-\frac{1}{3}} = \frac{3}{2} = 1\tfrac{1}{2}. \quad \textit{Ans.}$$

The error committed by taking this expression for the value of the sum of the n first terms, is expressed by

$$\frac{a}{1-r} \times r^n = \frac{3}{2}\left(\frac{1}{3}\right)^n.$$

First take $n = 5$; it becomes,

$$\frac{3}{2}\left(\frac{1}{3}\right)^5 = \frac{1}{2 \,.\, 3^4} = \frac{1}{162}.$$

When $n = 6$, we find,

$$\frac{3}{2}\left(\frac{1}{3}\right)^6 = \frac{1}{162} \times \frac{1}{3} = \frac{1}{486}.$$

Hence, we see, that the *error committed* by taking $\frac{3}{2}$ for the sum of a certain number of terms, is less in proportion as this number is greater.

2. Again, take the progression,

$$1 : \frac{1}{2} : \frac{1}{4} : \frac{1}{8} : \frac{1}{16} : \frac{1}{32} : \&c. \ldots$$

We have, $S = \dfrac{a}{1 - r} = \dfrac{1}{1 - \dfrac{1}{2}} = 2.$ *Ans.*

3. What is the sum of the progression,

$$1,\ \frac{1}{10},\ \frac{1}{100},\ \frac{1}{1000},\ \frac{1}{10000},\ \text{\&c., to infinity.}$$

$$S = \frac{a}{1 - r} = \frac{1}{1 - \dfrac{1}{10}} = 1\frac{1}{9}. \quad \textit{Ans.}$$

213. In the several questions of geometrical progression, there are five numbers to be considered:

1st.	The first term, . .	a.
2d.	The ratio, ...	r.
3d.	The number of terms,	n.
4th.	The last term, . .	l.
5th.	The sum of the terms,	S.

214. We shall terminate this subject by solving this problem:

To find a mean proportional between any two numbers, as m and n.

Denote the required mean by x. We shall then have (Art. 197),

$$x^2 = m \times n;$$

and hence, $$x = \sqrt{m \times n}.$$

---

213. How many numbers are considered in a geometrical progression? What are they?

214. How do you find a mean proportional between two numbers?

That is: *Multiply the two numbers together, and extract the square root of the product.*

1. What is the geometrical mean between the numbers 2 and 8?

$$\text{Mean} = \sqrt{8 \times 2} = \sqrt{16} = 4. \quad \textit{Ans.}$$

2. What is the mean between 4 and 16? *Ans.* 8.
3. What is the mean between 3 and 27? *Ans.* 9.
4. What is the mean between 2 and 72? *Ans.* 12.
5. What is the mean between 4 and 64? *Ans.* 16.

CHAPTER X.

OF LOGARITHMS.

215. THE nature and properties of the logarithms in common use, will be readily understood by considering attentively the different powers of the number 10. They are,

$$\begin{aligned} 10^0 &= 1 \\ 10^1 &= 10 \\ 10^2 &= 100 \\ 10^3 &= 1000 \\ 10^4 &= 10000 \\ 10^5 &= 100000 \\ \&c., &\quad \&c. \end{aligned}$$

It is plain that the *exponents* 0, 1, 2, 3, 4, 5, &c., form an arithmetical series of which the common difference is 1; and that the numbers 1, 10, 100, 1000, 10000, 100000, &c., form a geometrical progression of which the common ratio is 10. The number 10 is called the *base* of the system of logarithms; and the exponents 0, 1, 2, 3, 4, 5, &c., are the *logarithms* of

---

215. What relation exists between the exponents 1, 2, 3, &c.? How are the corresponding numbers 10, 100, 1000? What is the common difference of the exponents? What is the common ratio of the corresponding numbers? What is the base of the common system of logarithms? What are the exponents? Of what number is the exponent 1 the logarithm? The exponent 2? The exponent 3?

the numbers which are produced by raising 10 to the powers denoted by those exponents.

216. If we denote the logarithm of any number by m, then the number itself will be the mth power of 10; that is, if we represent the corresponding number by M,

$$10^m = M.$$

Thus, if we make $m = 0$, M will be equal to 1; if $m = 1$, M will be equal to 10, &c. Hence,

The logarithm of a number is the exponent of the power to which it is necessary to raise the base of the system in order to produce the number.

217. If, as before, 10 denotes the base of the system of logarithms, m any exponent, and M the corresponding number, we shall then have,

$$10^m = M, \qquad (1.)$$

in which m is the logarithm of M.

If we take a second exponent n, and let N denote the corresponding number, we shall have,

$$10^n = N, \qquad (2.)$$

in which n is the logarithm of N.

If, now, we multiply the first of these equations by the second, member by member, we have,

$$10^m \times 10^n = 10^{m+n} = M \times N;$$

but since 10 is the base of the system, $m + n$ is the logarithm $M \times N$; hence,

---

216. If we denote the base of a system by 10, and the exponent by m, what will represent the corresponding number? What is the logarithm of a number?

217. To what is the sum of the logarithms of any two numbers equal? To what, then, will the addition of logarithms correspond?

The sum of the logarithms of any two numbers is equal to the logarithm of their product.

Therefore, *the addition of logarithms corresponds to the multiplication of their numbers.*

218. If we divide Equation (1) by Equation (2), member by member, we have,

$$\frac{10^m}{10^n} = 10^{m-n} = \frac{M}{N};$$

but since 10 is the base of the system, $m - n$ is the logarithm of $\frac{M}{N}$; hence,

If one number be divided by another, the logarithm of the quotient will be equal to the logarithm of the dividend, diminished by that of the divisor.

Therefore, *the subtraction of logarithms corresponds to the division of their numbers.*

219. Let us examine further the equations,

$$\begin{aligned} 10^0 &= 1 \\ 10^1 &= 10 \\ 10^2 &= 100 \\ 10^3 &= 1000 \\ \&c., &\quad \&c. \end{aligned}$$

It is plain that the logarithm of 1 is 0, and that the logarithm of any number between 1 and 10, is greater than

---

218. If one number be divided by another, what will the logarithm of the quotient be equal to? To what, then, will the subtraction of logarithms correspond?

219. What is the logarithm of 1? Between what limits are the logarithms of all numbers between 1 and 10? How are they generally expressed?

0 and less than 1. The logarithm is generally expressed by decimal fractions; thus,

$$\log 2 = 0.301030.$$

The logarithm of any number greater than 10 and less than 100, is greater than 1 and less than 2, and is expressed by 1 and a decimal fraction; thus,

$$\log 50 = 1.698970.$$

The part of the logarithm which stands at the left of the decimal point, is called the *characteristic* of the logarithm. The characteristic is always *one less than the number of places of figures in the number whose logarithm is taken.*

Thus, in the first case, for numbers between 1 and 10, there is but one place of figures, and the characteristic is 0. For numbers between 10 and 100, there are two places of figures, and the characteristic is 1; and similarly for other numbers.

TABLE OF LOGARITHMS.

220. A table of logarithms is a table in which are written the logarithms of all numbers between 1 and some other given number. A table showing the logarithms of the numbers between 1 and 100 is annexed. The numbers are written in the column designated by the letter N, and the logarithms in the column designated by Log.

How is it with the logarithms of numbers between 10 and 100? What is that part of the logarithm called which stands at the left of the characteristic? What is the value of the characteristic?

220. What is a table of logarithms? Explain the manner of finding the logarithms of numbers between 1 and 100?

TABLE.

N.	Log.	N.	Log.	N.	Log.	N.	Log.
1	0.000000	26	1.414973	51	1.707570	76	1.880814
2	0.301030	27	1.431364	52	1.716003	77	1.886491
3	0.477121	28	1.447158	53	1.724276	78	1.892095
4	0 602060	29	1.462398	54	1.732394	79	1.897627
5	0.698970	30	1.477121	55	1.740363	80	1.903090
6	0.778151	31	1.491362	56	1.748188	81	1.908485
7	0.845098	32	1.505150	57	1.755875	82	1.913814
8	0.903090	33	1.518514	58	1.763428	83	1.919078
9	0.954243	34	1.531479	59	1.770852	84	1.924279
10	1.000000	35	1.544068	60	1.778151	85	1.929419
11	1.041393	36	1.556303	61	1.785330	86	1.934498
12	1.079181	37	1.568202	62	1.792392	87	1.939519
13	1.113943	38	1.579784	63	1.799341	88	1.944483
14	1.146128	39	1.591065	64	1.806180	89	1.949390
15	1.176091	40	1.602060	65	1.812913	90	1.954243
16	1.204120	41	1.621784	66	1.819544	91	1.959041
17	1.230449	42	1.623249	67	1.826075	92	1.963788
18	1.255273	43	1.633468	68	1.832509	93	1.968483
19	1.278754	44	1.643453	69	1.838849	94	1.973128
20	1.301030	45	1.653213	70	1.845098	95	1.977724
21	1.322219	46	1.662758	71	1.851258	96	1.982271
22	1.342423	47	1.672098	72	1.857333	97	1.986772
23	1.361728	48	1.681241	73	1.863323	98	1.991226
24	1.380211	49	1.690196	74	1.869232	99	1.995635
25	1.397940	50	1.698970	75	1.875061	100	2.000000

EXAMPLES.

1. Let it be required to multiply 8 by 9, by means of logarithms. We have seen, Art. 216, that the sum of the logarithms is equal to the logarithm of the product. Therefore, find the logarithm of 8 from the table, which is 0.903090, and then the logarithm of 9, which is 0.954243; and their sum, which is 1.857333, will be the logarithm of the product. In searching along in the table, we find that 72 stands opposite this logarithm; hence, 72 is the product of 8 by 9.

2. What is the product of 7 by 12?

Logarithm of 7 is, ...	0.845098
Logarithm of 12 is, ...	1.079181
Logarithm of their product, . .	1.924279

and the corresponding number is 84.

3. What is the product of 9 by 11?

Logarithm of 9 is, ...	0.954243
Logarithm of 11 is, . . .	1.041393
Logarithm of their product, .	1.995636

and the corresponding number is 99.

4. Let it be required to divide 84 by 3. We have seen in Art. 218, that the subtraction of Logarithms corresponds to the division of their numbers. Hence, if we find the logarithm of 84, and then subtract from it the logarithm of 3, the remainder will be the logarithm of the quotient.

The logarithm of 84 is, . . .	1.924279
The logarithm of 3 is, . . .	0.477121
Their difference is, ...	1.447158

and the corresponding number is 28.

5. What is the product of 6 by 7?

Logarithm of 6 is, ...	0.778151
Logarithm of 7 is, . . .	0.845098
Their sum is, ...	1.623249

and the corresponding number of the table, 42.

National Series of Standard School-Books,

PUBLISHED BY

A. S. BARNES & COMPANY.

THIS SERIES embraces more than Three Hundred Volumes of Standard Educational Works, composing the most complete and uniformly meritorious collection of text-books ever published by a single firm.

THE SERIES is *complete*, covering every variety and grade of science and literature, from the Primer which guides the lisping tongue of the infant, to the abstruse and difficult "West Point Course."

THE SERIES is *uniformly excellent*. Each volume, among so many, maintains the general standard of merit, and assists, in its place, to round the perfect whole.

THE SERIES is *known and popularly used in every section of the United States, and by every class of citizens*, representing all shades of political opinion and religious belief. In proof of this, it is only necessary to name the following popular works, with which *every one* is familiar, and which fairly represent the whole.

PARKER & WATSON'S NATIONAL and INDEPENDENT READERS & SPELLERS.
Two series; each complete in itself, and of high and lower gradation respectively.

DAVIES' Complete Unified "West Point Course" in MATHEMATICS.
Including Arithmetics, Algebra, Geometry, Calculus, Surveying, and all branches.

MONTEITH & McNALLY'S Unrivaled Course in GEOGRAPHY.
Three books, with optional volumes. This is the only series with Allen's Map-Drawing.

CLARK'S Diagram System for Learning ENGLISH GRAMMAR.
Two Grammars (Primary and Advanced); also Analysis, Key and Chart separate.

BARNES' Series of Brief Books for One Term of Study in HISTORY.
United States History, published in 1871. A General History and others to follow at once.

STEELE'S "Fourteen Weeks" in each branch of NATURAL SCIENCE.
Philosophy, Chemistry, Astronomy, and Geology now ready. Physiology in press.

WORMAN'S Popular Method for Teaching FOREIGN LANGUAGE.
A German Series is now nearly complete, and the French series happily commenced.

WILLARD'S Course of History.
PECK'S GANOT'S Natural Philosophy.
JARVIS' Physiology and Health.
WOOD'S Text-Books in Botany.
PORTER'S Principles of Chemistry.
CHAMBERS' Elements of Zoology.
PAGE'S Elements of Geology.
BEERS' Round-hand Penmanship.
SMITH'S Orthography and Etymology.
BOYD'S Course in English Literature.
SMITH & MARTIN'S Bookkeeping.
PAYSON, DUNTON & CO.'S Writing.
ANDREWS & STODDARD'S Latin Series
CROSBY'S Greek Series.
PUJOL'S Complete French Class Book.
ROOT'S School Music-Books.
SEARING'S Virgil and Classical Texts.
CLEVELAND'S Compendiums.
CHAPMAN'S American Drawing.
MANSFIELD'S Political Manual.
NORTHEND'S Graded Speakers.
CHAMPLIN'S Political Economy.
BERARD'S History of England.
SHERWOOD'S Writing Speller.
BROOKFIELD'S Child's Composition.
TRACY'S School Record.
PECK'S Applied Mathematics.
CHURCH'S Higher Mathematics.
BARTLETT'S College Philosophy.
MAHAN'S Mental Science.
ALDEN'S Text-Book of Ethics.
BROOKS' School Devotion.

☞ ANY BOOK SENT POSTPAID ON RECEIPT OF ITS PRICE. ☜

Who would know more of this unrivaled Series should consult for details, 1. THE DESCRIPTIVE CATALOGUE, free to Teachers; others, 5 cents. 2. THE ILLUSTRATED EDUCATIONAL BULLETIN, periodical organ of the Publishers; full of instruction for Teachers; subscription, 10 cents; sample free. Address

A. S. BARNES & COMPANY, New York, or Chicago.

MATHEMATICS.

DAVIES' NATIONAL COURSE.

ARITHMETIC.

			SLATED.
1.	Davies' Primary Arithmetic,	$ 25	$ 32
2.	Davies' Intellectual Arithmetic,	40	48
3.	Davies' Elements of Written Arithmetic,	50	60
4.	Davies' Practical Arithmetic,	90	1 00
	Key to Practical Arithmetic,	90	
5.	Davies' University Arithmetic,	1 40	1 50
	Key to University Arithmetic,	*1 40	

ALGEBRA.

1.	Davies' New Elementary Algebra,	*1 25	1 35
	Key to Elementary Algebra,	*1 25	
2.	Davies' University Algebra,	1 50	1 60
	Key to University Algebra,	*1 50	
3.	Davies' Bourdon's Algebra,	2 25	2 38
	Key to Bourdon's Algebra,	*2 25	

GEOMETRY.

1.	Davies' Elementary Geometry and Trigonometry,	1 40	1 50
2.	Davies' Legendre's Geometry,	2 25	2 38
3.	Davies' Analytical Geometry and Calculus,	2 50	2 63
4.	Davies' Descriptive Geometry,	2 75	2 88

MENSURATION.

1.	Davies' Practical Mathematics and Mensuration,	1 50	1 60
2.	Davies' Elements of Surveying,	2 50	2 63
3.	Davies' Shades, Shadows, and Perspective,	3 75	3 88

MATHEMATICAL SCIENCE.

Davies' Grammar of Arithmetic,	* 50
Davies' Outlines of Mathematical Science,	*1 00
Davies' Logic and Utility of Mathematics,	*1 50
Davies' Metric System,	*1 50
Davies & Peck's Dictionary of Mathematics,	*5 00

DAVIES' NATIONAL COURSE of MATHEMATICS.

ITS RECORD.

In claiming for this series the first place among American text-books, of what ever class, the Publishers appeal to the magnificent record which its volumes have earned during the *thirty-five years* of Dr. Charles Davies' mathematical labors. The unremitting exertions of a life-time have placed *the modern series* on the same proud eminence among competitors that each of its predecessors has successively enjoyed in a course of constantly improved editions, now rounded to their perfect fruition—for it seems almost that this science is susceptible of no further demonstration.

During the period alluded to, many authors and editors in this department have started into public notice, and by borrowing ideas and processes original with Dr. Davies, have enjoyed a brief popularity, but are now almost unknown. Many of the series of to-day, built upon a similar basis, and described as "modern books," are destined to a similar fate; while the most far-seeing eye will find it difficult to fix the time, on the basis of any data afforded by their past history, when these books will cease to increase and prosper, and fix a still firmer hold on the affection of every educated American.

One cause of this unparalleled popularity is found in the fact that the enterprise of the author did not cease with the original completion of his books. Always a practical teacher, he has incorporated in his text-books from time to time the advantages of every improvement in methods of teaching, and every advance in science. During all the years in which he has been laboring, he constantly submitted his own theories and those of others to the practical test of the class-room—approving, rejecting, or modifying them as the experience thus obtained might suggest. In this way he has been able to produce an almost perfect series of class-books, in which every department of mathematics has received minute and exhaustive attention.

Nor has he yet retired from the field. Still in the prime of life, and enjoying a ripe experience which no other living mathematician or teacher can emulate, his pen is ever ready to carry on the good work, as the progress of science may demand. Witness his recent exposition of the "Metric System," which received the official endorsement of Congress, by its Committee on Uniform Weights and Measures.

DAVIES' SYSTEM IS THE ACKNOWLEDGED NATIONAL STANDARD FOR THE UNITED STATES, for the following reasons:—

1st. It is the basis of instruction in the great national schools at West Point and Annapolis.

2d. It has received the *quasi* endorsement of the National Congress.

3d. It is exclusively used in the public schools of the National Capital.

4th. The officials of the Government use it as authority in all cases involving mathematical questions.

5th. Our great soldiers and sailors commanding the national armies and navies were educated in this system. So have been a majority of eminent scientists in this country. All these refer to "Davies" as authority.

6th. A larger number of American citizens have received their education from this than from any other series.

7th. The series has a larger circulation throughout the whole country than any other, being *extensively used in every State in the Union.*

Davies' National Course of Mathematics.

TESTIMONIALS.

From L. Van Bokkelen, *State Superintendent Public Instruction, Maryland.*

The series of Arithmetics edited by Prof. Davies, and published by your firm, have been used for many years in the schools of several counties, and the city of Baltimore, and have been approved by teachers and commissioners.

Under the law of 1865, establishing a uniform system of Free Public Schools, these Arithmetics were unanimously adopted by the State Board of Education, after a careful examination, and are now used in all the Public Schools of Maryland.

These facts evidence the high opinion entertained by the School Authorities of the value of the series theoretically and practically.

From Horace Webster, *President of the College of New York.*

The undersigned has examined, with care and thought, several volumes of Davies' Mathematics, and is of the opinion that, as a whole, it is the most complete and best course for Academic and Collegiate instruction, with which he is acquainted.

From David N. Camp, *State Superintendent of Common Schools, Connecticut.*

I have examined Davies' Series of Arithmetics with some care. The language is clear and precise; each principle is thoroughly analyzed, and the whole so arranged as to facilitate the work of instruction. Having observed the satisfaction and success with which the different books have been used by eminent teachers, it gives me pleasure to commend them to others.

From J. O. Wilson, *Chairman Committee on Text-Books, Washington, D. C.*

I consider Davies' Arithmetics decidedly superior to any other series, and in this opinion I am sustained, I believe, by the entire Board of Education and Corps of Teachers in this city, where they have been used for several years past.

From John L. Campbell, *Professor of Mathematics, Wabash College, Indiana.*

A proper combination of abstract reasoning and practical illustration is the chief excellence in Prof. Davies' Mathematical works. I prefer his Arithmetics, Algebras, Geometry and Trigonometry to all others now in use, and cordially recommend them to all who desire the advancement of sound learning.

From Major J. H. Whittlesey, *Government Inspector of Military Schools.*

Be assured, I regard the works of Prof. Davies, with which I am acquainted, as by far the best text-books in print on the subjects which they treat. I shall certainly encourage their adoption wherever a word from me may be of any avail.

From T. McC. Ballantine, *Prof. Mathematics Cumberland College, Kentucky.*

I have long taught Prof. Davies' Course of Mathematics, and I continue to like their working.

From John McLean Bell, B. A., *Prin. of Lower Canada College.*

I have used Davies' Arithmetical and Mathematical Series as text-books in the schools under my charge for the last six years. These I have found of great efficacy in exciting, invigorating, and concentrating the intellectual faculties of the young.

Each treatise serves as an introduction to the next higher, by the similarity of its reasonings and methods; and the student is carried forward, by easy and gradual steps, over the whole field of mathematical inquiry, and that, too, in a *shorter* time than is usually occupied in mastering a single department. I sincerely and heartily recommend them to the attention of my fellow-teachers in Canada.

From D. W. Steele, *Prin. Philekoian Academy, Cold Springs, Texas.*

I have used Davies' Arithmetics till I know them nearly by heart. A better series of school-books never were published. I have recommended them until they are now used in all this region of country.

A large mass of similar "Opinions" may be obtained by addressing the publishers for special circular for Davies' Mathematics. New recommendations are published in current numbers of the *Educational Bulletin.*

MATHEMATICS—Continued.

ARITHMETICAL EXAMPLES.

Reuck's Examples in Denominate Numbers	$ 50
Reuck's Examples in Arithmetic	1 00

These volumes diffei from the ordinary arithmetic in their peculiarly *practical* character. They are composed mainly of examples, and afford the most severe and thorough discipline for the mind. While a book which should contain a complete treatise of theory and practice would be too cumbersome for every-day use, the insufficiency of *practical* examples has been a source of complaint.

HIGHER MATHEMATICS.

Church's Elements of Calculus	2 50
Church's Analytical Geometry	2 50
Church's Descriptive Geometry, with Shades, Shadows, and Perspective	4 00

These volumes constitute the "West Point Course" in their several departments.

Courtenay's Elements of Calculus	3 00

A work especially popular at the South.

Hackley's Trigonometry	2 50

With applications to navigation and surveying, nautical and practical geometry and geodesy.

APPLIED MATHEMATICS.

Peck's Ganot's Popular Physics	1 75
Peck's Elements of Mechanics	2 00
Peck's Practical Calculus	1 75

Prof. W G. Peck, of Columbia College, has designed the first of these works for the ordinary wants of schools in the department of Natural Philosophy. The work enjoys a high reputation.

The Mechanics and Calculus are the briefest treatises on those subjects now published. Their methods are purely practical, and unembarrassed by the details which rather confuse than simplify science.

SLATED ARITHMETICS.

This consists of the application of an artificially slated surface to the inner cover of a book, with flap of the same opening outward, so that students may refer to the book and use the slate at one and the same time, and as though the slate were detached. When folded up, the slate preserves examples and memoranda til' needed. The material used is as durable as the stone slate. The additional cost of books thus improved is trifling.

HISTORY.

Monteith's Youth's History, ...$ 75

A History of the United States for beginners. It is arranged upon the catechetical plan, with illustrative maps and engravings, review questions, dates in parentheses (that their study may be optional with the younger class of learners), and interesting Biographical Sketches of all persons who have been prominently identified with the history of our country.

Willard's United States, School edition, . . . 1 40

Do. do. University edition, . 2 25

The plan of this standard work is chronologically exhibited in front of the title-page; the Maps and Sketches are found useful assistants to the memory, and dates, usually so difficult to remember, are so systematically arranged as in a great degree to obviate the difficulty. Candor, impartiality, and accuracy, are the distinguishing features of the narrative portion.

Willard's Universal History, ... 2 25

The most valuable features of the "United States" are reproduced in this. The peculiarities of the work are its great conciseness and the prominence given to the chronological order of events. The margin marks each successive era with great distinctness, so that the pupil retains not only the event but its time, and thus fixes the order of history firmly and usefully in his mind. Mrs. Willard's books are constantly revised, and at all times written up to embrace important historical events of recent date.

Berard's History of England, ... 1 75

By an authoress well known for the success of her History of the United States. The social life of the English people is felicitously interwoven, as in fact, with the civil and military transactions of the realm.

Ricord's History of Rome, ... 1 75

Possesses the charm of an attractive romance. The Fables with which this history abounds are introduced in such a way as not to deceive the inexperienced, while adding materially to the value of the work as a reliable index to the character and institutions, as well as the history of the Roman people.

Hanna's Bible History, ... 1 25

The only compendium of Bible narrative which affords a connected and chronological view of the important events there recorded, divested of all superfluous detail.

Summary of History, Complete 60

American History, $0 40. French and Eng. Hist. 35

A well proportioned outline of leading events, condensing the substance of the more extensive text-book in common use into a series of statements so brief, that every word may be committed to memory, and yet so comprehensive that it presents an accurate though general view of the whole continuous life of nations.

Marsh's Ecclesiastical History, ... 2 00

Questions to ditto, ... 75

Affording the History of the Church in all ages, with accounts of the pagan world during Biblical periods, and the character, rise, and progress of all Religions, as well as the various sects of the worshipers of Christ. The work is entirely non-sectarian, though strictly catholic.

HISTORY—Continued.

BARNES' ONE-TERM HISTORY.

A Brief History of the United States, . . . $1 50

This is probably the MOST ORIGINAL SCHOOL-BOOK published for many years, in any department. A few of its claims are the following:

1. **Brevity.**—The text is complete for Grammar School or intermediate classes, in 290 12mo pages, large type. It may readily be completed, if desired, in one term of study.

2. **Comprehensiveness.**—Though so brief, this book contains the pith of all the wearying contents of the larger manuals, and a great deal more than the memory usually retains from the latter.

3. **Interest** has been a prime consideration. Small books have heretofore been bare, full of dry statistics, unattractive. This one is charmingly written, replete with anecdote, and brilliant with illustration.

4. **Proportion of Events.**—It is remarkable for the discrimination with which the different portions of our history are presented according to their importance. Thus the older works being already large books when the civil war took place, give it less space than that accorded to the Revolution.

5. **Arrangement.**—In six epochs, entitled respectively, Discovery and Settlement. the Colonies, the Revolution, Growth of States, the Civil War, and Current Events.

6. **Catch Words.**—Each paragraph is preceded by its leading thought in prominent type, standing in the student's mind for the whole paragraph.

7. **Key Notes.**—Analogous with this is the idea of grouping battles, etc., about some central event, which relieves the sameness so common in such descriptions, and renders each distinct by some striking peculiarity of its own.

8. **Foot Notes.**—These are crowded with interesting matter that is not strictly a part of history proper. They may be learned or not, at pleasure. They are certain in any event to be read.

9. **Biographies** of all the leading characters are given in full in foot-notes.

10. **Maps.**—Elegant and distinct Maps from engravings on copper-plate, and beautifully colored, precede each epoch, and contain all the places named.

11. **Questions** are at the back of the book, to compel a more independent use of the text. Both text and questions are so worded that the pupil must give intelligent answers IN HIS OWN WORDS. "Yes" and "No" will not do.

12. **Historical Recreations.**—These are additional questions to test the student's knowledge, in review, as: "What trees are celebrated in our history?" "When did a fog save our army?" "What Presidents died in office?" "When was the Mississippi our western boundary?" "Who said, 'I would rather be right than President?'" etc.

13. **The Illustrations,** about seventy in number, are the work of our best artists and engravers, produced at great expense. They are vivid and interesting, and mostly upon subjects never before illustrated in a school-book.

14. **Dates.**—Only the leading dates are given in the text, and these are so associated as to assist the memory, but at the head of each page is the date of the event first mentioned, and at the close of each epoch a summary of events and dates.

15. **The Philosophy of History** is studiously exhibited—the causes and effects of events being distinctly traced and their interconnection shown.

16. **Impartiality.**—All sectional, partisan, or denominational views are avoided. Facts are stated after a careful comparison of all authorities without the least prejudice or favor.

17. **Index.**—A verbal index at the close of the book perfects it as a work of reference.

It will be observed that the above are all particulars in which School Histories have been signally defective, or altogether wanting. Many other claims to favor it shares in common with its predecessors.

BARNES' BRIEF UNITED STATES HISTORY.

Already prescribed under authority of law for exclusive and uniform use in the public schools of TWO STATES—Texas and Arkansas. Also adopted for HUNDREDS of important CITIES and TOWNS—among which we name Brooklyn, N.Y.; Jersey City, N.J.; Scranton, Pa.; Wilmington, Del.; Portland, Me.; Springfield, Mass.; Louisville, Ky.; Vicksburg, Miss.; Madison, Wis.; Rochester, Minn.; Macon, Mo.; Springfield, Ill., etc.

SOME TESTIMONIALS.

From HON. J. M. MCKENZIE, *Supt. Pub. Inst., Nebraska.*

I have examined your "Brief History of the United States," and like it *real well;* and were I teaching a graded school, I think I should use it as a text-book.

From HON. H. B. WILSON, *Supt. Pub. Inst., Minnesota.*

I have read with much interest the "One-Term History of the United States." I am much pleased with it. In my judgment, it contains all of the United States history that the majority of pupils in our common schools can spare time to study.

From PRES. EDWARD BROOKS, *Millersville State Normal School, Pa.*

It is a work that will be a favorite with teachers and pupils. Its scope and style especially adapt it for use in our public schools. I cordially commend it to teachers desiring to introduce an interesting and practical text-book upon this subject.

From PRES. BARKER, *Buffalo State Normal School, N.Y.*

In the copy of your "Brief History," before me, the important items to be learned in history seem most ingeniously brought out and kept in the foreground. These items are *time*, *persons*, *places*, and *events*. It has the appearance of an exceedingly fresh and systematic work. I think I shall put it into my classes

From PROF. WM. F. ALLEN, *State Univ. of Wisconsin.*

I think the author of the new "Brief History of the United States" has been very successful in combining brevity with sufficient fullness and interest. *Particularly*, he has avoided the excessive number of names and dates that most histories contain. Two features that I like *very much* are the *anecdotes* at the foot of the page and the "*Historical Recreations*" in the Appendix. The latter, I think, is quite a *new* feature, and the other is *very* well executed.

From S. G. WRIGHT, *Assist.-Supt. Pub. Inst., Kansas.*

It is with extreme pleasure we submit our recommendation of the "Brief History of the United States." It meets the needs of young and older children, combining concision with perspicuity, and if "brevity is the soul of wit," this "Brief History" contains not only that well-chosen ingredient, but wisdom sufficient to enlighten those students who are wearily longing for a "new departure" from certain old and uninteresting presentations of fossilized writers. We congratulate a progressive public upon a progressive book.

From HON. NEWTON BATEMAN, *Supt. Pub. Inst., Illinois.*

Barnes' One-Term History of the United States is an exceedingly attractive and spirited little book. Its claim to several new and valuable features seems well founded. Under the form of six well-defined Epochs, the History of the United States is traced tersely, yet pithily, from the earliest times to the present day. A good map precedes each epoch, whereby the history and geography of the period may be studied together, *as they always should be.* The syllabus of each paragraph is made to stand in such bold relief, by the use of large, heavy type, as to be of much *mnemonic* value to the student. The book is written in a sprightly and piquant style, the interest never flagging from beginning to end—a rare and difficult achievement in works of this kind.

From the "Chicago Schoolmaster" (Editorial).

A thorough examination of Barnes' Brief History of the United States brings the examiner to the conclusion that it is a superior book in almost every respect. The book is neat in form, and of good material. The type is clear, large, and distinct. The facts and dates are correct. The arrangement of topics is just the thing needed in a history text-book. By this arrangement the pupil can see at once what he is expected to do. The topics are well selected, embracing the leading ideas or principal events of American history. . . . The book as a whole is much superior to any I have examined. So much do I think this, that I have ordered it for my class, and shall use it in my school. (Signed) B. W. BAKER.

PENMANSHIP.

Beers' System of Progressive Penmanship.
Per dozen $1 68

This "round hand" system of Penmanship in twelve numbers, commends itself by its simplicity and thoroughness. The first four numbers are primary books. Nos. 5 to 7, advanced books for boys. Nos. 8 to 10, advanced books for girls. Nos. 11 and 12, ornamental penmanship. These books are printed from steel plates (engraved by McLees), and are unexcelled in mechanical execution. Large quantities are annually sold.

Beers' Slated Copy Slips, per set *50

All beginners should practice, for a few weeks, slate exercises, familiarizing them with the form of the letters, the motions of the hand and arm, &c., &c. These copy slips, 32 in number, supply all the copies found in a complete series of writing-books, at a trifling cost.

Payson, Dunton & Scribner's Copy-B'ks. P. doz. *1 80

The National System of Penmanship, in three distinct series—(1) Common School Series, comprising the first six numbers; (2) Business Series, Nos. 8, 11, and 12; (3) Ladies' Series, Nos. 7, 9, and 10.

Fulton & Eastman's Chirographic Charts, *3 75

To embellish the school room walls, and furnish class exercise in the elements of Penmanship.

Payson's Copy-Book Cover, per hundred .*4 00

Protects every page except the one in use, and furnishes "lines" with proper slope for the penman, under. Patented.

National Steel Pens, Card with all kinds . . . *15

Pronounced by competent judges the perfection of American-made pens, and superior to any foreign article.

SCHOOL SERIES.	
School Pen, per gross, . . $	60
Academic Pen, do . .	63
Fine Pointed Pen, per gross	70
POPULAR SERIES.	
Capitol Pen, per gross, . .	1 00
do do pr. box of 2 doz.	25
Bullion Pen (imit. gold) pr. gr.	75
Ladies' Pen do	63

Index Pen, per gross . . .	75
BUSINESS SERIES.	
Albata Pen, per gross, . .	40
Bank Pen, do . .	70
Empire Pen, do . .	70
Commercial Pen, per gross .	60
Express Pen, do .	75
Falcon Pen, do .	70
Elastic Pen, do .	75

Stimpson's Scientific Steel Pen, per gross .*2 00

One forward and two backward arches, ensuring great strength, well-balanced elasticity, evenness of point, and smoothness of execution. One gross in twelve contains a Scientific Gold Pen.

Stimpson's Ink-Retaining Holder, per doz. .*2 00

A simple apparatus, which does not get out of order, withholds at a single dip as much ink as the pen would otherwise realize from a dozen trips to the inkstand, which it supplies with moderate and easy flow.

Stimpson's Gold Pen, $3 00; **with Ink Retainer** *4 50

Stimpson's Penman's Card, ... * 50

One dozen Steel Pens (assorted points) and Patent Ink-retaining Pen holder.

BOOK-KEEPING.

Folsom's Logical Book-keeping, ...	$2 00
Folsom's Blanks to Book-keeping, ...	4 50

This treatise embraces the interesting and important discoveries of Prof. Folsom (of the Albany "Bryant & Stratton College"), the partial enunciation of which in lectures and otherwise has attracted so much attention in circles interested in commercial education.

After studying business phenomena for many years, he has arrived at the positive laws and principles that underlie the whole subject of Accounts; finds that the science is based in *Value* as a generic term; that value divides into *two classes* with varied species; that all the exchanges of values are reducible to nine equations; and that all the results of all these exchanges are limited to *thirteen* in number.

As accounts have been universally taught hitherto, without setting out from a radical analysis or definition of values, the science has been kept in great obscurity, and been made as difficult to impart as to acquire. On the new theory, however, these obstacles are chiefly removed. In reading over the first part of it, in which the governing laws and principles are discussed, a person with ordinary intelligence will obtain a fair conception of the *double entry* process of accounts. But when he comes to study thoroughly these laws and principles as there enunciated, and works out the examples and memoranda which elucidate the *thirteen results* of business, the student will neither fail in readily acquiring the science as it is, nor in becoming able intelligently to apply it in the interpretation of business.

Smith & Martin's Book-keeping, ...	1 25
Smith & Martin's Blanks, ...	*60

This work is by a practical teacher and a practical book-keeper. It is of a thoroughly popular class, and will be welcomed by every one who loves to see theory and practice combined in an easy, concise, and methodical form.

The Single Entry portion is well adapted to supply a want felt in nearly all other treatises, which seem to be prepared mainly for the use of wholesale merchants, leaving retailers, mechanics, farmers, etc., who transact the greater portion of the business of the country, without a guide. The work is also commended, on this account, for general use in Young Ladies' Seminaries, where a thorough grounding in the simpler form of accounts will be invaluable to the future housekeepers of the nation.

The treatise on Double Entry Book-keeping combines all the advantages of the most recent methods, with the utmost simplicity of application, thus affording the pupil all the advantages of actual experience in the counting-house, and giving a clear comprehension of the entire subject through a judicious course of mercantile transactions.

The shape of the book is such that the transactions can be presented as in actual practice; and the simplified form of Blanks—three in number—adds greatly to the ease experienced in acquiring the science.

DRAWING.

Chapman's American Drawing Book, . . . *$6 00

The standard American text-book and authority in all branches of art. A compilation of art principles. A manual for the amateur, and basis of study for the professional artist. Adapted for schools and private instruction.

CONTENTS.—"Any one who can Learn to Write can Learn to Draw."—Primary Instruction in Drawing.—Rudiments of Drawing the Human Head.—Rudiments in Drawing the Human Figure.—Rudiments of Drawing.—The Elements of Geometry.—Perspective.—Of Studying and Sketching from Nature.—Of Painting.—Etching and Engraving.—Of Modeling.—Of Composition.—Advice to the American Art-Student.

The work is of course magnificently illustrated with all the original designs.

Chapman's Elementary Drawing Book, . . 1 50

A Progressive Course of Practical Exercises, or a text-book for the training of the eye and hand. It contains the elements from the larger work, and a copy should be in the hands of every pupil; while a copy of the "American Drawing Book," named above, should be at hand for reference by the class.

The Little Artist's Portfolio, ... *50

25 Drawing Cards (progressive patterns), 25 Blanks, and a fine Artist's Pencil, all in one neat envelope.

Clark's Elements of Drawing, ... *1 00

A complete course in this graceful art, from the first rudiments of outline to the finished sketches of landscape and scenery.

Fowle's Linear and Perspective Drawing, . *60

For the cultivation of the eye and hand, with copious illustrations and directions for the guidance of the unskilled teacher.

Monk's Drawing Books—Six Numbers, per set, *2 25

Each book contains *eleven* large patterns, with opposing blanks. No. 1. Elementary Studies. No. 2. Studies of Foliage. No. 3. Landscapes. No. 4. Animals, I. No. 5. Animals, II. No. 6. Marine Views, etc.

Allen's Map-Drawing, . . . 25 cts.; Scale, 25

This method introduces a new era in Map-Drawing, for the following reasons:—1. It is a system. This is its greatest merit.—2. It is easily understood and taught.—3. The eye is trained to exact measurement by the use of a scale.—4. By no special effort of the memory, distance and comparative size are fixed in the mind.—5. It discards useless construction of lines.—6. It can be taught by any teacher, even though there may have been no previous practice in Map-Drawing.—7. Any pupil old enough to study Geography can learn by this System, in a short time, to draw accurate maps.—8. The System is not the result of theory, but comes directly from the school-room. It has been thoroughly and successfully tested there, with all grades of pupils.—9. It is economical, as it requires no mapping plates. It gives the pupil the ability of rapidly drawing accurate maps.

Ripley's Map-Drawing, ... 1 25

Based on the Circle. One of the most efficient aids to the acquirement of a knowledge of Geography is the practice of map-drawing. It is useful for the same reason that the best exercise in orthography is the *writing* of difficult words. Sight comes to the aid of hearing, and a double impression is produced upon the memory. Knowledge becomes less mechanical and more intuitive. The student who has sketched the outlines of a country, and dotted the important places, is little likely to forget either. The impression produced may be compared to that of a traveller who has been over the ground, while more comprehensive and accurate in detail.

MUSIC.

Jepson's Music Readers. 3 vols. . . . Each, 75 cts.

These are not books from which children simply learn songs, parrot-like, but teach the subject progressively—the scholar learning to read music by methods similar to those employed in teaching him to read printed language. Any teacher, however ignorant of music, provided he can, upon trial, simply sound the scale, may teach it without assistance, and will end by being a good singer himself. The "Elementary Music Reader," or first volume, heretofore issued by another publisher, has attained results in the State of Connecticut, where only it has been known, entirely unprecedented in the history of teaching music. The two companion volumes carry the same method into the higher grades.

Nash & Bristow's Cantara. No. 1, $1.15; No. 2, $1.40

The first volume is a complete musical text-book for schools of every grade. No. 2 is a choice selection of Solos and Part Songs. The authors are Directors of Music in the public schools of New York City, in which these books are the standard of instruction.

Curtis' Little Singer,	$0 60
Curtis' School Vocalist,	1 00
Kingsley's School-Room Choir,	60
Kingsley's Young Ladies' Harp,	1 00
Hager's Echo,	75
Perkins' Sabbath Carols,	35

For Sunday-schools.

DEVOTION.

Brooks' School Manual of Devotion, . . . $0 75

This volume contains daily devotional exercises, consisting of a hymn, selections of Scripture for alternate reading by teacher and pupils, and a prayer. Its value for opening and closing school is apparent.

Brooks' School Harmonist, ... *75

Contains appropriate *tunes* for each hymn in the "Manual of Devotion" described above.

NATURAL SCIENCE.

FAMILIAR SCIENCE.

Norton & Porter's First Book of Science, . $1 75

By eminent Professors of Yale College. Contains the principles of Natural Philosophy, Astronomy, Chemistry, Physiology, and Geology. Arranged on the Catechetical plan for primary classes and beginners.

Chambers' Treasury of Knowledge, ... 1 25

Progressive lessons upon—*first*, common things which lie most immediately around us, and first attract the attention of the young mind; *second*, common objects from the Mineral, Animal, and Vegetable kingdoms, manufactured articles, and miscellaneous substances; *third*, a systematic view of Nature under the various sciences. May be used as a Reader or Text-book.

NATURAL PHILOSOPHY.

Norton's First Book in Natural Philosophy, 1 00

By Prof. NORTON, of Yale College. Designed for beginners. Profusely illustrated, and arranged on the Catechetical plan.

Peck's Ganot's Course of Nat. Philosophy, . 1 75

The standard text-book of France, Americanized and popularized by Prof. PECK, of Columbia College. The most magnificent system of illustration ever adopted in an American school-book is here found. For intermediate classes.

Peck's Elements of Mechanics, ... 2 00

A suitable introduction to Bartlett's higher treatises on Mechanical Philosophy, and adequate in itself for a complete academical course.

Bartlett's SYNTHETIC AND ANALYTIC Mechanics, . each 5 00

Bartlett's Acoustics and Optics, ... 3 50

A system of Collegiate Philosophy, by Prof. BARTLETT, of West Point Military Academy.

Steele's 14 Weeks Course in Philos. (see p. 34) 1 50

Steele's Philosophical Apparatus, ... *125 00

Adequate to performing the experiments in the ordinary text-books. The articles will be sold separately, if desired. See special circular for details.

GEOLOGY.

Page's Elements of Geology, ... 1 25

A volume of Chambers' Educational Course. Practical, simple, and eminently calculated to make the study interesting.

Emmons' Manual of Geology, ... 1 25

The first Geologist of the country has here produced a work worthy of his reputation.

Steele's 14 Weeks Course (see p. 34) 1 50

Steele's Geological Cabinet, ...*40 00

Containing 125 carefully selected specimens. In four parts. Sold separately, if desired. See circular for details.

1

2.

3

Davies' Mathematics.

THE WEST POINT COURSE,

And Only Thorough and Complete Mathematical Series.

IN THREE PARTS.

I. COMMON SCHOOL COURSE.

Davies' Primary Arithmetic.—The fundamental principles displayed in Object Lessons.

Davies' Intellectual Arithmetic.—Referring all operations to the unit 1 as the only tangible basis for logical development.

Davies' Elements of Written Arithmetic.—A practical introduction to the whole subject. Theory subordinated to Practice.

Davies' Practical Arithmetic.*—The most successful combination of Theory and Practice, clear, exact, brief, and comprehensive.

II. ACADEMIC COURSE.

Davies' University Arithmetic.*—Treating the subject exhaustively as *a science*, in a logical series of connected propositions.

Davies' Elementary Algebra.*—A connecting link, conducting the pupil easily from arithmetical processes to abstract analysis.

Davies' University Algebra.*—For institutions desiring a more complete but not the fullest course in pure Algebra.

Davies' Practical Mathematics.—The science practically applied to the useful arts, as Drawing, Architecture, Surveying, Mechanics, etc.

Davies' Elementary Geometry.—The important principles in simple form, but with all the exactness of vigorous reasoning.

Davies' Elements of Surveying.—Re-written in 1870. The simplest and most practical presentation for youths of 12 to 13.

III. COLLEGIATE COURSE.

Davies' Bourdon's Algebra.*—Embracing Sturm's Theorem, and a most exhaustive and scholarly course.

Davies' University Algebra.*—A shorter course than Bourdon, for Institutions have less time to give the subject.

Davies' Legendre's Geometry.—Acknowledged *the only* satisfactory treatise of its grade. 300,000 copies have been sold.

Davies' Analytical Geometry and Calculus.—The shorter treatises, combined in one volume, are more available for American courses of study.

Davies' Analytical Geometry. } The original compendiums, for those de-
Davies' Diff. & Int. Calculus. } siring to give full time to each branch.

Davies' Descriptive Geometry.—With application to Spherical Trigonometry, Spherical Projections, and Warped Surfaces.

Davies' Shades, Shadows, and Perspective.—A succinct exposition of the mathematical principles involved.

Davies' Science of Mathematics.—For teachers, embracing

I. GRAMMAR OF ARITHMETIC, III. LOGIC AND UTILITY OF MATHEMATICS,
II. OUTLINES OF MATHEMATICS, IV. MATHEMATICAL DICTIONARY.

KEYS MAY BE OBTAINED FROM THE PUBLISHERS

BY TEACHERS ONLY.

This book should be returned to the Library on or before the last date stamped below.

A fine of five cents a day is incurred by retaining it beyond the specified time.

Please return promptly.

All

NATIONAL SERIES.

"Histo

THE UN

Monteith
upon the
the younge

2. Willard's
Designed
seeds of vi

3. Willard's
course. N
a series of

4. Summary
inent facts
batim, use
of America

ENGLAN

people, wit
literature,

2. Summary
A series o
attachment
well-propor

ROME. R
and doubtf
charm to the subject.

GENERAL. Willard's Universal History. A vast subject so arranged and illustrated as to be less difficult to acquire or retain. Its whole substance, in fact, is summarized on one page, in a grand "Temple of Time, or Picture of Nations.

2 General Summary of History. Being the Summaries of American, and of English and French History, bound in one volume. The leading events in the histories of these three nations epitomized in the briefest manner.

A. S. BARNES & CO.

PUBLISHERS.